

PMATH351 — Real Analysis

CLASSNOTES FOR FALL 2018

by

Johnson Ng

BMath (Hons), Pure Mathematics major, Actuarial Science Minor

University of Waterloo

Table of Contents

<i>Table of Contents</i>	2
<i>List of Definitions</i>	7
<i>List of Theorems</i>	10
<i>List of Procedures</i>	15
1 Lecture 1 Sep 06th	17
1.1 Course Logistics	17
1.2 Preview into the Introduction	17
2 Lecture 2 Sep 10th	19
2.1 Basic Set Theory	19
2.2 Products of Sets	23
3 Lecture 3 Sep 12th	27
3.1 Axiom of Choice	27
3.2 Relations	29
4 Lecture 4 Sep 14th	35
4.1 Zorn's Lemma	35
4.2 Cardinality	38
4.2.1 Equivalence Relation	38
5 Lecture 5 Sep 17th	41
5.1 Cardinality (Continued)	41
6 Lecture 6 Sep 19th	47

6.1	Cardinality (Continued 2)	47
7	Lecture 7 Sep 21st	53
7.1	Cardinality (Continued 3)	53
7.1.1	Cardinal Arithmetic	55
8	Lecture 8 Sep 24th	57
8.1	Cardinality (Continued 4)	57
8.1.1	Cardinal Arithmetic (Continued)	57
9	Lecture 9 Sep 26th	61
9.1	Introduction to Metric Spaces	61
10	Lecture 10 Sep 28th	67
10.1	Introduction to Metric Spaces (Continued)	67
11	Lecture 11 Oct 01st	75
11.1	Introduction to Metric Spaces (Continued 2)	75
12	Lecture 12 Oct 03rd	81
12.1	Introduction to Metric Spaces (Continued 3)	81
12.2	Topology on Metric Spaces	82
13	Lecture 13 Oct 05th	87
13.1	Topology on Metric Spaces (Continued)	87
14	Lecture 14 Oct 12th	91
14.1	Topology on Metric Spaces (Continued 2)	91
15	Lecture 15 Oct 15th	97
15.1	Topology on Metric Spaces (Continued 3)	97
15.2	Convergences of Sequences	100
16	Lecture 16 Oct 17th	103
16.1	Convergences of Sequences (Continued)	103
17	Lecture 17 Oct 19th	107
17.1	Induced Metric and Topologies	107
17.2	Continuity on Metric Spaces	108
18	Lecture 18 Oct 22nd	111

18.1	Continuity on Metric Spaces (Continued)	111
19	Lecture 19 Oct 24th	115
19.1	Completeness of Metric Spaces	115
19.1.1	Basic Properties of Cauchy Sequences	116
19.1.2	Examples of Complete Spaces	117
19.1.2.1	Completeness of \mathbb{R}	117
20	Lecture 20 Oct 26th	119
20.1	Completeness of Metric Spaces (Continued)	119
20.1.1	Examples of Complete Spaces (Continued)	119
20.1.1.1	Completeness of ℓ_p	119
20.1.1.2	Completeness of $(C_b(X), \ \cdot\ _\infty)$	120
21	Lecture 21 Oct 31st	123
21.1	Completeness of Metric Spaces (Continued 2)	123
21.1.1	Examples of Complete Spaces (Continued 2)	123
21.1.1.1	Completeness of $(C_b(X), \ \cdot\ _\infty)$ (Continued)	123
21.1.2	Characterizations of Completeness	124
22	Lecture 22 Nov 02nd	127
22.1	Completeness of Metric Spaces (Continued 3)	127
22.1.1	Characterizations of Completeness (Continued)	127
23	Lecture 23 Nov 05th	131
23.1	Completeness of Metric Spaces (Continued 4)	131
23.1.1	Characterizations of Completeness (Continued 2)	131
24	Lecture 24 Nov 07th	135
24.1	Completions of Metric Spaces	135
24.2	Banach Contractive Mapping Theorem	138
25	Lecture 25 Nov 09th	141
25.1	Banach Contractive Mapping Theorem (Continued)	141
25.2	Baire Category Theorem	145
26	Lecture 26 Nov 12th	147
26.1	Baire Category Theorem (Continued)	147
27	Lecture 27 Nov 14th	155

27.1	Baire Category Theorem (Continued 2)	155
27.2	Compactness	157
28	Lecture 28 Nov 16th	161
28.1	Compactness (Continued)	161
29	Lecture 29 Nov 19th	169
29.1	Compactness (Continued 2)	169
30	Lecture 30 Nov 21st	175
30.1	Compactness (Continued 3)	175
30.2	Finite Dimensional Normed Linear Spaces	175
31	Lecture 31 Nov 23rd	181
31.1	Finite Dimensional Normed Linear Space (Continued)	181
31.2	Uniform Continuity	182
31.3	The Space $(C(X), \ \cdot\ _\infty)$	184
31.3.1	Weierstrass Approximation Theorem	184
32	Lecture 32 Nov 26th	187
32.1	The Space $(C(X), \ \cdot\ _\infty)$ (Continued)	187
32.1.1	Weierstrass Approximation Theorem (Continued)	187
33	Lecture 33 Nov 28th	193
33.1	The Space $(C(X), \ \cdot\ _\infty)$ (Continued 2)	193
33.1.1	Weierstrass Approximation Theorem (Continued 2)	193
33.1.2	Stone-Weierstrass Theorem	193
33.1.2.1	Lattice Version	194
33.1.2.2	Subalgebra Version	197
34	Lecture 34 Nov 30th	201
34.1	The Space $(C(X), \ \cdot\ _\infty)$ (Continued 3)	201
34.1.1	Stone-Weierstrass Theorem (Continued)	201
34.1.1.1	Subalgebra Version (Continued)	201
35	Lecture 35 Dec 03rd	207
35.1	The Space $(C(X), \ \cdot\ _\infty)$ (Continued 4)	207
35.1.1	Compactness in $C(X)$ and the Ascoli-Arzelà Theorem	207
A	Useful Theorems from Earlier Calculus	213

6 TABLE OF CONTENTS

B Assignment 1	215
C Assignment 2	219
D Assignment 3	223
Bibliography	233
Index	234

List of Definitions


























1	Definition (Universal Set)	19
2	Definition (Union)	20
3	Definition (Intersection)	20
4	Definition (Set Difference)	20
5	Definition (Symmetric Difference)	20
6	Definition (Set Complement)	21
7	Definition (Empty Set)	21
8	Definition (Power Set)	21
9	Definition (Product of Sets)	23
10	Definition (Relation)	24
11	Definition (Function)	24
12	Definition (Choice Function)	25
13	Definition (Relations)	29
14	Definition (Partially Ordered Sets)	30
15	Definition (Totally Ordered Sets / Chains)	32
16	Definition (Bounds)	32
17	Definition (Maximal Element)	35
18	Definition (Well-Ordered)	36
19	Definition (Equivalence Relation)	38
20	Definition (Equivalence Class)	38
21	Definition (Partition)	39
22	Definition (Finite Sets)	41
23	Definition (Cardinality)	41
24	Definition (Infinite Sets)	42
25	Definition (Countable)	44
26	Definition (Smaller Cardinality)	44

8 LIST OF DEFINITIONS






























27	Definition (Uncountable)	49
28	Definition (Sum of Cardinals)	55
29	Definition (Multiplication of Cardinals)	56
30	Definition (Exponentiation of Cardinals)	57
31	Definition (Metric & Metric Space)	61
32	Definition (Norm & Normed Linear Space)	63
33	Definition ($\ \cdot\ _p$ -norm)	67
34	Definition (Open & Closed)	82
35	Definition (Topology)	84
36	Definition (Closure)	91
37	Definition (Interior)	91
38	Definition (Neighbourhood)	91
39	Definition (Boundary Point)	92
40	Definition (Separable)	94
41	Definition (Dense)	95
42	Definition (Limit Points)	97
43	Definition (Convergence)	101
44	Definition (Induced Metric & Induced Topology)	107
45	Definition (Continuity)	108
46	Definition (Continuity on a Space)	111
47	Definition (Homeomorphism)	113
48	Definition (Equivalent Metric Spaces)	113
49	Definition (Continuity on a set)	114
50	Definition (Cauchy)	115
51	Definition (Complete Metric Spaces)	116
52	Definition (Boundedness)	117
53	Definition (Convergence of Functions)	120
54	Definition (Diameter of a Set)	125
55	Definition (Formal Sum)	129

56	Definition (Isometry)	135
57	Definition (Completion)	135
58	Definition (Uniformly Continuous Functions)	136
59	Definition (Fixed Point)	139
60	Definition (Lipschitz)	141
61	Definition (Contraction)	141
62	Definition (Points of Discontinuity)	145
63	Definition (F_σ Sets)	147
64	Definition (G_δ Sets)	147
65	Definition (Nowhere Dense)	148
66	Definition (First Category)	148
67	Definition (Second Category)	148
68	Definition (Residual)	148
69	Definition (Uniformly Convergent Sequence of Functions on a Point)	153
70	Definition (Cover)	158
71	Definition (Compact)	158
72	Definition (Sequential Compactness)	162
73	Definition (Bolzano-Weierstrass Property (BWP))	162
74	Definition (Finite Intersection Property (FIP))	163
75	Definition (ϵ -net)	166
76	Definition (Totally Bounded)	166
77	Definition (Bounded Linear Map)	175
78	Definition (Point-Separating)	194
79	Definition (Lattice)	194
80	Definition (Subalgebra)	197
81	Definition (Equicontinuity)	207
82	Definition (Pointwise Bounded Functions)	209
83	Definition (Relatively Compact Sets)	209


List of Theorems

1	 Theorem (De Morgan's Laws)	22
2	 Axiom (Zermelo's Axiom of Choice)	27
3	 Axiom (Zermelo's Axiom of Choice v2)	27
4	 Axiom (Zorn's Lemma)	36
5	 Theorem (★ Non-Zero Vector Spaces has a Basis)	36
6	 Axiom (Well-Ordering Principle)	37
7	 Theorem (Axioms of Choice and Its Equivalents)	37
8	 Proposition (Characterization of An Equivalence Relation)	39
9	 Theorem (Pigeonhole Principle)	41
10	 Corollary (Pigeonhole Principle (Finite Case))	42
11	 Proposition (\mathbb{N} is the Smallest Infinite Set)	43
12	 Corollary (Infinite Sets are Equivalent to Its Proper Subsets)	43
13	 Proposition (Injectivity is Surjectivity Reversed)	44
14	 Theorem (★ ★ ★ Cantor-Schröder-Bernstein Theorem (CSB))	47
15	 Corollary (Cantor-Schröder-Bernstein Theorem - Restated)	48
16	 Proposition (Denumerability Check)	49
17	 Theorem (Cantor's Diagonal Argument)	49
18	 Corollary (Uncountability of \mathbb{R})	50
19	 Theorem (★ Comparability of Cardinals)	53
20	 Theorem (Sums of Cardinals)	55
21	 Theorem (Multiplication of Cardinals)	56
22	 Theorem (Exponentiation of Cardinals)	58
23	 Theorem ($2^{\aleph_0} = c$)	58
24	 Theorem (Russell's Paradox)	59
25	 Axiom (Continuum Hypothesis)	60

26	🛡️ Axiom (Generalized Continuum Hypothesis)	60
27	🌲 Lemma (Young’s Inequality)	67
28	📄 Theorem (Hölder’s Inequality)	68
29	📄 Theorem (Minkowski’s Inequality)	69
30	📄 Theorem (Hölder’s Inequality v2)	77
31	📄 Theorem (Minkowski’s Inequality v2)	78
32	💧 Proposition (Properties of Open Sets)	82
33	👉 Corollary (Properties of Closed Sets)	83
34	📄 Theorem (Open Balls are Open)	87
35	💧 Proposition (Closed Sets Include Its Boundary Points)	92
36	💧 Proposition (Closures include the Boundary Points of a Set)	92
37	💧 Proposition (Closed Sets Include Its Limit Points)	98
38	💧 Proposition (Mixing the notions)	98
39	💧 Proposition (More on Closures and Interiors)	99
40	📄 Theorem (Uniqueness of Limits of Sequences)	101
41	📄 Theorem (Sequential Characterizations of Limit Points, Boundaries, and Closedness)	104
42	📄 Theorem (The Metric Topology of a Subset is Its Induced Topology)	108
43	📄 Theorem (Continuity and Neighbourhoods)	109
44	📄 Theorem (★ Sequential Characterization of Continuity)	110
45	📄 Theorem (Analogue of Sequential Characterization of Continuity on a Space, and Continuity and Neighbourhoods)	112
46	📄 Theorem (Convergent Sequences are Cauchy)	115
47	📄 Theorem (★★★ Convergent Cauchy Subsequences)	116
48	💧 Proposition (Cauchy Sequences are Bounded)	117
49	📄 Theorem (Bolzano-Weierstrass)	117
50	📄 Theorem (\mathbb{R} is complete)	118
51	📄 Theorem (★ Completeness of ℓ_p)	119
52	📄 Theorem (★★★ Uniformly Convergent Pointwise Continuous Functions have a Pointwise Continuous Limit)	121

53	 Theorem (★ ★ ★ Completeness for $C_b(X)$)	123
54	 Theorem (Nested Interval Theorem)	124
55	 Proposition (Diameters of Subsets)	125
56	 Theorem (Cantor's Intersection Principle)	127
57	 Theorem (★ ★ Weierstrass M-test)	129
58	 Proposition (Subsets of Complete Spaces are Complete if they are Closed)	135
59	 Theorem (Completion Theorem)	137
60	 Theorem (Banach Contractive Mapping Theorem)	142
61	 Theorem (Set of Points of Discontinuity is F_σ)	149
62	 Theorem (Baire Category Theorem I)	149
63	 Theorem (★ Baire Category Theorem II)	151
64	 Corollary (\mathbb{Q} is not G_δ)	152
65	 Corollary (There are no Functions Discontinuous on all Irrational Numbers)	152
66	 Theorem (Limit of Sequence of Continuous Functions that Converges Pointwise is Continuous)	153
67	 Theorem (Uniform Convergence of A Sequence of Continuous Functions that Converges Pointwise)	155
68	 Corollary (Continuity of the Limit of a Sequence of Pointwise Convergent Functions on a Residual Set)	156
69	 Corollary (Derivative of a Function is Continuous on a dense G_δ set in \mathbb{R})	157
70	 Theorem (Heine-Borel Theorem)	158
71	 Proposition (Compact Spaces are Closed and Bounded)	159
72	 Proposition (Closed Subsets of Compact Sets are Compact)	161
73	 Theorem (Sequential Compactness is Equivalent to BWP)	163
74	 Theorem (FIP and Compactness)	164
75	 Corollary (Generalized Nested Interval Theorem for Compact Metric Spaces)	165
76	 Corollary (Compact Metric Spaces are Complete)	165
77	 Theorem (Compact Sets are Totally Bounded)	166
78	 Proposition (A Set is Totally Bounded iff Its Closure is Totally Bounded)	167
79	 Theorem (Compact Sets have BWP)	169
80	 Proposition (Sequential Compactness \implies Completeness and Total Boundedness)	170
81	 Theorem (Continuity Preserves Sequential Compactness)	170

82	👉 Corollary (Extreme Value Theorem)	171
83	☕ Theorem (Lesbesgue)	171
84	☕ Theorem (Lesbesgue-Borel)	172
85	☕ Theorem (Compactness \iff Completeness + Totally Bounded)	173
86	☕ Theorem (Continuity Preserves Compactness)	175
87	☕ Theorem (Boundedness is Equivalent to Continuity in Finite Dimensional Normed Linear Spaces) 176	
88	🌲 Lemma (Continuity of the Norm)	176
89	💧 Proposition (Linear Map Between Spaces of Different Dimensions is Bounded)	176
90	☕ Theorem (Boundedness of Functions between n -dimensional Vector Spaces and n -dimensional Normed Linear Spaces)	177
91	☕ Theorem (The Basis of a Infinite Dimensional Banach Spaces is Uncountable)	178
92	☕ Theorem (All Linear Maps Between Finite Dimensional Normed Linear Spaces are Bounded) 178	
93	👉 Corollary (All Linear Maps from A Finite Dimensional Normed Linear Space to Any Normed Linear Space is Bounded)	179
94	☕ Theorem (Completeness of Finite Dimensional Normed Linear Spaces)	182
95	☕ Theorem (Sequential Characterization of Uniform Continuity)	182
96	☕ Theorem (Continuous Functions from a Compact Set Is Uniformly Continuous)	183
97	☕ Theorem (Continuous Bijections from a Compact Space is a Homeomorphism)	184
98	🌲 Lemma (Lemma for Weierstrass Approximation)	187
99	☕ Theorem (★ ★ ★ Weierstrass Approximation Theorem)	188
100	💧 Proposition (Moments)	189
101	☕ Theorem (Banach-Mazurkiewickz Theorem)	190
102	👉 Corollary (Separability of $(C[a, b], \ \cdot\ _\infty)$)	193
103	💧 Proposition ($C(X)$ is Point-Separating)	194
104	☕ Theorem (Stone-Weierstrass Theorem — Lattice Version)	196
105	🌲 Lemma (Closure of a Subalgebra is a Subalgebra)	198
106	☕ Theorem (Stone-Weierstrass Theorem — Subalgebra Version)	201
107	☕ Theorem (Stone-Weierstrass Theorem — Complex Version)	204
108	💧 Proposition (Equicontinuity in a Compact Set is Uniform)	208
109	💧 Proposition (Pointwise Bounded Equicontinuous Functions in a Compact Set are Uniformly Bounded)	209
110	☕ Theorem (Arzelà-Ascoli)	210

A.1  Theorem (Monotone Convergence Theorem) 213

List of Procedures

1 Lecture 1 Sep 06th

1.1 Course Logistics

No content is covered in today's lecture so this chapter will cover some of the important logistical highlights that were mentioned in class.

- Assignments are designed to help students understand the content.
- Due to shortage of manpower, not all assignment questions will be graded; however, students are encouraged to attempt all of the questions.
- To further motivate students to work on ungraded questions, the midterm and final exam will likely recycle some of the assignment questions.
- There are no required text, but the professor has prepared course notes for reading. The course notes are self-contained.
- The approach of the class will be more interactive than most math courses.
- Due to the size of the class, students are encouraged to utilize Waterloo Learn for questions, so that similar questions by multiple students can be addressed at the same time.

1.2 Preview into the Introduction

How do we compare the size of two sets?

- If the sets are finite, this is a relatively easy task.
- If the sets are infinite, we will have to rely on functions.
 - Injective functions tell us that the **domain is of size that is lesser than or equal to the codomain**.
 - Surjective functions tell us that the **codomain is of size that is lesser than or equal to the domain**.
 - So does a bijective function tell us that the domain and codomain have the same size? Yes, although this is not as intuitive as it looks, as it relies on **Cantor-Schröder-Bernstein Theorem**.

Now, given two arbitrary sets, are we guaranteed to always be able to compare their sizes? It is tempting to immediately say yes, but to do that, one would have to agree on the **Axiom of Choice**. Fortunately, within the realm of this course, the Axiom of Choice is taken for granted.

2 Lecture 2 Sep 10th

2.1 Basic Set Theory

We shall use the following notations for some of the common set of numbers that we are already familiar with:

- \mathbb{N} denotes the set of natural numbers $\{1, 2, 3, \dots\}$;
- \mathbb{Z} denotes the set of integers $\{\dots, -2, -1, 0, 1, 2, \dots\}$;
- \mathbb{Q} denotes the set of rational numbers $\{\frac{a}{b} \mid a \in \mathbb{Z}, b \in \mathbb{N}\}$; and
- \mathbb{R} denotes the set of real numbers.

We shall start with having certain basic properties of \mathbb{N} , \mathbb{Z} , and \mathbb{Q} .

WE WILL USE the notation $A \subset B$ and $A \subseteq B$ interchangeably to mean that A is a subset of B with the possibility that $A = B$. When we wish to explicitly emphasize this possibility, we shall use $A \subseteq B$. When we wish to explicitly state that A is a **proper subset** of B , we will either specify that $A \neq B$ or simply $A \subsetneq B$.

Definition 1 (Universal Set)

A universal set, which we shall generally give the label X , is a set that contains all the mathematical objects that we are interested in.

This is a hand-wavy definition, but it is not in the interest of this course to further explore on this topic.

With a universal set in place, we can have the following defini-

tions:

Definition 2 (Union)

Let X be a set. If $\{A_\alpha\}_{\alpha \in I}$ such that $A_\alpha \subset X$, then the **union** for all A_α is defined as

$$\bigcup_{\alpha \in I} A_\alpha := \{x \in X \mid \exists \alpha \in I, x \in A_\alpha\}.$$

Definition 3 (Intersection)

Let X be a set. If $\{A_\alpha\}_{\alpha \in I}$ such that $A_\alpha \subset X$, then the **intersection** for all A_α is defined as

$$\bigcap_{\alpha \in I} A_\alpha := \{x \in X \mid \forall \alpha \in I, x \in A_\alpha\}.$$

Definition 4 (Set Difference)

Let X be a set and $A, B \subseteq X$. The **set difference** of A from B is defined as

$$A \setminus B := \{x \in X \mid x \in A, x \notin B\}.$$

On a similar notion:

Definition 5 (Symmetric Difference)

Let X be a set and $A, B \subseteq X$. The **symmetric difference** of A and B is defined as

$$A \Delta B := \{x \in X \mid (x \in A \wedge x \notin B) \vee (x \notin A \wedge x \in B)\}.$$

We can also talk about the non-members of a set:

In words, for an element in the symmetric difference of two sets, the element is either in A or B but not both. We can also think of the symmetric difference as

$$(A \cup B) \setminus (A \cap B)$$

or

$$(A \setminus B) \cup (B \setminus A).$$

Definition 6 (Set Complement)

Let X be a set and $A \subset X$. The set of all non-members of A is called the *complement* of A , which we denote as

$$A^c := \{x \in X \mid x \notin A\}.$$

Note 2.1.1

Note that

$$(A^c)^c = \{x \in X \mid x \notin A^c\} = \{x \in X \mid x \in A\} = A.$$

Now taking a step away from that, we define the following:

Definition 7 (Empty Set)

An *empty set*, denoted by \emptyset , is a set that contains nothing.

Note 2.1.2

The empty set is set to be a subset of all sets.

Definition 8 (Power Set)

Let X be a set. The *power set* of X is the set that contains all subsets of X , i.e.

$$\mathcal{P}(X) := \{A \mid A \subset X\}.$$

Note 2.1.3


A power set is always non-empty, since $\emptyset \in \mathcal{P}(\emptyset)$, and since $\emptyset \subset X$ for any set X , we have $\emptyset \in \mathcal{P}(X)$.

Example 2.1.1

Let $X = \{1, 2, \dots, n\}$. There are several ways we can show that the size of $\mathcal{P}(X)$ is 2^n . One of the methods is by using a characteristic function that maps from A to $\{0, 1\}$, defined by

$$X_A : A \rightarrow \{0, 1\}$$

$$X_A(x) = \begin{cases} 1 & x \in A \\ 0 & x \notin A \end{cases}.$$

Using this function, each element in X has 2 states: one being in the subset, and the other being not in the subset, which are represented by 1 and 0 respectively. It is then clear that there are 2^n of such configurations. 

 Theorem 1 (De Morgan's Laws)

Let X be a set. Given $\{A_\alpha\}_{\alpha \in I} \subset \mathcal{P}(X)$, we have

1. $\left(\bigcup_{\alpha \in I} A_\alpha\right)^c = \bigcap_{\alpha \in I} A_\alpha^c$; and
2. $\left(\bigcap_{\alpha \in I} A_\alpha\right)^c = \bigcup_{\alpha \in I} A_\alpha^c$.

 Proof

1. Note that

$$\begin{aligned} x \in \left(\bigcup_{\alpha \in I} A_\alpha\right)^c &\iff \nexists \alpha \in I \ x \in A_\alpha \\ &\iff \forall \alpha \in I \ x \notin A_\alpha \\ &\iff \forall \alpha \in I \ x \in A_\alpha^c \text{ by set complementation} \\ &\iff x \in \bigcap_{\alpha \in I} A_\alpha^c. \end{aligned}$$


2. Observe that, by part 1,

$$\left(\bigcap_{\alpha \in I} A_\alpha\right)^c = \left(\left(\bigcup_{\alpha \in I} A_\alpha^c\right)^c\right)^c = \bigcup_{\alpha \in I} A_\alpha^c.$$

Example 2.1.2

Suppose $I = \emptyset$. Then what is $\bigcup_{\alpha \in \emptyset} A_\alpha$? It is sensible to think that all we are left with is simply a union of empty sets, and so

$$\bigcup_{\alpha \in \emptyset} A_\alpha = \emptyset. \quad (2.1)$$

And what about $\bigcap_{\alpha \in \emptyset} A_\alpha$? By  Theorem 1, it is quite clear from Equation (2.1) that

$$\bigcap_{\alpha \in \emptyset} A_\alpha = X. \quad \rightarrow$$

2.2 Products of Sets

Definition 9 (Product of Sets)

Given 2 sets X and Y , the **product** of X and Y is given by

$$X \times Y := \{(x, y) \mid x \in X, y \in Y\}.$$

We often refer to elements of $X \times Y$ as **tuples**.

Note 2.2.1

Now if

$$X = \{x_1, x_2, \dots, x_n\},$$

$$Y = \{y_1, y_2, \dots, y_m\},$$

then

$$X \times Y = \{(x_i, y_j) \mid i = 1, 2, \dots, n, j = 1, 2, \dots, m\}$$

and so the size of $X \times Y$ is mn .

Consequently, we can think of tuples as two elements being in some "relation".

Definition 10 (Relation)

A *relation* on sets X and Y is a subset R of the product $X \times Y$. We write

$$xRy \text{ if } (x, y) \in R \subset X \times Y.$$

We call

- $\{x \in X \mid \exists y \in Y, (x, y) \in R\}$ as the **domain** of R ; and
- $\{y \in Y \mid \exists x \in X, (x, y) \in R\}$ as the **range** of R .

In relation to that, functions are, essentially, relations.

Definition 11 (Function)

A *function* from X to Y is a relation R such that

$$\forall x \in X \exists! y \in Y (x, y) \in R.$$

SUPPOSE X_1, X_2, \dots, X_n are non-empty¹ sets. We can define

$$X_1 \times X_2 \times \dots \times X_n = \prod_{i=1}^n X_i := \{(x_1, x_2, \dots, x_n) \mid x_i \in X_i\}.$$

Now if $X_i = X_j = X$ for all $i, j = 1, 2, \dots, n$, we write

$$\prod_{i=1}^n X_i = \prod_{i=1}^n X = X^n.$$

¹ We are typically only interested in non-empty sets, since empty sets usually lead us to vacuous truths, which are not interesting.

AND NOW COMES THE PROBLEM: given a collection $\{X_\alpha\}_{\alpha \in I}$ of non-empty sets², what do we mean by

² i.e. we now talk about arbitrary $\alpha \in I$.

$$\prod_{\alpha \in I} X_\alpha?$$

To motivate for what comes next, consider

$$\prod_{i=1}^n X_i = X_1 \times \dots \times X_n = \{(x_1, \dots, x_n) \mid x_i \in X_i\}.$$

Choose $(x_1, \dots, x_n) \in \prod_{i=1}^n X_i$. This induces a function

$$f_{(x_1, \dots, x_n)} : \{1, \dots, n\} \rightarrow \bigcup_{i=1}^n X_i$$

with

$$\begin{aligned} f(1) &= x_1 \in X_1 \\ f(2) &= x_2 \in X_2 \\ &\vdots \\ f(n) &= x_n \in X_n \end{aligned}$$

Now assume for a more general f such that

$$f : \{1, \dots, n\} \rightarrow \bigcup_{i=1}^n X_i$$


is defined by

$$f(i) \in X_i.$$

Then, we have

$$(f(1), f(2), \dots, f(n)) \in \prod_{i=1}^n X_i,$$

which leads us to the following notion:

 **Definition 12 (Choice Function)**

Given a collection $\{X_\alpha\}_{\alpha \in I}$ of non-empty sets, let

$$\prod_{\alpha \in I} X_\alpha = \left\{ f : I \rightarrow \bigcup_{\alpha \in I} X_\alpha \right\}$$

such that $f(\alpha) \in X_\alpha$. Such an f is called a *choice function*.

And so we may ask a similar question as before: if each X_α is non-empty, is $\prod_{\alpha \in I} X_\alpha$ non-empty? Turns out this is not as easy to show. In fact, it is essentially impossible to show, because this is exactly the **Axiom of Choice**.

3 Lecture 3 Sep 12th

3.1 Axiom of Choice

RECALL our final question of last lecture: If $\{X_\alpha\}_{\alpha \in I}$ is a non-empty collection of non-empty sets, is

$$\prod_{\alpha \in I} X_\alpha \neq \emptyset ?$$

Turns out this is widely known (in the world of mathematics) as the **Axiom of Choice**.

♣ Axiom 2 (Zermelo's Axiom of Choice)

If $\{X_\alpha\}_{\alpha \in I}$ is a non-empty collection of non-empty sets, then

$$\prod_{\alpha \in I} X_\alpha \neq \emptyset.$$

An equivalent statement of the above axiom is:

♣ Axiom 3 (Zermelo's Axiom of Choice v2)

$X \neq \emptyset \implies$

$$\exists f : \mathcal{P}(X) \setminus \{\emptyset\} \rightarrow X \quad \forall A \in \mathcal{P}(X) \setminus \{\emptyset\} \quad f(A) \in A$$

where f is the choice function.

Exercise 3.1.1

Prove that * Axiom 2 and * Axiom 3 are equivalent.

 **Proof****From * Axiom 2 to * Axiom 3:**

Since $X \neq \emptyset$, we have that $\mathcal{P}(X) \setminus \{\emptyset\}$ is a non-empty collection of non-empty sets. Therefore,

$$\prod_{A \in \mathcal{P}(X) \setminus \{\emptyset\}} A \neq \emptyset.$$

So we know that

$$\exists (x_A)_{A \in \mathcal{P}(X) \setminus \{\emptyset\}} \in \prod_{A \in \mathcal{P}(X) \setminus \{\emptyset\}} A.$$

We then simply need to choose the choice function $f : \mathcal{P}(X) \setminus \{\emptyset\} \rightarrow X$ such that

$$f(A) = x_A \in A.$$

From * Axiom 3 to * Axiom 2:

Let $X_\alpha \in \mathcal{P}(X)$ for $\alpha \in I$, where I is some index set. We know that not all $X_\alpha = \emptyset$ since $X \neq \emptyset$. Choose $J \subseteq I$ such that $\{X_\alpha\}_{\alpha \in J}$ is a non-empty collection of non-empty sets. Let $f : \mathcal{P}(X) \setminus \{\emptyset\}$ be any choice function. By * Axiom 3,


$$\forall X_\alpha \in \mathcal{P}(X) \setminus \{\emptyset\} \quad f(X_\alpha) \in X_\alpha.$$

Therefore,

$$(f(X_\alpha))_{\alpha \in J} \in \prod_{\alpha \in J} X_\alpha.$$

3.2 Relations

Now, it is in our interest to start talking about comparisons or relations between the mathematical objects that we have defined.

 **Definition 13 (Relations)**

A relation R on a set X is ¹


- **(Reflexive)** $\forall x \in X \ xRx$;
- **(Symmetric)** $\forall x, y \in X \ xRy \iff yRx$;
- **(Anti-symmetric)** $\forall x, y \in X \ xRy \wedge yRx \implies x = y$;
- **(Transitive)** $\forall x, y, z \in X \ xRy \wedge yRz \implies xRz$.

¹ We can look at this definition as $R \subseteq X \times X$. Under such a definition, we would have

- **(Reflexive)** $\forall x \in X \ (x, x) \in R$;
- **(Symmetric)** $\forall x, y \in X \ (x, y) \in R \iff (y, x) \in R$;
- **(Anti-symmetric)** $\forall x, y \in X \ (x, y), (y, x) \in R \implies x = y$;
- **(Transitive)** $\forall x, y, z \in X \ (x, y), (y, z) \in R \implies (x, z) \in R$.


Example 3.2.1

Let $X = \mathbb{R}$, and let $xRy \iff x \leq y$, where \leq is the notion of “less than or equal to”, which we shall assume that it has the meaning that we know. Observe that \leq is:

- reflexive: $\forall x \in \mathbb{R} \ x \leq x$ is true;
- anti-symmetric: $\forall x, y \in \mathbb{R} \ x \leq y \wedge y \leq x \implies x = y$; and
- transitive: $\forall x, y, z \in \mathbb{R} \ x \leq y \wedge y \leq z \implies x \leq z$. 


Example 3.2.2

Let $Y \neq \emptyset$, $X = \mathcal{P}(Y)$, with $ARB \iff A \subseteq B$. Observe that \subseteq is:

- reflexive: $\forall A \in \mathcal{P}(Y) \ ARA \iff A \subseteq A$ is true;
- anti-symmetric: $\forall A, B \in \mathcal{P}(Y) \ ARB \wedge BRA \iff A \subseteq B \wedge B \subseteq A \implies A = B$;
- transitive: $\forall A, B, C \in \mathcal{P}(Y) \ ARB \wedge BRC \iff A \subseteq B \wedge B \subseteq C \implies A \subseteq C$. 

Example 3.2.3

Let $Y \neq \emptyset$, $X = \mathcal{P}(Y)$, with $ARB \iff A \supseteq B$. Observe that \supseteq is:

- reflexive: $\forall A \in \mathcal{P}(Y) \ ARA \iff A \subseteq A$;
- anti-symmetric: $\forall A, B \in \mathcal{P}(Y) \ ARB \wedge BRA \iff A \supseteq B \wedge B \supseteq A \implies A = B$;
- transitive: $\forall A, B, C \in \mathcal{P}(Y) \ ARB \wedge BRC \iff A \supseteq B \wedge B \supseteq C \implies A \supseteq C$. 

All the above examples are also known as *partially ordered sets*.

Definition 14 (Partially Ordered Sets)

The set X with the relation R on X is called a *partially ordered set* (or a *poset*) if R is

- reflexive;
- anti-symmetric; and
- transitive.

We denote a poset by (X, R) .

The “partial” in ‘partially ordered’ indicates that not every pair of elements need to be comparable, i.e. there may be pairs for which neither precedes the other (anti-symmetry).

Note 3.2.1

If (X, R) is a poset, then if $A \subseteq X$, and $R_1 = R \upharpoonright_{A \times A}$, then (A, R_1) is also a poset.

Example 3.2.4


How many possible relations can we define on these sets to make them into posets?

1. $X = \emptyset$

Solution


We have that $R = \emptyset \times \emptyset$, and so the only relation we have is an empty relation. Then it is vacuously true that (X, R) a poset.

2. $X = \{x\}$

 **Solution**

We have that $R = X \times X = \{(x, x)\}$. It is clear that (X, R) is a poset.


3. $X = \{x, y\}$

 **Solution**


There are 3 possible relations:

- a relation where xRx and yRy ;
- a relation where xRy ; or
- a relation where yRx .


3 possibilities represented as graphs (known as Hasse diagram), separated by lines:



4. $X = \{x, y, z\}$

 **Solution**

The following are all the possibilities represented by graphs, where the underlined numbers represent the number of ways we can rearrange the elements for unique relations:




Therefore, we see that there are a total of

$$1 + 3 + 3 + 6 + 6 = 19 \text{ relations.}$$

Exercise 3.2.1

How many possible relations can we define on a set of 6 elements to the set into a poset?

 **Solution**

to be added

Definition 15 (Totally Ordered Sets / Chains)

The set X with the relation R on X is called a **totally ordered set** (or a **chain**) if (X, R) is a poset with the exception that, for any $x, y \in X$, either xRy or yRx but not both.

Definition 16 (Bounds)

Let (X, \leq) be a poset. Let $A \subset X$. We say $x_0 \in X$ is an **upper bound** for A if

$$\forall a \in A \quad a \leq x_0.$$

If A has an upper bound, we say that A is **bounded above**. If A is bounded above, then x_0 is the **least upper bound** (or **supremum**) of A if for any $x_1 \in X$ that is an upper bound of A , we have

$$x_0 \leq x_1.$$

We write $x_0 = \text{lub}(A) = \text{sup}(A)$. If $\text{sup}(A) \in A$, then $\text{sup}(A) = \text{max}(A)$ is the **maximum** of A .

We can analogously define for:

upper bound \rightarrow lower bound

bounded above \rightarrow bounded below

least upper bound, lub \rightarrow greatest lower bound, glb


supremum, sup \rightarrow infimum, inf

maximum, max \rightarrow minimum, min

Note 3.2.2

By **anti-symmetry** of posets, we have that max, sup, min, inf are all unique if they exist.

Example 3.2.5 (Least Upper Bound Property of \mathbb{R})

Let $X = \mathbb{R}$, and \leq be the order that we have defined. Every bounded non-empty subset of X has a supremum. 

Example 3.2.6

Let $Y \neq \emptyset$, and $X = \mathcal{P}(Y)$, and \subseteq the ordering by inclusion. We know that Y is the maximum element of (X, \subseteq) . Then the collection $\{A_\alpha\}_{\alpha \in I} \subset \mathcal{P}(Y)$ is bounded above by Y , and we have that


$$\sup(\{A_\alpha\}_{\alpha \in I}) = \bigcup_{\alpha \in I} A_\alpha$$

$$\inf(\{A_\alpha\}_{\alpha \in I}) = \bigcap_{\alpha \in I} A_\alpha$$

Now if $Y = \emptyset$, we would end up having

$$\sup(\{A_\alpha\}_{\alpha \in I}) = \emptyset$$

$$\inf(\{A_\alpha\}_{\alpha \in I}) = X$$

This makes sense, since the empty set would be the least of upper bounds, and since $X = \mathcal{P}(Y)$ would have to be the greatest of lower bounds. 

4 Lecture 4 Sep 14th

4.1 Zorn's Lemma





Definition 17 (Maximal Element)

Let (X, \leq) be a poset. An element $x \in X$ is *maximal* if whenever $y \in X$ is such that $x \leq y$, we must have $y = x$.

Example 4.1.1

Looking back at Example 3.2.4, on the set $X = \{x, y, z\}$, we have that the maximal element in each possible poset is/are:


$\bullet x \quad \bullet y \quad \bullet z$ x, y, z are all maximal



This shows to us that the maximal element does not have to be unique.




Example 4.1.2


- Given $X \neq \emptyset$, the maximal element of the poset $(\mathcal{P}(X), \subseteq)$ is X .
- Given $X \neq \emptyset$, the maximal element of the poset $(\mathcal{P}(X), \supseteq)$ is \emptyset .
- The poset (\mathbb{R}, \leq) has no maximal element. 

 Axiom 4 (Zorn's Lemma)

If (X, \leq) is a non-empty poset such that every chain $S \subset X$ has an upper bound, then (X, \leq) has a maximal element.

 Theorem 5 (★ Non-Zero Vector Spaces has a Basis)

Every non-zero vector space, V , has a basis.

 Proof (★)

Let


$$\mathcal{L} := \{A \subset V \mid A \text{ is linearly independent}\}.$$

Note that $\mathcal{L} \neq \emptyset$ since $V \neq \{0\}$. Now order elements of \mathcal{L} with \subseteq . It suffices to show that (\mathcal{L}, \subseteq) has a maximal element, since this maximal element must be a basis. Otherwise, we would contradict the maximality of such an element.¹

Now let $S = \{A_\alpha\}_{\alpha \in I}$ be a chain in \mathcal{L} . Let

$$A_0 = \bigcup_{\alpha \in I} A_\alpha.$$

Require clarification before proceeding... □

 Definition 18 (Well-Ordered)

We say that a poset (X, \leq) is *well-ordered* if every non-empty subset $A \subset X$ has a least/minimal element in A .

The flow of this proof is a typical approach when Zorn's Lemma is involved.


¹ This is the key to this proof.

Exercise 4.1.1

Prove that well-ordered sets are chains.

Example 4.1.3

(\mathbb{N}, \leq) is well-ordered. 

 Axiom 6 (Well-Ordering Principle)

Every non-empty set can be well-ordered.


 Theorem 7 (Axioms of Choice and Its Equivalents)

TFAE:

1. *Axiom of Choice*, * *Axiom 2*
2. *Zorn's Lemma*, * *Axiom 4*
3. *Well-Ordering Principle*, * *Axiom 6*.

Exercise 4.1.2

Prove  *Theorem 7*

 Proof

(3) \implies (1) is simple; let the choice function be such that we pick the minimal element from each set among a non-empty collection of non-empty sets. It is clear that the product of these sets will always have an element, in particular the tuple where each component is the minimal element of each set.

The rest will be added once I've worked it out □


Example 4.1.4

Let $X = \mathbb{Q}$. Let $\varphi : \mathbb{Q} \rightarrow \mathbb{N}$ be defined such that

$$\varphi\left(\frac{m}{n}\right) = \begin{cases} 2^m 5^n & m > 0 \\ 1 & m = 0 \\ 3^{-m} 7^n & m < 0 \end{cases}$$

By the **unique prime factorization of natural numbers** (or **Fundamental Theorem of Arithmetic**), we have that φ is injective. In fact,

$$r \leq s \iff \varphi(r) \leq \varphi(s),$$

showing to us that we have a well-ordering on \mathbb{Q} . 

4.2 Cardinality

4.2.1 Equivalence Relation

Definition 19 (Equivalence Relation)

Let X be non-empty set. A relation \sim on X is an **equivalence relation** if it is

- reflexive;
- symmetric; and
- transitive.

Definition 20 (Equivalence Class)

Let X be a non-empty set, and $x \in X$. An **equivalence class** of x under the equivalence relation \sim is defined as

$$[x] := \{y \in X \mid x \sim y\}.$$

Note 4.2.1

Note that we either have $[x] = [y]$ or $[x] \cap [y] = \emptyset$. This is sensible, since if $w \in [x]$, then $w \sim x$. If $w \in [y]$, then we are done. If $w \notin [y]$, suppose $\exists v \in [y]$ such that $w \sim v$, which then implies $w \in [y]$ which is a contradiction.

This results shows to us that

$$X = \bigcup_{x \in X} [x],$$

or in words, equivalence classes partition the set.

Definition 21 (Partition)

Let $X \neq \emptyset$. A *partition* of X is a collection $\{A_\alpha\}_{\alpha \in I} \subset \mathcal{P}(X)$ such that

1. $A_\alpha \neq \emptyset$;
2. $A_\alpha \cap A_\beta = \emptyset$ if $\alpha \neq \beta$ in I ; and
3. $X = \bigcup_{\alpha \in I} A_\alpha$.

With this, we have ourselves another method to show that \sim is an equivalence relation.

Proposition 8 (Characterization of An Equivalence Relation)

If $\{A_\alpha\}_{\alpha \in I}$ is a partition of X and $x \sim y \iff x, y \in A_\alpha$, then \sim is an equivalence relation.

The proof of this statement has been stated above.

Similar to when we defined partial orders, we can ask ourselves the following question:

Example 4.2.1

How many equivalence relations are there on the set $X = \{1, 2, 3\}$?²

Solution

Note that we can partition X as

$$\{\{1\}, \{2\}, \{3\}\}, \{\{1, 2, 3\}\},$$

and

$$\{\{1, 2\}, \{3\}\},$$

² By Proposition 8, this question is equivalent to asking for the number of partitions we can create from the set X . The study of counting partitions is what is covered by Bell's Number.

which we can rearrange in 3 different ways. Therefore, there are 5 different equivalence relations that we can define on X .

Example 4.2.2

Let X be any set. Consider $\mathcal{P}(X)$. Define \sim on $\mathcal{P}(X)$ by

$$A \sim B \iff \exists f : A \rightarrow B$$

such that f is surjective³. It is easy to verify that \sim is an equivalence relation.

³ \sim partitions X into sets that have the same number of elements.



5 Lecture 5 Sep 17th

5.1 Cardinality (Continued)

Definition 22 (Finite Sets)

A set X is *finite* if $X = \emptyset$ or if $X \sim \{1, 2, \dots, n\}$ for some $n \in \mathbb{N}$, where \sim is the equivalence relation defined in Example 4.2.2.

Definition 23 (Cardinality)

If $X \sim n$, we say X has *cardinality* n and write $|X| = n$. We also let $|\emptyset| = 0$.

NOW A GOOD QUESTION here is: if $n \neq m$, is $\{1, 2, \dots, n\} \sim \{1, 2, \dots, m\}$?

Theorem 9 (Pigeonhole Principle)

The set $\{1, 2, \dots, n\}$ is not equivalent to any of its proper subset.

Proof

We shall prove this by induction on n .

Base case: $\{1\} \approx \emptyset$.

This is a **proof by contradiction**, using the fact that we cannot find an injective function from a “larger” set to a “smaller” set.

We can assume that the function f is not surjective, since if the larger set is indeed equivalent to the smaller set, then it should not matter if f is surjective or not. In particular, we only require that there be an injective function.

Requires clarification and confirmation

Assume that the statement holds for $\{1, \dots, k\}$. Suppose we have an injective function

$$f : \{1, 2, \dots, k, k + 1\} \rightarrow \{1, 2, \dots, k, k + 1\}$$

that is not surjective.

Case 1: $k + 1 \notin \text{range}(f)$, where $\text{range}(f)$ is the range of f . Then we have

Note: \upharpoonright is the restriction sign.

$$f \upharpoonright_{\{1, \dots, k\}} : \{1, \dots, k\} \rightarrow \{1, \dots, k\} \setminus \{f(k + 1)\}.$$

However, f is an injective function and clearly

$$\{1, \dots, k\} \setminus \{f(k + 1)\} \subseteq \{1, \dots, k\},$$

a contradiction.

Case 2: $k + 1 \in \text{range}(f)$. Then $\exists j_0 \in \{1, \dots, k, k + 1\}$ such that $f(j_0) = k + 1$, and since f is not surjective, $\exists m \in \{1, \dots, k\}$ such that $m \notin \text{range}(f)$. Then consider a new function $g : \{1, \dots, k, k + 1\} \rightarrow \{1, \dots, k\}$ such that

$$g(a) = \begin{cases} m & a = k + 1 \\ f(k + 1) & a = j_0 \\ f(a) & a \neq j_0, k + 1 \end{cases} \quad \square$$


➡ Corollary 10 (Pigeonhole Principle (Finite Case))

If the set X is finite, then X is not equivalent to any proper subset.

Exercise 5.1.1

Prove ➡ Corollary 10.


Sketch of proof:




📖 Definition 24 (Infinite Sets)

X is *infinite* if it is not finite.

Example 5.1.1

Observe that we can construct a function $f : \mathbb{N} \rightarrow \{2, 3, \dots\}$ by $f(n) = n + 1$. It is clear that f is a bijective function, and so $\mathbb{N} \sim \{2, 3, \dots\}$. 

 **Proposition 11 (ℕ is the Smallest Infinite Set)**

Every infinite set contains a subset $A \sim \mathbb{N}$.

 **Proof**

Suppose X is infinite. Let

$$f : \mathcal{P}(X) \setminus \{\emptyset\} \rightarrow X$$

such that for $S \subset X$ where $S \neq \emptyset$, $f(S) \in S$ ¹. Let $x_1 = f(X)$. Let $x_2 = f(X \setminus \{x_1\})$. Recursively, define


¹* Axiom 3 ahoy!

$$x_n = f(X \setminus \{x_1, \dots, x_{n-1}\}).$$

This gives us a sequence

$$X \supset S = \{x_1, \dots, x_n, \dots\}$$

which is equivalent to \mathbb{N} via the map $n \mapsto x_n$.

 **Corollary 12 (Infinite Sets are Equivalent to Its Proper Subsets)**

Every infinite set X is equivalent to a proper subset of X .

 **Proof**

Given such an X , we construct a sequence $\{x_n\}$ as in the previous proof. Define $f : X \rightarrow X \setminus \{x_n\}$ by

$$f(x) = \begin{cases} x_{n+1} & x \in \{x_n\} \\ x & x \notin \{x_n\}. \end{cases}$$

Clearly so, f is injective.

Definition 25 (Countable)

We say that a set is *countable* (or *denumerable*) is either X is finite or if $X \sim \mathbb{N}$. If $X \sim \mathbb{N}$, we can say that X is *countably infinite* and write $|X| = |\mathbb{N}| = \aleph_0$.

Definition 26 (Smaller Cardinality)

Given 2 sets X, Y , we write

$$|X| \leq |Y|$$

if $\exists f : X \rightarrow Y$ injective.

Proposition 13 (Injectivity is Surjectivity Reversed)

TFAE

1. $\exists f : X \rightarrow Y$ injective
2. $\exists g : Y \rightarrow X$ surjective

Proof

(1) \implies (2): Define

$$g(y) = \begin{cases} x & \exists x \in X \text{ } f(x) = y \\ x_0 & \text{any } x_0 \in X \end{cases}$$

Clearly g is surjective.

(2) \implies (1): Since g is surjective, for each $x \in X$, we have that²

$$g^{-1}(\{x\}) = \{y \in Y : g(y) = x\} \neq \emptyset.$$

By the Axiom of Choice, there exists a choice function $h : \mathcal{P}(Y) \setminus \{\emptyset\} \rightarrow Y$ such that for each $A \subset Y$, $h(A) \in A$. Then, let $f : X \rightarrow Y$ such that

$$f(x) = h(g^{-1}(\{x\})).$$

Clearly so, f is injective.

² The idea here is to collect the preimages into a set, and use the choice function as an injective map.

“ Note 5.1.1

Note that we have $|\mathbb{N}| \leq |\mathbb{Q}|$, since we can define an injective function $f : \mathbb{N} \rightarrow \mathbb{Q}$ such that $f(n) = \frac{n}{1}$.

We have also shown that $|\mathbb{Q}| \leq |\mathbb{N}|$ using our injective function $g : \mathbb{Q} \rightarrow \mathbb{N}$, given by

$$g\left(\frac{m}{n}\right) = \begin{cases} 2^m 3^n & m > 0 \\ 1 & m = 0 \\ 5^{-m} 7^n & m < 0 \end{cases}$$

QUESTION: Is $|\mathbb{N}| = |\mathbb{Q}|$? In other words, given $|X| \leq |Y| \wedge |Y| \leq |X|$, is $|X| = |Y|$?

6 Lecture 6 Sep 19th

6.1 Cardinality (Continued 2)

Before delving into resolving our last question in the previous lecture, note the following:

“ Note 6.1.1

Suppose $f : X \rightarrow Y$ is bijective. Let $A \subseteq B$, then

$$f(B \setminus A) = f(B) \setminus f(A).$$

Prove this observation as an exercise:

Exercise 6.1.1

Prove the note on the left.

📖 Theorem 14 (★ ★ ★ Cantor-Schröder-Bernstein Theorem (CSB))

Let $A_2 \subset A_1 \subset A_0 = A$. Assume that $A_2 \sim A_0$. Then $A_0 \sim A_1$.

✎ Proof

Let $\varphi : A_0 \rightarrow A_2$ be bijective, by assumption. Since $A_1 \subset A_0$, let $A_3 = \varphi(A_1) \subset A_2$, and since $A_2 \subset A_0$, let $A_4 = \varphi(A_2) \subset A_3$. Recursively so, let

$$A_{n+2} = \varphi(A_n)$$




Figure 6.1: The core idea of the proof for Cantor-Schröder-Bernstein Theorem

Notice that

$$A_0 = (A_0 \setminus A_1) \cup (A_1 \setminus A_2) \cup (A_2 \setminus A_3) \cup (A_3 \setminus A_4) \cup \dots \bigcap_{n=0}^{\infty} A_n$$

$$A_1 = (A_1 \setminus A_2) \cup (A_2 \setminus A_3) \cup (A_3 \setminus A_4) \cup (A_4 \setminus A_5) \cup \dots \bigcap_{n=1}^{\infty} A_n$$

Observe that

$$\bigcap_{n=0}^{\infty} A_n = \bigcap_{n=1}^{\infty} A_n.$$

¹Define $f : A \rightarrow A_1$ by

$$f(x) = \begin{cases} x & x \in \bigcap_{n=0}^{\infty} A_n \\ x & x \in A_{2k+1} \setminus A_{2k+2}, k = 0, 1, 2, \dots \\ \varphi(x) & x \in A_{2k} \setminus A_{2k+1}, k = 0, 1, 2, \dots \end{cases} \quad \square$$

¹ Here, we employ the idea from Figure 6.1.

Corollary 15 (Cantor-Schröder-Bernstein Theorem - Restated)

If $A_1 \subset A \wedge B_1 \subset B \wedge A \sim B_1 \wedge B \sim A_1$, then $A \sim B$.²

² This is equivalent to the statement

$$|A| \leq |B| \wedge |B| \leq |A| \implies |A| = |B|.$$

Proof

By assumption, let $f : A \rightarrow B_1$ be bijective, and let $g : B \rightarrow A_1$ be bijective. Let $A_2 = g(B_1) \subseteq A_1 \subset A$. Let $A_2 = g(B_1) \subseteq A_1 \subset A$. Then the composite function $g \circ f : A \rightarrow A_2$ is bijective, and so $A \sim A_2$. By Theorem 14, we have

$$A \sim A_2 \sim A_1 \sim B.$$

Example 6.1.1

Our question from last lecture now has an answer: by Theorem 14, we have that $|\mathbb{Q}| = |\mathbb{N}|$.³

³ Now that we know that they have the same cardinality:

Proposition 16 (Denumerability Check)


Exercise 6.1.2

Find a bijection between \mathbb{Q} and \mathbb{N} .

If X is infinite, then

$$|X| = |\mathbb{N}| = \aleph_0 \iff \exists f : X \rightarrow \mathbb{N} \text{ bijective.}$$

 **Proof**


(\implies) is immediate. For (\impliedby), suppose $f : X \rightarrow \mathbb{N}$, which implies that $|X| \leq |\mathbb{N}|$. By  Proposition 11, $|\mathbb{N}| \leq |X|$. Therefore, $|X| = |\mathbb{N}| = \aleph_0$.

Example 6.1.2


$\mathbb{N} \times \mathbb{N}$ is countable. The function

$$f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N} \text{ given by } f(m, n) = 2^n 3^m$$

is injective. 

 **Definition 27 (Uncountable)**

A set X is *uncountable* if it is not countable.

 **Theorem 17 (Cantor's Diagonal Argument)**

$(0, 1)$ is uncountable.

 **Proof**

Suppose, for contradiction, that $(0, 1)$ is countable. Then we can write

$$a_1 = .a_{11}a_{12}a_{13} \dots$$

$$a_2 = .a_{21}a_{22}a_{23} \dots$$

$$\begin{array}{c} \vdots \\ a_n = .a_{n1}a_{n2}a_{n3} \dots \\ \vdots \end{array}$$

in $(0, 1)$. This representation is unique if we do not allow the representation to end in a string of 9's. Let $b \in (0, 1)$, expressed as $b = .b_1b_2b_3 \dots$ such that

$$b_i = \begin{cases} 5 & a_i \neq 5 \\ 2 & a_i = 5 \end{cases}$$

However, $b \notin (0, 1)$, otherwise b would be one of the a_n 's, a contradiction.

Corollary 18 (Uncountability of \mathbb{R})

\mathbb{R} is uncountable.

Proof

Let $f : (0, 1) \rightarrow \mathbb{R}$ be given by

$$f(x) = \tan\left(\pi x - \frac{\pi}{2}\right).$$

Clearly so, $(0, 1)$ is bijective.

Note 6.1.2

We denote $|\mathbb{R}| = c$.

QUESTION: Given sets X, Y , is it always true that either⁴

1. $|X| = |Y|$;

⁴ As compare to $\leq, <$ implies that there is no surjection from the set on the LHS to the RHS.

7 Lecture 7 Sep 21st

7.1 Cardinality (Continued 3)

Theorem 19 (★ Comparability of Cardinals)

If X and Y are non-empty, then either

$$|X| \leq |Y| \vee |Y| \leq |X|.$$

Proof

Let

$$S = \{(A, B, f) \mid A \subseteq X, B \subseteq Y, f : A \rightarrow B \text{ bijective}\}.$$

Note that $S \neq \emptyset$, since X and Y are non-empty, and so we can have

$f(a) = b$ for $A = \{a\} \subset X$ and $B = \{b\} \subset Y$.¹ We order S as

follows: we say

$$(A_1, B_1, f_1) \leq (A_2, B_2, f_2)$$

if

$$A_1 \subseteq A_2, B_1 \subseteq B_2, f_1 = f_2 \upharpoonright_{A_1}.$$

Let $C = \{(A_\alpha, B_\alpha, f_\alpha)\}_{\alpha \in I}$ be a chain in (S, \leq) . Let $A_0 = \bigcup_{\alpha \in I} A_\alpha$,

$B_0 = \bigcup_{\alpha \in I} B_\alpha$, and define $f_0 : A_0 \rightarrow B_0$ by

$$f_0(x) = f_{\alpha_0}(x) \text{ if } x \in A_{\alpha_0}.$$

¹ We want to use the maximal element to obtain our result. To that end, we need Zorn's Lemma. So we need S to build this up.

Now if $x \in A_{\alpha_1}$, $x \in A_{\alpha_2}$ and

$$(A_{\alpha_1}, B_{\alpha_1}, f_{\alpha_1}) \leq (A_{\alpha_2}, B_{\alpha_2}, f_{\alpha_2}),$$

we have that

$$f_{\alpha_1}(x) = f_{\alpha_2} \upharpoonright_{A_{\alpha_1}}(x) = f_{\alpha_2}(x),$$

i.e. f_0 is well-defined.

Claim 1: $f_0 : A_0 \rightarrow B_0$ is injective.

Let $x_1, x_2 \in A_0$ such that $x_1 \neq x_2$.

$$\implies \exists \alpha_1, \alpha_2 \in I \quad x_1 \in A_{\alpha_1} \wedge x_2 \in A_{\alpha_2} \wedge A_{\alpha_1} \subseteq A_{\alpha_2} \text{ (wlog)}$$

$$\implies x_1, x_2 \in A_{\alpha_2}$$

$$\implies (\because f_{\alpha_2} \text{ injective} \implies f_{\alpha_2}(x_1) \neq f_{\alpha_2}(x_2))$$

$$\implies f_0(x_1) \neq f_0(x_2) \implies f_0 \text{ injective.}$$

Claim 2: $f_0 : A_0 \rightarrow B_0$ is surjective.

Let $y_0 \in B_0$

$$\implies \exists \alpha_0 \in I \quad y_0 \in B_{\alpha_0}$$

$$\implies \exists x_0 \in A_{\alpha_0} \quad f_{\alpha_0}(x_0) = y_0 \text{ (}\because f_{\alpha_0} \text{ surjective)}$$

$$\implies f_0(x_0) = y_0$$

$\therefore (A_0, B_0, f_0)$ is an upper bound for C . Then by Zorn's Lemma, (S, \leq) has a maximal element (A, B, f) .

Case 1: If $A = X$, then injectivity of f implies $|X| \leq |Y|$.

Case 2: If $B = Y$, then surjectivity of f implies $|Y| \leq |A| \leq |X|$.

Case 3: If $A \neq X \wedge B \neq Y$, then $X \setminus A \neq \emptyset \wedge Y \setminus B \neq \emptyset$. Let $x_0 \in X \setminus A$, $y_0 \in Y \setminus B$. Let $A^* = A \cup \{x_0\}$, $B^* = B \cup \{y_0\}$, and $f^* : A^* \rightarrow B^*$ such that

$$f^*(x) = \begin{cases} f(x) & x \in A \\ y_0 & x = x_0 \end{cases}$$

Then $(A, B, f) \leq (A^*, B^*, f^*)$, contradicting maximality.

7.1.1 Cardinal Arithmetic

Sum of Cardinals Observe that if $X = \{x_1, \dots, x_n\}$, $Y = \{y_1, \dots, y_m\}$, and $X \cap Y = \emptyset$, then $|X| = n$, $|Y| = m$ and $|X \cup Y| = n + m$. This motivates us to provide the following definition:

Definition 28 (Sum of Cardinals)

Assume that X and Y are such that $X \cap Y = \emptyset$. We define

$$|X| + |Y| = |X \cup Y|.$$

QUESTION: So what about $\aleph_0 + \aleph_0$?

A thought that motivates us to give the following answer lies in the observation that: if X is the set of even natural numbers and Y the odd natural numbers, then $X \cup Y$ is the set of all natural numbers. All three sets are countably infinite, i.e. they have cardinality \aleph_0 .

QUESTION: What about $c + c$?

A similar motivation comes from the observation that: given $X = (0, 1)$ and $Y = (1, 2)$, we have

$$c = |X| \leq |X| + |Y| \leq |R| = c,$$

and so $|X| = |Y| = c \implies |X \cup Y| = c$.

Theorem 20 (Sums of Cardinals)

Given sets X and Y , if X is infinite, then

1. $|X| + |X| = |X|$
 2. $|X| + |Y| = \max(|X|, |Y|)$
-

Exercise 7.1.1

Prove **Theorem 20** as an exercise.

Multiplication of Cardinals Given

$$X = \{x_1, x_2, \dots, x_n\}$$

$$Y = \{y_1, y_2, \dots, y_m\}$$

we have that

$$X \times Y = \{(x_i, y_j) \mid i = 1, 2, \dots, n, j = 1, 2, \dots, m\}$$

and so

$$|X \times Y| = nm.$$

Definition 29 (Multiplication of Cardinals)

Given sets X and Y , define

$$|X| |Y| = |X \times Y|.$$

Example 7.1.1

We have $|\mathbb{N} \times \mathbb{N}| = \aleph_0$ since the function $f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ given by

$$f(n, m) = 2^n 3^m$$

is injective. 


QUESTION: What about $c \cdot c$?

Theorem 21 (Multiplication of Cardinals)

If X is infinite and $Y \neq \emptyset$, then

- $|X \times X| = |X| \implies |X| |X| = |X|$;
- $|X \times Y| = \max(|X|, |Y|)$.

Exercise 7.1.2

Prove  Theorem 21 as an exercise.

8 Lecture 8 Sep 24th

8.1 Cardinality (Continued 4)

8.1.1 Cardinal Arithmetic (Continued)

Exponentiation of Cardinals Recall if $\{Y_x\}_{x \in X}$ is a collection of non-empty sets, we have¹

¹ This should remind you of * Axiom 3

$$\prod_{x \in X} Y_x = \{f : X \rightarrow \bigcup_{x \in X} Y_x \mid f(x) \in Y_x\}.$$

Now if $Y = Y_x$ for all $x \in X$, we have

$$Y^X = \prod_{x \in X} Y = \{f : X \rightarrow Y\}.$$

Example 8.1.1

Given

$$Y = \{1, \dots, m\} \quad X = \{1, \dots, n\}$$

we have

$$Y^X = \{f : \{1, \dots, n\} \rightarrow \{1, \dots, m\}\}.$$

Observe that Y^X is similar to Y^n . So $|Y^X| = m^n$.²



² Need better explanation.

Definition 30 (Exponentiation of Cardinals)

Given sets X and Y , define

$$|Y|^{|X|} := |Y^X|.$$

Theorem 22 (Exponentiation of Cardinals)

If X, Y, Z are non-empty sets, then

- $|Y|^{|X|} \cdot |Y|^{|Z|} = |Y|^{|X|+|Z|}$;
- $\left(|Y|^{|X|}\right)^{|Z|} = |Y|^{|X|\cdot|Z|}$.

Theorem 23 ($2^{\aleph_0} = c$)

We have that $2^{\aleph_0} = c$.

Proof

Note that $2^{\aleph_0} = |\{0, 1\}^{\mathbb{N}}|$, where³

$$|\{0, 1\}^{\mathbb{N}}| = |\{f : \mathbb{N} \rightarrow \{0, 1\}\}| = |\{\{a_n\}_{n=1}^{\infty} \mid a_i = 0, 1\}|$$

Given a sequence $\{a_n\} \in \{0, 1\}^{\mathbb{N}}$, define $\varphi : \{0, 1\}^{\mathbb{N}} \rightarrow (0, 1)$ such that⁴

$$\varphi(\{a_n\}) := \sum_{i=1}^{\infty} \frac{a_n}{3^n}.$$

which is injective since there are no trailing 2's. Therefore $2^{\aleph_0} \leq c$.

Given $\alpha \in (0, 1)$, let⁵

$$\alpha = \sum_{n=1}^{\infty} \frac{b_n}{2^n},$$

where $b_n = 0, 1$. Let $\psi : (0, 1) \rightarrow \{0, 1\}^{\mathbb{N}}$ such that

$$\psi(\alpha) = \psi\left(\sum_{i=1}^{\infty} \frac{b_n}{2^n}\right) = \{b_n\}$$

Then ψ is injective, and so $c \leq 2^{\aleph_0}$. Thus $2^{\aleph_0} = c$ as required.

Exercise 8.1.1

Prove **Theorem 22**.

This requires closer studying.

³ Explain 2nd equality.

⁴ This is a base 3 representation (of what?)

⁵ This is a base 2 representation.

Example 8.1.2

Find $|\mathbb{N}_0^{\mathbb{N}_0}|$ and $c^{\mathbb{N}_0}$.



Solution

We have that

$$c = 2^{\mathbb{N}_0} \leq \mathbb{N}_0^{\mathbb{N}_0} \leq c^{\mathbb{N}_0} = (2^{\mathbb{N}_0})^{\mathbb{N}_0} = 2^{\mathbb{N}_0 \cdot \mathbb{N}_0} = 2^{\mathbb{N}_0} = c$$

Example 8.1.3

Show $|\mathcal{P}(X)| = 2^{|X|} = |2^X|$.



Solution

Given $f : X \rightarrow \{0, 1\}$, let⁶

⁶ A is a collection of all x 's that gets mapped to f .

$$A = \{x \in X \mid f(x) = 1\} \subset X.$$

Define $\Gamma : 2^X \rightarrow \mathcal{P}(X)$ by

$$\Gamma(f) = f^{-1}(\{1\})$$

Γ is injective⁷.

⁷ Why?

Conversely, given $A \subset X$, define the characteristic function

$$\chi_A(x) = \begin{cases} 1 & x \in A \\ 0 & x \notin A \end{cases} \in 2^X.$$

Then define $\Phi : \mathcal{P}(X) \rightarrow 2^X$ such that

$$\Phi(A) = \chi_A.$$

Clearly so, Φ is injective.

Theorem 24 (Russell's Paradox)

For any X , we have $|X| < |\mathcal{P}(X)| = 2^{|X|}$.

Proof

Let $f : X \rightarrow \mathcal{P}(X)$ be $f(x) = \{x\}$. Clearly, f is injective, and so $|X| < |\mathcal{P}(X)|$.

Claim: $\nexists g : X \rightarrow \mathcal{P}(X)$ surjective.

Suppose not. Let⁸

$$A = \{x \in X \mid x \notin g(x)\}$$

Pick $x_0 \in X$ with $g(x_0) = A$. Now if $x_0 \in A$, then $x_0 \in g(x_0)$, but this implies that $x_0 \notin A$, a contradiction.

So $x_0 \notin A$, i.e. $x_0 \notin g(x_0)$, which in turn implies that $x_0 \in A$, yet another contradiction. Therefore such a function g cannot exist, as claimed.

Therefore, we have $|X| < |\mathcal{P}(X)|$ as required.

⁸ By the **Bounded Separation Axiom** (see ZF Set Theory), this is a set, and since it is a subset of X , it is a valid element of $\mathcal{P}(X)$. Thus, we can consider such a set.

QUESTION: Is there anything between \aleph_0 and c ?

♣ Axiom 25 (Continuum Hypothesis)

If $\aleph_0 \leq \gamma \leq c$, then either $\gamma = \aleph_0$ or $\gamma = c$.

♣ Axiom 26 (Generalized Continuum Hypothesis)

If $|X| \leq \gamma \leq 2^{|X|}$, then either $\gamma = |X|$ or $\gamma = 2^{|X|}$.

In this course, we shall assume that the Continuum Hypothesis is true.

9 Lecture 9 Sep 26th

9.1 Introduction to Metric Spaces

Definition 31 (Metric & Metric Space)

Given a set X , a *metric* on X is a function $d : X \times X \rightarrow \mathbb{R}$ such that

1. (*positive definiteness*) $d(x, y) \geq 0$ and $d(x, y) = 0 \iff x = y$;
2. (*symmetry*) $d(x, y) = d(y, x)$; and
3. (*triangle inequality*) $d(x, y) \leq d(x, z) + d(z, y)$.

The pair (X, d) is called a *metric space*.

Remark 9.1.1

A metric is an abstract notion of distance.

Example 9.1.1 (Standard Metric on \mathbb{R})

Let $X = \mathbb{R}$, and let $d(x, y) = |x - y|$.

Clearly so, the first 2 criterias are satisfied:

- $|\cdot| \geq 0$ and $|x - y| = 0 \iff x = y$; and
- $|x - y| = |y - x|$.

The triangle inequality property is the usual triangle inequality of the absolute value function, i.e.

$$|x - y| \leq |x| + |y|.$$



QUESTION: For an arbitrary set X , can we define a metric on X ? The


following example shows that we can,

Example 9.1.2 (Discrete Metric)

Let X be any set. We can simply define

$$d(x, y) = \begin{cases} 1 & x \neq y \\ 0 & x = y \end{cases}$$

This metric clearly satisfies all 3 criterias of being a metric:

- $d : X \times X \rightarrow \{0, 1\}$ and so $d(x, y) \geq 0$, and by definition, we have $d(x, y) = 0 \iff x = y$;
- By definition, $d(x, y) = d(y, x)$ as it does not matter how the pair is ordered; and
- Since $d(x, y) \geq 0$, we have that $d(x, y) \leq d(x, z) + d(y, z)$. 

Example 9.1.3 (Euclidean Metric / 2-metric on \mathbb{R}^n)


Let $X = \mathbb{R}^n$. Let $\vec{x} = \{x_1, x_2, \dots, x_n\}$ and $\vec{y} = \{y_1, y_2, \dots, y_n\}$. Define

$$d_2(\vec{x}, \vec{y}) = \sqrt{\sum_{i=1}^n (x_i - y_i)^2}$$

Note that in \mathbb{R}^2 , this is our regular (Euclidean) distance between two points.

It is not difficult to see that d_2 satisfies the first 2 criterion to being a metric:

- d_2 is the square root of the sum of squares, and so $d_2(\vec{x}, \vec{y}) \geq 0$ for any $\vec{x}, \vec{y} \in \mathbb{R}^n$, and $d_2(\vec{x}, \vec{y}) = 0 \iff \forall i \in \{1, \dots, n\} x_i = y_i \iff \vec{x} = \vec{y}$;
- Since $(x_i - y_i)^2 = (y_i - x_i)^2$ for any $x_i, y_i \in \mathbb{R}$, we have that $d_2(\vec{x}, \vec{y}) = d_2(\vec{y}, \vec{x})$.

However, it is not immediately clear that d_2 satisfies the triangle inequality criterion, especially if $n \geq 3$. If $n = 2$, heuristically, the triangle inequality simply tells that the length of any one side of a triangle is less than or equal to the sum of the other two, e.g. Figure 9.1. 





Figure 9.1: A visualization of the triangle inequality in \mathbb{R}^2 .

Remark 9.1.2

Many of the important examples of metric spaces are vector spaces with an abstract length function, or *norm*. 

Definition 32 (Norm & Normed Linear Space)

Given a vector space V (usually over \mathbb{R}), a *norm* on V is a function

$$\|\cdot\| : V \rightarrow \mathbb{R}$$

such that

1. (*positive definiteness*) $\|v\| \geq 0$ and $\|v\| = 0 \iff v = 0$;
2. (*scalar multiplication*) $\|\alpha \cdot v\| = |\alpha| \|v\|$; and
3. (*triangle inequality*) $\|v + w\| \leq \|v\| + \|w\|$.

The pair $(V, \|\cdot\|)$ is called a *normed linear space*.

Remark 9.1.3

Given a normed linear space $(V, \|\cdot\|)$, a natural metric, $d_{\|\cdot\|}$, on V induced by $\|\cdot\|$ can be defined as

$$d_{\|\cdot\|}(x, y) = \|x - y\|.$$

Exercise 9.1.1

Prove that $d_{\|\cdot\|}$ is indeed a metric.

Proof (Exercise 9.1.1)

1. (positive definiteness) It is clear from the definition of a norm that $d_{\|\cdot\|}(x, y) = \|x - y\| \geq 0$, and $d_{\|\cdot\|}(x, y) = 0 \iff x - y = 0 \iff x = y$.
2. (symmetry) Symmetry follows simply from definition, as $\|x - y\| = \|y - x\|$.
3. (triangle inequality) For $x, y, z \in V$, we have

$$\begin{aligned} d_{\|\cdot\|}(x, y) &= \|x - y\| = \|x - z + z - y\| \\ &\leq \|x - z\| + \|z - y\| \quad \because \text{triangle inequality of norms} \\ &= \|x - z\| + \|y - z\| \quad \because \text{symmetry} \\ &= d_{\|\cdot\|}(x, z) + d_{\|\cdot\|}(y, z) \end{aligned}$$

Example 9.1.4 (Euclidean Norm)

Let $X = \mathbb{R}^n$, and $\vec{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$. Define $\|\cdot\|_2$ such that

$$\|(x_1, \dots, x_n)\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

From Example 9.1.3, we are given the triangle inequality property, in which we have yet to prove. Positive definiteness is clear. For scalar multiplication, let $\vec{x} = (x_1, \dots, x_n)$, and notice that

$$\|\alpha \cdot \vec{x}\|_2 = \sqrt{\sum_{i=1}^n (\alpha x_i)^2} = \sqrt{\alpha^2 \sum_{i=1}^n x_i^2} = |\alpha| \sqrt{\sum_{i=1}^n x_i^2} = |\alpha| \|\vec{x}\|_2.$$

Thus $\|\cdot\|_2$ is indeed a norm. We call $\|\cdot\|_2$ the **2-norm** or the **Euclidean norm**.

We observe that, in comparison with Example 9.1.3, we have that

$$d_2(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|_2.$$

Example 9.1.5 (1-norm)

Let $X = \mathbb{R}^n$, and $\vec{x} = (x_1, \dots, x_n)$. Define

$$\|\vec{x}\|_1 := \sum_{i=1}^n |x_i|.$$


Clearly so, $\|\cdot\|_1$ is a norm:

- **(positive definiteness)** This is true by the absolute value function, i.e. every $|x_i| \geq 0$, and so the sum over these x_i 's is also non-negative, and $\sum_{i=1}^n |x_i| = 0 \iff \forall i \in \{1, \dots, n\} x_i = 0 \iff \vec{x} = 0$.
- **(scalar multiplication)** This uses a similar argument as in the previous example.
- **(triangle inequality)** This is true by, again, the triangle inequality on absolute values.

We call $\|\cdot\|_1$ the **1-norm**.

Thus, we can define

$$d_1(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|_1$$

and it can easily be verified that d_1 is indeed a metric. 

Example 9.1.6

Let $X = \mathbb{R}^n$ and $\vec{x} = (x_1, \dots, x_n)$. Define

$$\|\vec{x}\|_\infty = \max\{|x_i|\}$$

Again, it is easy to see that $\|\cdot\|_\infty$ is a norm;

- **(positive definiteness)** $\because \forall i \in \{1, \dots, n\} \quad |x_i| \geq 0 \implies \max\{|x_i|\} \geq 0$ and $\max\{|x_i|\} = 0 \iff x_i = 0 \iff \vec{x} = 0$.
- **(scalar multiplication)** Notice that


$$\|\alpha \cdot \vec{x}\|_\infty = \max\{|\alpha x_i|\} = |\alpha| \max\{|x_i|\} = |\alpha| \|\vec{x}\|_\infty.$$

- **(triangle inequality)** This is once again true by the triangle inequality on the absolute value function, i.e.

$$\begin{aligned} \because \forall i \in \{1, \dots, n\} \quad |x_i + y_i| &\leq |x_i| + |y_i| \\ \max\{|x_i + y_i|\} &\leq \max\{|x_i|\} + \max\{|y_i|\}. \end{aligned}$$

We can then define

$$d_\infty(\vec{x}, \vec{y}) = \max\{|x_i - y_i|\},$$

which we can easily verify that it is indeed a metric¹. 

¹ Symmetry holds by the property of the absolute value function.

HERE'S an interesting notion: let

$$S_i = \{\vec{x} \in \mathbb{R}^2 \mid \|\vec{x}\|_i = 1\}, \quad i = 1, 2, \infty$$

Notice that we would then have the following graph: In fact, it is true that if we let $i \in \mathbb{N} \setminus \{0\}$, as suggested by Figure 9.2, we would see that the “diamond” would grow into a “circle” as in S_2 , and as $i \geq 3$, the unit ball will expand and approach the “square”, which is S_∞ .




Figure 9.2: Unit ball depending on $\|\vec{x}\|_i$

Another observation that we can make is if we can show that a set is open for a “smaller” S_i , then the same set is open for any S_j for $j \geq i$.

If we apply these norms into metrics, we have

$$d_\infty \leq d_2 \leq d_1$$

where we say that d_∞ is the **least sensitive**, and d_1 being the **most sensitive**².


Example 9.1.7

For $1 < p < \infty$, define on \mathbb{R}^n

$$\|\vec{x}\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}$$

Continuing with the same idea as in previous examples, we can let

$$d_p(\vec{x}, \vec{y}) = \left(\sum_{i=1}^n |x_i - y_i|^p \right)^{\frac{1}{p}}$$

In the next lecture, we will go into proving that this is indeed a norm, and so we can define a metric using this norm. 

Note that if we allow for $0 < i < 1$, then we would have a graph that looks like the following, which is a convex graph, i.e. we cannot create well-defined norms.




Figure 9.3: $\|\cdot\|_p$ for $0 < p < 1$

² For sufficiently close points, we see that d_∞ would reflect the least change, while we can see change in d_1 for every two points that we take.

10 Lecture 10 Sep 28th

10.1 Introduction to Metric Spaces (Continued)

Definition 33 ($\|\cdot\|_p$ -norm)

Given $\vec{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$, we define, for $1 < p < \infty$, the $\|\cdot\|_p$ -norm to be

$$\|\vec{x}\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}$$

We asked the question: why is $\|\cdot\|_p$ a norm?

Lemma 27 (Young's Inequality)

If $1 < p < \infty$,

$$\frac{1}{p} + \frac{1}{q} = 1,$$

and if $\alpha, \beta > 0$, then

$$\alpha \cdot \beta \leq \frac{\alpha^p}{p} + \frac{\beta^q}{q}.$$

Proof

Motivated by Figure 10.1, using notions from calculus, we have from calculus,




Figure 10.1: Motivation for Lemma 27.

$$\begin{aligned}
\alpha\beta &\leq \int_0^\alpha t^{p-1} dt + \int_0^\beta u^{q-1} du \\
&= \frac{t^p}{p} \Big|_0^\alpha + \frac{u^q}{q} \Big|_0^\beta \\
&= \frac{\alpha^p}{p} + \frac{\beta^q}{q},
\end{aligned}$$

where we note that

$$\begin{aligned}
\frac{1}{p} + \frac{1}{q} &= 1 \\
\frac{q}{p} &= q - 1 \\
\frac{p}{q} &= p - 1 \\
1 &= (p - 1)(q - 1)
\end{aligned}$$

Theorem 28 (Hölder's Inequality)

For $1 < p < \infty$, let $\frac{1}{p} + \frac{1}{q} = 1$ ¹. Let

$$\vec{x} = (x_1, \dots, x_n), \quad \vec{y} = (y_1, \dots, y_n).$$

Then

$$\sum_{i=1}^n |x_i y_i| \leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} \cdot \left(\sum_{i=1}^n |y_i|^q \right)^{\frac{1}{q}}.$$

¹ We also call q the conjugate of p .

Note 10.1.1

Note that $p = 2$ is just the *Cauchy-Schwarz Inequality*:

$$\begin{aligned}
\sum_{i=1}^n |x_i y_i| &\leq \left(\sum_{i=1}^n |x_i|^2 \right)^{\frac{1}{2}} \cdot \left(\sum_{i=1}^n |y_i|^2 \right)^{\frac{1}{2}} \implies \\
\left(\sum_{i=1}^n |x_i y_i| \right)^2 &\leq \left(\sum_{i=1}^n |x_i|^2 \right) \cdot \left(\sum_{i=1}^n |y_i|^2 \right)
\end{aligned}$$

 **Proof**

Since if either \vec{x} or \vec{y} is zero, then we have that the inequality is trivially true, we can suppose that $\vec{x} \neq 0 \neq \vec{y}$. Now, note that for $\alpha, \beta \neq 0$, we have that²

$$\begin{aligned} \sum_{i=1}^n |x_i y_i| &\leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} \cdot \left(\sum_{i=1}^n |y_i|^q \right)^{\frac{1}{q}} \\ &\iff \\ \sum_{i=1}^n |\alpha x_i \cdot \beta y_i| &\leq \left(\sum_{i=1}^n |\alpha x_i|^p \right)^{\frac{1}{p}} \cdot \left(\sum_{i=1}^n |\beta y_i|^q \right)^{\frac{1}{q}}. \end{aligned}$$

² In the second inequality, notice that we can easily get back to the first equation by dividing both sides by $\alpha\beta$.

So we can assume that

$$\left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} = 1 = \left(\sum_{i=1}^n |y_i|^q \right)^{\frac{1}{q}}, \quad (10.1)$$

and if not, we can simply choose $\alpha, \beta \neq 0$ to scale these values to become one. By Lemma 27, we have

$$|x_i y_i| \leq \frac{|x_i|^p}{p} + \frac{|y_i|^q}{q}.$$

Hence

$$\begin{aligned} \sum_{i=1}^n |x_i y_i| &\leq \sum_{i=1}^n \frac{|x_i|^p}{p} + \sum_{i=1}^n \frac{|y_i|^q}{q} = \frac{1}{p} + \frac{1}{q} \quad \because \text{Equation (10.1)} \\ &= 1 = \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} \cdot \left(\sum_{i=1}^n |y_i|^q \right)^{\frac{1}{q}} \end{aligned}$$

as required.

We are now ready to prove our long-awaited result.

 **Theorem 29 (Minkowski's Inequality)**

Let $1 < p < \infty$. If

$$\vec{x} = (x_1, \dots, x_n), \quad \vec{y} = (y_1, \dots, y_n)$$

in \mathbb{R}^n , then

$$\left(\sum_{i=1}^n |x_i + y_i|^p \right)^{\frac{1}{p}} \leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}},$$

i.e.

$$\|\vec{x} + \vec{y}\|_p \leq \|\vec{x}\|_p + \|\vec{y}\|_p.$$

Proof

Let

$$\frac{1}{p} + \frac{1}{q} = 1.$$

Once again, we may assume that $\vec{x} \neq 0 \neq \vec{y}$, as otherwise the inequality is true trivially so. Now, notice that

$$\begin{aligned} \sum_{i=1}^n |x_i + y_i|^p &= \sum_{i=1}^n |x_i + y_i| |x_i + y_i|^{p-1} \\ &\leq \sum_{i=1}^n |x_i| |x_i + y_i|^{p-1} + \sum_{i=1}^n |y_i| |x_i + y_i| \quad \text{triangle inequality} \\ &\leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n |x_i + y_i|^{(p-1)q} \right)^{\frac{1}{q}} \\ &\quad + \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n |x_i + y_i|^{(p-1)q} \right)^{\frac{1}{q}} \end{aligned}$$

where the last step is by Hölder's Inequality. Note that $\frac{1}{p} + \frac{1}{q} = 1 \implies p = q(p-1)$. Thus

$$\begin{aligned} \sum_{i=1}^n |x_i + y_i|^p &\leq \left[\left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}} \right] \cdot \left(\sum_{i=1}^n |x_i + y_i|^p \right)^{\frac{1}{q}} \\ &\implies \left(\sum_{i=1}^n |x_i + y_i|^p \right)^{1 - \frac{1}{q}} \leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}} \end{aligned}$$

Note 10.1.2

With this we have that $\|\cdot\|_p$ satisfies the triangle inequality condition, and so $\|\cdot\|_p$ is a norm on \mathbb{R}^n .

Note 10.1.3

Given $1 \leq p \leq q < \infty$, we have³

$$\|\cdot\|_\infty \leq \|\cdot\|_q \leq \|\cdot\|_p \leq \|\cdot\|_1.$$

³ For a visual representation of this result, see Figure 9.2.

 **Proof**

It is quite clear that $\forall p \geq 1$,

$$\|\cdot\|_\infty = \max\{|\cdot|\} \leq \left(\sum |\cdot|^p\right)^{\frac{1}{p}} = \|\cdot\|_p.$$

For $1 \leq p \leq q < \infty$, consider Holder's Inequality, where we have

$$\sum_{i=1}^n |a_i| |b_i| \leq \left(\sum_{i=1}^n |a_i|^r\right)^{\frac{1}{r}} \cdot \left(\sum_{i=1}^n |b_i|^{\frac{r}{r-1}}\right)^{1-\frac{1}{r}}.$$

Let $|a_i| = |x_i|^p$, $|b_i| = 1$ and $r = \frac{q}{p} \geq 1$ ⁴. Then we have

⁴ Note that this is true by $p \leq q$.

$$\begin{aligned} \sum_{i=1}^n |x_i|^p &\leq \left(\sum_{i=1}^n |x_i|^q\right)^{\frac{p}{q}} \cdot \left(\sum_{i=1}^n 1^{\frac{q}{q-p}}\right)^{1-\frac{p}{q}} \\ &= n^{1-\frac{p}{q}} \cdot \left(\sum_{i=1}^n |x_i|^q\right)^{\frac{p}{q}} \end{aligned}$$

Therefore, for $\vec{x} = (x_1, \dots, x_n) \in \mathbb{R}$,

$$\begin{aligned} \|\vec{x}\|_p &= \left(\sum_{i=1}^n |x_i|^p\right)^{\frac{1}{p}} \leq \left(n^{1-\frac{p}{q}} \cdot \left(\sum_{i=1}^n |x_i|^q\right)^{\frac{p}{q}}\right)^{\frac{1}{p}} \\ &= n^{\frac{1}{p}-\frac{1}{q}} \cdot \left(\sum_{i=1}^n |x_i|^q\right)^{\frac{1}{q}} = n^{\frac{1}{p}-\frac{1}{q}} \cdot \|\vec{x}\|_q. \end{aligned}$$

Thus, we have

$$\|\cdot\|_q \leq \|\cdot\|_p.$$

The chain of inequality follows.

Example 10.1.1 (Sequence Spaces)

1. Let $\ell_1 = \left\{ \{x_i\} \mid \sum_{i=1}^{\infty} |x_i| < \infty \right\}$. Define

$$\|\{x_i\}\|_1 = \sum_{i=1}^{\infty} |x_i|$$

Let $\{x_i\}, \{y_i\} \in \ell_1$. Observe that $\forall n \in \mathbb{N}$, we have

$$\sum_{i=1}^n |x_i + y_i| \leq \sum_{i=1}^n |x_i| + \sum_{i=1}^n |y_i| \leq \|\{x_i\}\|_1 + \|\{y_i\}\|_1.$$

Then by the Monotone Convergence Theorem, we have that

$$\sum_{i=1}^{\infty} |x_i + y_i| \leq \|\{x_i\}\|_1 + \|\{y_i\}\|_1.$$

Thus $\{x_i + y_i\} \in \ell_1$ and

$$\|\{x_i + y_i\}\|_1 \leq \|\{x_i\}\|_1 + \|\{y_i\}\|_1.$$

Let $\{x_n\} \in \ell_1$ and $\alpha \in \mathbb{R}$. Then

$$\sum_{i=1}^{\infty} |\alpha x_i| = |\alpha| \sum_{i=1}^{\infty} |x_i|.$$

Therefore $\{\alpha x_n\} \in \ell_1$ and $\|\{\alpha x_n\}\|_1 = |\alpha| \|\{x_n\}\|_1$.

Thus, we have that ℓ_1 is a vector space, and $(\ell_1, \|\cdot\|_1)$ is a normed linear space.

2. Let $\ell_{\infty}(\mathbb{N}) = \ell_{\infty} = \left\{ \{x_i\} \mid \{x_i\} \text{ is bounded} \right\}$. Define

$$\|\{x_i\}\|_{\infty} = \sup\{|x_i| \mid i \in \mathbb{N}\}.$$

Observe that $\forall \{x_i\}, \{y_i\} \in \ell_{\infty}$, then $\forall i \in \mathbb{N}$, we have

$$|x_i + y_i| \leq |x_i| + |y_i| \leq \|\{x_i\}\|_{\infty} + \|\{y_i\}\|_{\infty}.$$

So $\{x_i + y_i\} \in \ell_{\infty}$, and

$$\|\{x_i + y_i\}\|_{\infty} \leq \|\{x_i\}\|_{\infty} + \|\{y_i\}\|_{\infty}.$$

11 Lecture 11 Oct 01st

11.1 Introduction to Metric Spaces (Continued 2)

We wondered about $\ell_p(\mathbb{R})$ in the last lecture but let us consider a case that is even more general.

QUESTION: Can we define $\ell_p(\Gamma)$ for any set Γ ?

Example 11.1.1

Let $\ell_\infty(\Gamma) = \{f : \Gamma \rightarrow \mathbb{R} \mid f(\Gamma) \text{ is bounded}\}$. If $f \in \ell_\infty(\Gamma)$, define

$$\|f\|_\infty = \sup\{|f(x)| \mid x \in \Gamma\}.$$

Notice that for $f, g \in \ell_\infty(\Gamma)$, and $\alpha \in \mathbb{R}$, then we have, by the Triangle Inequality,

$$\begin{aligned}\|f + g\|_\infty &= \sup\{|(f + g)(x)| \mid x \in \Gamma\} \\ &= \sup\{|f(x) + g(x)| \mid x \in \Gamma\} \\ &\leq \sup\{|f(x)| \mid x \in \Gamma\} + \sup\{|g(x)| \mid x \in \Gamma\} \\ &= \|f\|_\infty + \|g\|_\infty.\end{aligned}$$

So $f + g \in \ell_\infty(\Gamma)$, and


$$\|f + g\|_\infty \leq \|f\|_\infty + \|g\|_\infty.$$

Also, we have

$$\begin{aligned}\|\alpha f\|_\infty &= \sup\{|(\alpha f)(x)| \mid x \in \Gamma\} \\ &= \sup\{|\alpha| |f(x)| \mid x \in \Gamma\}\end{aligned}$$

$$\begin{aligned}
 &= |\alpha| \sup\{|f(x)| \mid x \in \Gamma\} \\
 &= |\alpha| \|f\|_\infty.
 \end{aligned}$$

So $\alpha f \in \ell_\infty(\Gamma)$, and $\|\alpha f\|_\infty = |\alpha| \|f\|_\infty$.


Therefore, $(\ell_\infty(\Gamma), \|\cdot\|_\infty)$ is a normed linear space. 

Example 11.1.2

Let $\ell_1(\Gamma) = \{f : \Gamma \rightarrow \mathbb{R} \mid P(f)\}$, where $P(f)$ is the statement

$$\|f\|_1 = \sup \left\{ \sum_{i=1}^n |f(x_i)| \mid x_1, \dots, x_n \in \Gamma, n \in \mathbb{N} \setminus \{0\} \right\} < \infty.$$

It is clear that $\ell_1(\Gamma) \subseteq \ell_\infty(\Gamma)$, where $\ell_\infty(\Gamma)$ is from Example 11.1.1.

Consequently, $(\ell_1(\Gamma), \|\cdot\|_1)$ is a normed linear space. 

We can extend the same idea onto ℓ_p spaces.

Example 11.1.3

Let $X = C[a, b] = \{f : [a, b] \rightarrow \mathbb{R} \mid f \text{ is continuous on } [a, b]\}$, and define¹

$$\begin{aligned}
 \|f\|_\infty &= \sup\{|f(x)| \mid x \in [a, b]\} \\
 &= \max\{|f(x)| \mid x \in [a, b]\}
 \end{aligned}$$

By (regular) Triangle Inequality, for any $f, g \in C[a, b]$, we have

$$\begin{aligned}
 \|f + g\|_\infty &= \max\{|f(x) + g(x)| \mid x \in [a, b]\} \\
 &\leq \max\{|f(x)| \mid x \in [a, b]\} + \max\{|g(x)| \mid x \in [a, b]\} \\
 &= \|f\|_\infty + \|g\|_\infty,
 \end{aligned}$$

and, for $\alpha \in \mathbb{R}$,

$$\begin{aligned}
 \|\alpha f\|_\infty &= \max\{|\alpha f(x)| \mid x \in [a, b]\} \\
 &= |\alpha| \max\{|f(x)| \mid x \in [a, b]\} \\
 &= |\alpha| \|f\|_\infty.
 \end{aligned}$$

Thus $\|\cdot\|_\infty$ is a norm on $C[a, b]$, and $(C[a, b], \|\cdot\|_\infty)$ is a normed linear space^{2,3}.

Require clarification Notice that $\forall f \in \ell_1(\Gamma)$, for each $n \in \mathbb{N}$,

$$A_n = \{x \in \Gamma \mid |f(x)| \geq \frac{1}{n}\} \text{ is finite.}$$

So

$$A_0 = \bigcup_{n=1}^{\infty} A_n \text{ is countable.}$$

and

$$A_0 = \{x \in \Gamma \mid |f(x)| \neq 0\}$$

¹ Note in this case sup is also max, since we are on a closed interval.

² This space is complete.

³ This space is important for us for the purpose of this course.

Also, observe that

$$C[a, b] \subset \ell_\infty([a, b]).$$

Example 11.1.4

Let $X = C[a, b]$ ⁴ have the same definition as the previous example, but this time define

$$\|f\|_1 = \int_a^b |f(t)| dt.$$

By **linearity of integration**, both the triangle equality and scalar multiplication hold, and so $(C[a, b], \|\cdot\|_1)$ is a normed linear space⁵.

⁴ Some authors also write this as $L^1[a, b]$.

⁵ Compared to the last example, this is not a complete space.

Example 11.1.5

Let $X = C[a, b]$, and $1 < p < \infty$. Define

$$\|f\|_p = \left(\int_a^b |f(x)|^p dx \right)^{\frac{1}{p}}$$

Again, by linearity of integration, scalar multiplication holds. However, it is not as easy to show for the triangle inequality; we are now asking the same question as we did before for ℓ_p , which we solved using Hölder's Inequality and Minkowski's Inequality. But now, instead of summations, we have integrations.


Theorem 30 (Hölder's Inequality v2)

Let $1 < p < \infty$ and $\frac{1}{p} + \frac{1}{q} = 1$. For each $f, g \in C[a, b]$, we have

$$\int_a^b |f(t)g(t)| dt \leq \left(\int_a^b |f(t)|^p dt \right)^{\frac{1}{p}} \left(\int_a^b |g(t)|^q dt \right)^{\frac{1}{q}}.$$

Proof

If either $f(x) = 0$ or $g(x) = 0$ for all $x \in [a, b]$, then the inequality holds trivially so. Thus, we may assume that $\forall x \in [a, b], f(x) \neq 0 \neq g(x)$. By the linearity of integration, we can apply the same

reasoning as we did in  Theorem 28, and assume that

$$\int_a^b |f(t)|^p dt = 1 = \int_a^b |g(t)|^q dt$$

By Lemma 27, we have

$$|f(t)g(t)| \leq \frac{|f(t)|^p}{p} + \frac{|g(t)|^q}{q}.$$

Thus

$$\begin{aligned} \int_a^b |f(t)g(t)| dt &\leq \int_a^b \left(\frac{|f(t)|^p}{p} + \frac{|g(t)|^q}{q} \right) dt \\ &= \frac{1}{p} + \frac{1}{q} = 1 \\ &= \left(\int_a^b |f(t)|^p dt \right)^{\frac{1}{p}} \left(\int_a^b |g(t)|^q dt \right)^{\frac{1}{q}} \end{aligned}$$


as required.

Theorem 31 (Minkowski's Inequality v2)


Let $1 < p < \infty$. If $f, g \in C[a, b]$, then

$$\left(\int_a^b |(f+g)(t)|^p dt \right)^{\frac{1}{p}} \leq \left(\int_a^b |f(t)|^p dt \right)^{\frac{1}{p}} + \left(\int_a^b |g(t)|^p dt \right)^{\frac{1}{p}}.$$

Proof

The proof is similar to the one we had in  Theorem 29; if $\forall x \in [a, b]$, either $f(x) = 0$ or $g(x) = 0$, then the inequality holds trivially so. Thus we may assume that $\forall x \in [a, b]$, $f(x) \neq 0 \neq g(x)$.

Now, notice that by (regular) Triangle Inequality and, later on,

 Theorem 30,

$$\begin{aligned} &\int_a^b |(f+g)(t)|^p dt \\ &= \int_a^b |(f+g)(t)| |(f+g)(t)|^{p-1} dt \\ &\leq \int_a^b |f(t)| |(f+g)(t)|^{p-1} dt + \int_a^b |g(t)| |(f+g)(t)|^{p-1} dt \end{aligned}$$

$$\begin{aligned} &\leq \left(\int_a^b |f(t)|^p dt \right)^{\frac{1}{p}} \left(\int_a^b |(f+g)(t)|^{q(p-1)} dt \right)^{\frac{1}{q}} \\ &\quad + \left(\int_a^b |g(t)|^p dt \right)^{\frac{1}{p}} \left(\int_a^b |(f+g)(t)|^{q(p-1)} dt \right)^{\frac{1}{q}} \\ &= \left[\left(\int_a^b |f(t)|^p dt \right)^{\frac{1}{p}} + \left(\int_a^b |g(t)|^p dt \right)^{\frac{1}{p}} \right] \\ &\quad \cdot \left(\int_a^b |(f+g)(t)|^p dt \right)^{\frac{1}{q}} \end{aligned}$$

where we note that $\frac{1}{p} + \frac{1}{q} = 1 \implies p = q(p-1)$. Consequently, since $\frac{1}{p} = 1 - \frac{1}{q}$,

$$\left(\int_a^b |(f+g)(t)|^p dt \right)^{\frac{1}{p}} \leq \left(\int_a^b |f(t)|^p dt \right)^{\frac{1}{p}} \cdot \left(\int_a^b |g(t)|^p dt \right)^{\frac{1}{p}},$$

as required.

This shows that our definition of $\|\cdot\|_p$ on $C[a, b]$ is indeed a norm, and so $(C[a, b], \|\cdot\|_p)$ is a normed linear space.


Example 11.1.6 (Bounded Operator)

Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be normed linear spaces. Let $T : X \rightarrow Y$ be linear. Define

$$\|T\| = \sup\{\|T_X\|_Y \mid \|x\|_X < 1\}.$$

We say that T is bounded if $\|T\| < \infty$. Let

$$B(X, Y) = \{T : X \rightarrow Y \mid T \text{ is bounded}\}.$$

In the next lecture, we shall show that $(B(X, Y), \|\cdot\|)$ is a normed linear space. 

QUESTION: Consider the transformation $\begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$. What is

a norm $\left\| \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix} \right\|$ that works?

Exercise 11.1.1

Show that there exists an injection from $(C[a, b], \|\cdot\|_2)$ to $\ell_2(\mathbb{N})$. Note that this does not work for $p \geq 3$.

12 Lecture 12 Oct 03rd

12.1 Introduction to Metric Spaces (Continued 3)

Example 12.1.1 (Bounded Operator)

Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be normed linear spaces. Let $T : X \rightarrow Y$ be linear. Define

$$\|T\| = \sup\{\|T(x)\|_Y \mid \|x\|_X < 1\}.$$

We say that T is bounded if $\|T\| < \infty$. Let

$$B(X, Y) = \{T : X \rightarrow Y \mid T \text{ is bounded}\}.$$

To show that $B(X, Y)$ is a normed linear space, let $S, T \in B(X, Y)$, and let $\|x\|_X \leq 1$. Then


$$\begin{aligned} \|(S+T)(x)\|_Y &= \|S(x) + T(x)\|_Y \\ &\leq \|S(x)\|_Y + \|T(x)\|_Y \quad \because \|\cdot\|_Y \text{ is a norm} \\ &\leq \|S\| + \|T\| \end{aligned}$$

and so $S+T \in B(X, Y)$ and $\|S+T\| \leq \|S\| + \|T\|$. For $\alpha \in \mathbb{R}$, we have

$$\begin{aligned} \|\alpha S\| &= \sup\{\|(\alpha S)(x)\|_Y \mid \|x\|_X \leq 1\} \\ &= |\alpha| \sup\{\|S(x)\|_Y \mid \|x\|_X \leq 1\} \quad \because \|\cdot\|_Y \text{ is a norm} \\ &= |\alpha| \|S\|. \end{aligned}$$

So $(\alpha S) \in B(X, Y)$ and $\|\alpha S\| = |\alpha| \|S\|$. It is clear that due to $\|\cdot\|_Y$ being a norm, and so $\|\cdot\|$ is also positive definite. Thus $B(X, Y)$ is a

In this example, we look at how we can apply a translation of norms from X to Y that preserves the norm.

normed linear space as claimed. 

12.2 Topology on Metric Spaces

Definition 34 (Open & Closed)

Let (X, d) be a metric space. If $x_0 \in X$, then

$$B(x_0, \varepsilon) = \{y \in X \mid d(x, y) < \varepsilon\}$$

is called the **open ball** centered at x_0 with radius $\varepsilon > 0$.

$$B[x_0, \varepsilon] = \{y \in X \mid d(x, y) \leq \varepsilon\}$$

is called the **closed ball** centered at x_0 with radius $\varepsilon > 0$.

We say that $U \subset X$ is **open** if

$$\forall x \in U \exists \varepsilon_0 > 0 \quad B(x_0, \varepsilon_0) \subset U.$$

We say that $F \subset X$ is **closed** if F^C is open.

Proposition 32 (Properties of Open Sets)

Let (X, d) be a metric space.

1. X, \emptyset are open,
2. If $\{U_\alpha\}_{\alpha \in I}$ is a collection of open sets, then $U = \bigcup_{\alpha \in I} U_\alpha$ is open.
3. If $\{U_1, \dots, U_n\}$ is a collection of open sets, then $U = \bigcap_{i=1}^n U_i$ is open.

Proof

1. If $x_0 \in X$, then $B(x_0, 1) \subseteq X$, and so X is open. The empty set is open vacuously so.
2. Let $U = \bigcup_{\alpha \in I} U_\alpha$ and $x_0 \in U$. Then $\exists \alpha_0 \in I$ such that $x_0 \in U_{\alpha_0}$.

Then $\exists \varepsilon_0 > 0$ such that

$$B(x_0, \varepsilon_0) \subset U_{\alpha_0} \subset U.$$

3. Let $x_0 \in U = \bigcap_{i=1}^n U_i$. Then for each $i \in \{1, \dots, n\}$, $\exists \varepsilon_i > 0$ such that $B(x_0, \varepsilon_i) \subset U_i$. Let

$$\varepsilon_0 = \min\{\varepsilon_1, \dots, \varepsilon_n\}.$$

Then we have that $\forall i \in \{1, \dots, n\}$, $\varepsilon_0 \leq \varepsilon_i$. Thus

$$B(x_0, \varepsilon_0) \subset B(x_0, \varepsilon_i) \subset U_i$$


for each i . Therefore $B(x_0, \varepsilon_0) \subset U$.

Corollary 33 (Properties of Closed Sets)

Let (X, d) be a metric space.

1. X, \emptyset are closed.
2. If $\{F_\alpha\}_{\alpha \in I}$ is a collection of closed sets, then $F = \bigcap_{\alpha \in I} F_\alpha$ is closed.
3. If $\{F_1, \dots, F_n\}$ is a collection of closed sets, then $F = \bigcup_{i=1}^n F_i$ is closed.

Proof

The proof follows from De Morgan's Laws,  Proposition 32, and by taking set complements.


Example 12.2.1

Let X be any set and d the discrete metric

$$d(x, y) = \begin{cases} 1 & x \neq y \\ 0 & x = y \end{cases}$$

We want to know what sets are open on X under d . Notice that any

Exercise 12.2.1



Write out the full proof for  Corollary 33 as an exercise.

set of just a singleton is open, since

$$B\left(x_0, \frac{1}{2}\right) \subset X.$$

Consequently, any $A \in \mathcal{P}(X)$ is an arbitrary union of open sets, i.e.

$$A = \bigcup_{x \in A} \{x\}.$$

Thus by  Proposition 32, A is open. 

Note 12.2.1

On \mathbb{R} , only \emptyset and \mathbb{R} itself are both open and closed. This can be proven using the *Intermediate Value Theorem*.

Definition 35 (Topology)

Given any X , a set $\tau \subset \mathcal{P}(X)$ is called a **topology** on X is

1. $X, \emptyset \in \tau$
2. If $\{U_\alpha\}_{\alpha \in I}$ such that for each $\alpha \in I$, $U_\alpha \in \tau$, then

$$U = \bigcup_{\alpha \in I} U_\alpha \in \tau.$$

3. If $\{U_1, \dots, U_n\}$ such that $U_i \in \tau$ for each $i \in \{1, \dots, n\}$, then

$$U = \bigcap_{i=1}^n U_i \in \tau.$$

If (X, d) is a metric space, then

$$\tau_d = \{U \subset X \mid U \text{ open in } (X, d)\}$$

is called a **metric topology**, or **d -topology**, associated with the metric d .

We call (X, τ) a **topological space**.

Example 12.2.2

Given X ,

1. $\mathcal{P}(X)$ is a topology on X , and it is called the **discrete topology**;
2. $\{\emptyset, X\}$ is a topology on X , and it is called the **indiscrete topology**.



13 Lecture 13 Oct 05th

13.1 Topology on Metric Spaces (Continued)

Theorem 34 (Open Balls are Open)

1. $B(x_0, \varepsilon)$ is open.
2. $B[x_0, \varepsilon]$ is closed.
3. Every open set is the union of open balls.
4. $\forall x \in X, \{x\}$ is closed.

Proof

1. Consider $x \in B(x_0, \varepsilon)$ and let $r = d(x, x_0)$.

Let $\alpha = \varepsilon - r$. Assume that $y \in B(x, \alpha)$. By the Triangle Inequality,

$$d(x_0, y) \leq d(x_0, x) + d(x, y) < r + \alpha = \varepsilon.$$

2. Let $y \in B[x_0, \varepsilon]^c$, and let $r = d(x_0, y)$.

Let $\alpha = r - \varepsilon$. Assume $z \in B(y, \alpha)$, and suppose, for contradiction, that $z \in B[x_0, \varepsilon]$. Then

$$r = d(x_0, y) \leq d(x_0, z) + d(z, y) < \varepsilon + \alpha = r,$$

but $r < r$ contradicts the fact that $r = r$.




Figure 13.1: Idea of proof for 1. in \mathbb{R}^2 .




Figure 13.2: Idea of proof for 2. in \mathbb{R}^2 .

3. Let $U \subset X$ be open. $\forall x \in U$, let $\varepsilon_x > 0$ be such that $B(x, \varepsilon_x) \subset U$.

Then

$$U = \bigcup_{x \in U} B(x, \varepsilon_x).$$


4. Let $y \in X$ such that $y \neq x$. Let $r = d(y, x)$. Then $x \notin B(y, \frac{r}{2})$, and so

$$B(y, \frac{r}{2}) \subset \{x\}^c. \quad \square$$

Example 13.1.1 (Open Intervals are Open)

Let $X = \mathbb{R}$, and $d(x, y) = |x - y|$, the standard metric. Let $I = (a, b)$, for some $a, b \in \mathbb{R} \cup \{\pm\infty\}$. Let $x \in I$. Now let

$$\varepsilon = \min\{1, |x - a|, |x - b|\}.$$

Then, clearly so, $B(x, \varepsilon) \subset I$. 

If $U \subset \mathbb{R}$ is open, and if we define \sim on U by $x \sim y$ iff $(x, y), (y, x) \subset U$. This is what we did in Q1. We proved that \sim is an equivalence relation. Let $I_x = [x]$ be the interval defined by \sim . We proved that I_x is an open interval.

Consequently, if we have U being open in \mathbb{R} , then U can be expressed as the union of a countable collection $\{I_\alpha, \alpha \in I\}$ of open intervals, which are pairwise disjoint.

QUESTION: Given $U = \{(x, y) \mid |x|, |y| < 1\}$, can we do the same as above, i.e. can we use a countable collection of disjoint open sets to express U , or, in other words, cover U ?

Example 13.1.2 (Cantor Set)

Let's consider the closed interval $[0, 1]$, of which we shall label as P_0 .




Figure 13.3: Cantor set showing up to $n = 2$, with the excluded interval in $n = 3$ shown.

Define P_1 by removing an open interval of length $\frac{1}{3}$ sitting in the

middle of P_0 , i.e.

$$P_1 = [0, 1] \setminus \left(\frac{1}{3}, \frac{2}{3}\right) = \left[0, \frac{1}{3}\right] \cup \left[\frac{2}{3}, 1\right]$$

Define P_2 by removing an open interval of length $\frac{1}{3^2}$ sitting in the middle of each of the 2 closed intervals in P_1 , ie.


$$P_2 = \left[0, \frac{1}{9}\right] \cup \left[\frac{2}{9}, \frac{1}{3}\right] \cup \left[\frac{2}{3}, \frac{7}{9}\right] \cup \left[\frac{8}{9}, 1\right]$$

Recursively so, define P_{n+1} by removing an open interval of length $\frac{1}{3^{n+1}}$ sitting in the middle of each of the 2^n closed intervals in P_n .

Let P , the **Cantor Set** (or **Cantor Ternary Set**), be defined as

$$P = \bigcap_{n=0}^{\infty} P_n.$$

The following are some properties of P :

1. P is closed, since it is closed under an arbitrary number of closed sets (see  Corollary 33).


2. We have

$$x \in P \iff x = \sum_{i=1}^{\infty} \frac{a_i}{3^i}$$

where $a_n = 0, 2$. In other words, every element of P is a ternary number.

3. $|P| = 2^{\aleph_0} = c$.

4. P_n does not contain any interval of length greater than or equal to $\frac{1}{3^n}$.

5. Consequently, the length of P is 0. 

14 Lecture 14 Oct 12th

14.1 Topology on Metric Spaces (Continued 2)

Definition 36 (Closure)

Let $A \subseteq (X, d)$. We define the **closure** \bar{A} of A to be

$$\bar{A} = \cap \{F \subset X \mid F \text{ is closed, } A \subset F\}.$$

\bar{A} is the smallest closed set that contains A .

Definition 37 (Interior)

Let $A \subseteq (X, d)$. We define the **interior** A° of A to be

$$A^\circ = \cup \{U \subset X \mid U \text{ is open, } U \subset A\}.$$

A° is the largest open set contained in A .

Remark 14.1.1

We have that

$$A^\circ \subset A \subset \bar{A}$$

Definition 38 (Neighbourhood)

We say that a set A is a **neighbourhood** of a point $x \in X$ if $x \in A^\circ$.¹

¹ A neighbourhood is **not necessarily open**; the definition applies to elements in the interior after all.

“ Note 14.1.1

A is a neighbourhood of $x \in X$ if and only if $\exists \varepsilon > 0$ such that $B(x, \varepsilon) \subset A$.

📖 Definition 39 (Boundary Point)

Given $A \subset (X, d)$, a point x is called a **boundary point** for A if

$$\forall \varepsilon > 0 \quad B(x, \varepsilon) \cap A \neq \emptyset \wedge B(x, \varepsilon) \cap A^c \neq \emptyset.$$

We denote the collection of all boundary points of A by $\text{bdy}(A)$.

💧 Proposition 35 (Closed Sets Include Its Boundary Points)

Let (X, d) be a metric space and $A \subset X$. Then A is closed \iff
 $\text{bdy}(A) \subset A$.

✎ Proof

(1) \implies (2): Suppose $x \in A^c$, which is open. Then $\exists \varepsilon > 0$ such that $B(x, \varepsilon) \subset A^c$. Then $x \notin \text{bdy}(A)$, i.e. $\text{bdy}(A) \subset A$.²

(2) \implies (3): ³Let $x \in A^c$. Then, by assumption, $x \notin \text{bdy}(A)$. Then $\exists \varepsilon > 0$ such that either $B(x, \varepsilon) \subset A$ or $B(x, \varepsilon) \subset A^c$. But since $x \notin A$, we must have $B(x, \varepsilon) \subset A^c$, i.e. A^c is open.

² The idea of this proof is to show that it is impossible for the boundary to be in A^c .

³ To show that A is closed, we should show that A^c is open.

💧 Proposition 36 (Closures include the Boundary Points of a Set)

Given $A \subset (X, d)$, we have $\bar{A} = A \cup \text{bdy}(A)$.

 **Proof**

By definition, $A \subseteq \bar{A}$, so it suffices to show that $\text{bdy}(A) \subset \bar{A}$ to show that $A \cup \text{bdy}(A) \subseteq \bar{A}$.


⁴Assume that $x \notin \bar{A}$, i.e. $x \in \bar{A}^C$, which is open since \bar{A} is closed by definition. Then $\exists \varepsilon > 0$ such that $B(x, \varepsilon) \subset \bar{A}^C$. Since $x \notin A \subset \bar{A}$, we have that $B(x, \varepsilon) \cap A = \emptyset$, i.e. $x \notin \text{bdy}(A)$. Therefore $\text{bdy}(A) \subset \bar{A}$, and so $A \cup \text{bdy}(A) \subseteq \bar{A}$ as claimed.

⁴ Here, we employ the same proof as the previous proposition.

⁵Let $x \in \text{bdy}(A \cup \text{bdy}(A))$. Then $\forall \varepsilon > 0$, we have

$$B(x, \varepsilon) \cap (A \cup \text{bdy}(A)) \neq \emptyset \quad (14.1)$$

$$\begin{aligned} &\wedge \\ B(x, \varepsilon) \cap (A \cup \text{bdy}(A))^C &\neq \emptyset. \end{aligned} \quad (14.2)$$

⁵ For this part, if we can show that $A \cup \text{bdy}(A)$ is closed, then by definition, $\bar{A} \subseteq A \cup \text{bdy}(A)$ since \bar{A} is the smallest such set that contains A . To show that $A \cup \text{bdy}(A)$ is closed, we can either show that $(A \cup \text{bdy}(A))^C$ is open, or use  Proposition 35 to show that $\text{bdy}(A \cup \text{bdy}(A)) \subset (A \cup \text{bdy}(A))$. We shall show for the more complicated expression.

Note that by De Morgan's Laws, we have that

$$(A \cup \text{bdy}(A))^C = A^C \cap \text{bdy}(A)^C.$$

Then (14.2) would be

$$B(x, \varepsilon) \cap A^C \cap \text{bdy}(A)^C \neq \emptyset,$$

and so

$$B(x, \varepsilon) \cap A^C \neq \emptyset \quad (14.3)$$

$$\begin{aligned} &\wedge \\ B(x, \varepsilon) \cap \text{bdy}(A)^C &\neq \emptyset. \end{aligned} \quad (14.4)$$

From (14.1), we have

$$B(x, \varepsilon) \cap A \neq \emptyset \vee B(x, \varepsilon) \cap \text{bdy}(A) \neq \emptyset.$$


If $B(x, \varepsilon) \cap A \neq \emptyset$, then \therefore (14.3), $x \in \text{bdy}(A)$, and so

$$\text{bdy}(A \cup \text{bdy}(A)) \subseteq (A \cup \text{bdy}(A)). \quad (\dagger)$$

If $B(x, \varepsilon) \cap \text{bdy}(A) \neq \emptyset$, let $z \in B(x, \varepsilon) \cap \text{bdy}(A)$. $\therefore z \in B(x, \varepsilon)$, let $r = d(x, z)$, and $\alpha = \varepsilon - r > 0$. Let $z_0 \in B(z, \alpha)$. Then by the


Triangle Inequality

$$d(x, z_0) \leq d(x, z) + d(z, z_0) < r + \alpha = \varepsilon.$$

Thus $z_0 \in B(x, \varepsilon) \implies (B(z, \alpha) \subseteq B(x, \varepsilon))$. Then $\because z \in \text{bdy}(A)$, we have $B(z, \alpha) \cap A \neq \emptyset$, and so $B(x, \varepsilon) \cap A \neq \emptyset$. Then we can just follow the argument we did in (†) and arrive at the same conclusion. Consequently, by  Proposition 35, $A \cup \text{bdy}(A)$ is closed as claimed.


Example 14.1.1


Let $X = \mathbb{R}$ and $A = [0, 1)$. We have that

- $\text{bdy}(A) = \{0, 1\}$;
- $A^\circ = (0, 1)$; and
- $\bar{A} = [0, 1]$. 

Example 14.1.2

Let $X = \mathbb{R}$ and $A = \mathbb{Q}$. We have that

- $\text{bdy}(A) = \mathbb{R}$ since every open ball around $a \in A$ will always contain elements in \mathbb{Q} and \mathbb{Q}^c ;
- $A^\circ = \emptyset$ since $A^\circ = A \setminus \text{bdy}(A)$; and
- $\bar{A} = \mathbb{R}$ since $\bar{A} = A \cup \text{bdy}(A)$. 


 **Definition 40 (Separable)**

A metric space (X, d) is *separable* if there exists a countable set $A \subset X$ such that $\bar{A} = X$, and call the metric space *non-separable* otherwise.


Example 14.1.3

Every finite metric space (X, d) is separable.

This is true since every subset A of X is countable since X itself


is countable. Consequently, if we pick A to be a subset that takes every other element in X , then it is clear that $\bar{A} = X$, and so (X, d) is separable. 

Example 14.1.4

\mathbb{R} is separable as shown in Example 14.1.2.⁶ 

6

Example 14.1.5

\mathbb{R}^n is separable if d_p for all $1 \leq p \leq \infty$. We can apply the same argument that we had for Example 14.1.2 and apply it component-wise. Consequently, $\overline{\mathbb{Q}^n} = \mathbb{R}^n$. In other words, for any $(x_1, \dots, x_n) \in (\mathbb{R}^n, d_p)$, we can pick a $(r_1, \dots, r_n) \in \mathbb{Q}^n$ that is as close to (x_1, \dots, x_n) as possible. 


Exercise 14.1.1

Prove that $\bar{\mathbb{Q}} = \mathbb{R}$ using the Archimedean Property of \mathbb{R} .

Remark 14.1.2

Notice that

$$\bar{A} = X \iff \forall x \in X \forall \varepsilon > 0 B(x, \varepsilon) \cap A \neq \emptyset. \quad \text{💬}$$

 **Definition 41 (Dense)**

A is *dense* in (X, d) if $\bar{A} = X$. Equivalently, A is dense if for every open set $W \subset X$, $W \cap A \neq \emptyset$.

QUESTION: Is $(\ell_1, \|\cdot\|_1)$ separable? Is $(\ell_\infty, \|\cdot\|_\infty)$ separable?

Recall Example 10.1.1.

15 Lecture 15 Oct 15th

15.1 Topology on Metric Spaces (Continued 3)

Definition 42 (Limit Points)

Let (X, d) be a metric space, and $A \subset X$. We say that x_0 is a **limit point** for A if for any neighbourhood of x_0 , we have that

$$N \cap (A \setminus \{x_0\}) \neq \emptyset.$$

Equivalently, $\forall \varepsilon > 0, \exists x \in A$, where $x \neq x_0$, such that $x \in B(x_0, \varepsilon)$.¹
We sometimes call limit points as **cluster points**. We denote the set of limit points of A as $\text{Lim}(A) \subset X$ ²

¹ This also means that $B(x_0, \varepsilon)$ must have infinitely many points close to x_0 , for otherwise, we would be able to find some $\varepsilon > \varepsilon_0 > 0$ such that $B(x_0, \varepsilon_0) \cap A = \emptyset$.

² Note that the set of limit points is not necessarily a subset of A .

Example 15.1.1

Let $X = \mathbb{R}$, and $A = [0, 1) \subset \mathbb{R}$. We have that

$$\text{Lim}[0, 1) = [0, 1].$$

Example 15.1.2

Let $X = \mathbb{R}$ and $A = \mathbb{N} \subset \mathbb{R}$. Since $\forall n \in \mathbb{N}, \exists \varepsilon = \frac{1}{2}$ such that $\forall m \in \mathbb{N} \setminus \{n\}$, we have that $m \notin B\left(n, \frac{1}{2}\right)$, we have

$$\text{Lim } \mathbb{N} = \emptyset.$$

💧 Proposition 37 (Closed Sets Include Its Limit Points)

Let $A \subset (X, d)$. Then

1. A is closed $\iff \text{Lim}(A) \subset A$;
2. $\bar{A} = A \cup \text{Lim}(A)$.

Proof

1. For the (\implies) direction, suppose A is closed. ³Let $x_0 \in A^C$. Then $\exists \varepsilon > 0$ such that $B(x_0, \varepsilon) \cap A = \emptyset$. Thus, by definition, we have that $x_0 \notin \text{Lim}(A)$ ⁴. Therefore, $\text{Lim}(A) \subset A$.


For the (\impliedby) direction, suppose $\text{Lim}(A) \subset A$. Let $x_0 \in A^C$. Then $x_0 \notin \text{Lim}(A)$, which means that $\exists \varepsilon > 0$ such that $B(x_0, \varepsilon) \cap A = \emptyset$, i.e. $B(x_0, \varepsilon) \subset A^C$. Thus A is closed.

2. ⁵It is clear that $A \subset \bar{A}$. Let $x_0 \in \bar{A}^C$. Then $\exists \varepsilon > 0$ such that $B(x_0, \varepsilon) \subset \bar{A}^C$. In particular, we have that $B(x_0, \varepsilon) \cap A = \emptyset$, i.e. $x_0 \notin \text{Lim}(A)$. Thus $\text{Lim}(A) \subset \bar{A}$.

Again, it suffices to show that $A \cup \text{Lim}(A)$ is closed to CTP. Let $x_0 \in (A \cup \text{Lim}(A))^C$ ⁶. Then $\exists \varepsilon > 0$ such that $B(x_0, \varepsilon) \cap A = \emptyset$. If $z \in \text{Lim}(A)$ and $z \in B(x_0, \varepsilon)$, then we have $B(x_0, \varepsilon)$ is a neighbourhood of z , and so we must have that $B(x_0, \varepsilon) \cap A \neq \emptyset$, which is a contradiction. Thus $(A \cup \text{Lim}(A))^C$ is open, and so $A \cup \text{Lim}(A)$ is closed, as required. \square

³ This uses a reversed way of thinking: if we want to show that $\text{Lim}(A) \subset A$, then instead of trying to directly show the containment, we show that all elements in A^C are in fact not limit points due to A being closed.

⁴ Notice there that there are no elements in A that are **close to** x_0 , and so it's not a limit point.

⁵ This proof is similar to that of  Proposition 36.


⁶ It is clear by De Morgan's Law that $x_0 \in A^C$ and $x_0 \notin \text{Lim}(A)$, which implies that $\text{Lim}(A) \subset A$. But this does not give us a clear geometrical picture of the notion.

💧 Proposition 38 (Mixing the notions)

Let $A \subseteq B \subseteq (X, d)$.

1. $\bar{A} \subseteq \bar{B}$;
2. $A^\circ \subset B^\circ$;
3. $A^\circ = A \setminus \text{bdy}(A)$;
4. $\text{bdy}(A) = \text{bdy}(A^C)$;


Exercise 15.1.1

Prove  Proposition 38.

5. $A^\circ = (\overline{A^c})^c.$

 **Proof**

1. It is clear that $A \subset B \subset \overline{B}$. Suppose $\text{Lim}(A)$ is not a subset of \overline{B} . Then $\exists x \in \text{Lim}(A) \setminus \overline{B}$, i.e. $x \in \overline{B}^c$. Since \overline{B} is closed, B^c is open and so $\exists \varepsilon > 0$ such that $B(x, \varepsilon) \subset B^c$. Since $x \in \text{Lim}(A)$, $\exists a \in A$ such that $a \in B(x, \varepsilon) \subset B^c$, but $A \subset B$, a contradiction. Thus $\text{Lim}(A) \subset \overline{B}$.
2. $a \in A^\circ \implies \exists \varepsilon > 0 B(a, \varepsilon) \subset A \subset B \implies a \in B^\circ \dashv$
3. $x \in A \setminus \text{bdy}(A) \implies \exists \varepsilon > 0 B(x, \varepsilon) \cap A^c = \emptyset \implies x \in A^\circ \dashv$
 $x \in A^\circ \implies \exists \varepsilon_0 > 0 B(x, \varepsilon_0) \subset A$
 Sps $x \in \text{bdy}(A)$. Then $\forall \varepsilon > 0 B(x, \varepsilon) \cap A^c \neq \emptyset \implies B(x, \varepsilon_0) \cap A^c = \emptyset \nmid B(x, \varepsilon_0) \subset A \dashv$
4. $x \in \text{bdy}(A) \implies \forall \varepsilon > 0 B(x, \varepsilon) \cap A \neq \emptyset \wedge B(x, \varepsilon) \cap A^c \neq \emptyset$
 $x \notin \text{bdy}(A^c) \implies \exists \varepsilon_0 > 0 B(x, \varepsilon_0) \cap A = \emptyset \vee B(x, \varepsilon_0) \cap A^c = \emptyset$
 But $B(x, \varepsilon_0) \cap A = \emptyset \nmid \forall \varepsilon > 0 B(x, \varepsilon) \cap A^c \neq \emptyset$
 and $B(x, \varepsilon_0) \cap A^c = \emptyset \nmid \forall \varepsilon > 0 B(x, \varepsilon) \cap A^c \neq \emptyset$
 $\implies x \in \text{bdy}(A^c) \dashv$. The converse is a similar argument.
5. $(\overline{A^c})^c = (A^c \cup \text{bdy}(A^c))^c = A \cap \text{bdy}(A)^c = A \setminus \text{bdy}(A) = A^\circ \dashv$

 **Proposition 39 (More on Closures and Interiors)**

Let $A, B \subseteq (X, d)$.

1. $\overline{A \cup B} = \overline{A} \cup \overline{B}$;
2. $(A \cap B)^\circ = A^\circ \cap B^\circ$.

Exercise 15.1.2

Prove Item 2 for  Proposition 39.

 **Proof**

1. We have that $A \subseteq \bar{A}$ and $B \subseteq \bar{B}$, so $A \cup B \subseteq \bar{A} \cup \bar{B}$. Since $\overline{A \cup B}$ is closed, we must have that $\overline{A \cup B} \subseteq \bar{A} \cup \bar{B}$. Similarly so, we have

$$A \subseteq A \cup B \implies \bar{A} \subseteq \overline{A \cup B}$$

$$B \subseteq A \cup B \implies \bar{B} \subseteq \overline{A \cup B}$$

and so $\bar{A} \cup \bar{B} \subseteq \overline{A \cup B}$.

2. Since $A^\circ \subseteq A$ and $B^\circ \subseteq B$, and $A^\circ \cap B^\circ$ is open, we must have that $A^\circ \cap B^\circ \subseteq (A \cap B)^\circ$. On the other hand, since $(A \cap B)^\circ \subseteq A^\circ$ and $(A \cap B)^\circ \subseteq B^\circ$, we have that $(A \cap B)^\circ \subseteq A^\circ \cap B^\circ$.

QUESTION: Is $\overline{A \cap B} = \bar{A} \cap \bar{B}$? No.

Example 15.1.3

Let $X = \mathbb{R}$, $A = \mathbb{Q}$ and $B = \mathbb{Q}^c$. We know that $\bar{A} = \mathbb{R} = \bar{B}$. But, observe that

$$\overline{A \cap B} = \emptyset \text{ while } \bar{A} \cap \bar{B} = \mathbb{R}. \quad \blacktriangleright$$

However, we do have that $\overline{A \cap B} \subseteq \bar{A} \cap \bar{B}$.

QUESTION: Given (X, d) a metric space, is

$$B(x_0, \varepsilon) = B[x_0, \varepsilon]$$

true? Again, no.

Example 15.1.4

Let X be a set with $|X| \geq 2$, and d the discrete metric. We have that

$$B(x_0, 1) = \{x_0\} \text{ but } B[x_0, 1] = X. \quad \blacktriangleright$$

Definition 43 (Convergence)

Given a sequence $\{x_n\} \subset (X, d)$ and $x_0 \in X$, we say that the sequence *converges* to x_0 if

$$\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall n \geq N_0 d(x_n, x_0) < \varepsilon.$$

This is equivalent to saying that the sequence $\{d(x_n, x_0)\}$ converges to 0 in X . We denote this by

$$x_0 = \lim_{n \rightarrow \infty} x_n \text{ or } x_n \rightarrow x_0.$$

If no such x_0 exists, we say that the sequence *diverges*.

Theorem 40 (Uniqueness of Limits of Sequences)

If $\{x_n\}$ is a sequence in (X, d) with $x_n \rightarrow x_0$ and $x_n \rightarrow y_0$, then $x_0 = y_0$.

Proof

$$x_0 \neq y_0 \implies \exists \varepsilon = d(x_0, y_0) \implies B(x_0, \frac{\varepsilon}{2}) \cap B(y_0, \frac{\varepsilon}{2}) = \emptyset$$

However, $\exists N_0 \in \mathbb{N} \forall n \geq N_0$

$$x_n \in B(x_0, \frac{\varepsilon}{2}) \wedge x_n \in B(y_0, \frac{\varepsilon}{2})$$

which is impossible. Thus $x_0 = y_0$.





Figure 15.1: A geometric representation of the proof for  Theorem 40.

16 Lecture 16 Oct 17th

16.1 Convergences of Sequences (Continued)

Example 16.1.1

Let $X = \mathbb{R}^n$, $d = d_p$, for $1 \leq p \leq \infty$, and $\vec{x}_k = \{(x_{k,1}, x_{k,2}, \dots, x_{k,n})\}$.

Claim :

$$\vec{X}_k \xrightarrow{\ell_p} \vec{x}_0 = (x_{0,1}, x_{0,2}, \dots, x_{0,n}) \iff \forall j \in \{1, \dots, n\} x_{k,j} \rightarrow x_{0,j}.$$

Note : In general, we have

$$|x_{k,j} - x_{0,j}| \leq \|\vec{x}_k - \vec{x}_0\|_p$$

So it is clear that the (\implies) direction is true, i.e.

$$\vec{X}_k \rightarrow \vec{x}_0 \implies \forall j \in \{1, \dots, n\} x_{k,j} \rightarrow x_{0,j}.$$

For the other direction, we look at the different p 's to see how it works differently: in all cases, assume that $x_{k,j} \rightarrow x_{0,j}$ for all j , and that $\varepsilon > 0$

$p = \infty$: we have that $\exists k_0 \in \mathbb{N}$ such that $\forall k \geq k_0$,

$$|x_{k,j} - x_{0,j}| < \varepsilon \text{ for } j \in \{1, \dots, n\},$$

and so

$$\|\vec{x}_k - \vec{x}_0\|_\infty = \max \left\{ |x_{k,j} - x_{0,j}| : 1 \leq j \leq n \right\} < \varepsilon.$$

$p = 1$: if we assume that for each j ,

$$|x_{k,j} - x_{0,j}| < \frac{\varepsilon}{n},$$

then


$$\|\vec{x}_k - \vec{x}_0\|_1 = \sum_{j=1}^n |x_{k,j} - x_{0,j}| < \sum_{j=1}^n \frac{\varepsilon}{n} = \varepsilon.$$

$1 < p < \infty$: this time, we assume that for each j ,

$$|x_{k,j} - x_{0,j}| < \frac{\varepsilon}{\sqrt[p]{n}}.$$

Then

$$\|\vec{x}_k - \vec{x}_0\|_p = \left(\sum_{j=1}^n |x_{k,j} - x_{0,j}|^p \right)^{\frac{1}{p}} < \left(\sum_{j=1}^n \left(\frac{\varepsilon}{\sqrt[p]{n}} \right)^p \right)^{\frac{1}{p}} = \varepsilon.$$

This completes the proof of our claim. 

Example 16.1.2

Let $X = (C[a, b], \|\cdot\|_\infty)$. Then

$$f_n \rightarrow f \iff \|f_n - f\|_\infty \rightarrow 0.$$

Notice that for the (\implies) direction,¹

$$\begin{aligned} (\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall n \geq N_0 |f_n - f| < \varepsilon) \\ \implies \|f_n - f\|_\infty = \max\{|f_n(x) - f(x)| : x \in [a, b]\} < \varepsilon. \end{aligned}$$

¹ Note that this is **uniform convergence**, which implies **pointwise convergence**.

The (\impliedby) direction is easy, since

$$|f_n(x) - f(x)| \leq \max\{|f_n(x) - f(x)| : x \in [a, b]\} < \varepsilon. \quad \img alt="arrow" data-bbox="600 690 620 710"/>$$

Theorem 41 (Sequential Characterizations of Limit Points, Boundaries, and Closedness)

Given $A \subset (X, d)$,

$$1. \ x_0 \in \text{Lim}(A) \iff \exists \{x_n\} \subset A \ (x_n \neq x_0) \wedge (x_n \rightarrow x_0);$$

2. $x_0 \in \text{bdy}(A) \iff \exists \{x_n\} \subset A, \{y_n\} \subset A^C (x_n \rightarrow x_0) \wedge (y_n \rightarrow x_0)$;
3. A is closed $\iff (\forall \{x_n\} \subset A \ x_n \rightarrow x_0 \in X \implies x_0 \in A)$

 **Proof**

1. $x_0 \in \text{Lim}(A) \implies \forall n \in \mathbb{N} \ x_n \in B\left(x_0, \frac{1}{n}\right) \setminus \{x_0\} \implies d(x_n, x_0) < \frac{1}{n} \implies x_n \rightarrow x_0 \dashv$

$$\begin{aligned} & \{x_n\} \subset A \ (x_n \rightarrow x_0) \wedge (x_n \neq x_0) \implies \\ & \forall \varepsilon > 0 \ \exists N_0 \in \mathbb{N} \ \forall n \geq N_0 \ x_n \in B(x_0, \varepsilon) \dashv \end{aligned}$$

2. $x \in \text{bdy}(A) \implies$
 $(\because \forall \varepsilon > 0 \ A \cap B(x, \varepsilon) \neq \emptyset) \ \exists x_n \in A \cap B\left(x, \frac{1}{n}\right) \wedge$
 $(\because \forall \varepsilon > 0 \ A^C \cap B(x, \varepsilon) \neq \emptyset) \ \exists y_n \in A^C \cap B\left(x, \frac{1}{n}\right)$
 $\implies (\{x_n\} \subset A \wedge x_n \rightarrow x) \wedge (\{y_n\} \subset A^C \wedge y_n \rightarrow x) \dashv$

$$\begin{aligned} & (\{x_n\} \subset A \wedge x_n \rightarrow x) \wedge (\{y_n\} \subset A^C \wedge y_n \rightarrow x) \\ & \implies \forall \varepsilon > 0 \ \exists N_0 \in \mathbb{N} \ \forall n \geq N_0 \ x_n, y_n \in B(x, \varepsilon) \\ & \implies x_0 \in \text{bdy}(A) \dashv \end{aligned}$$

3. Sps A is closed and $(\{x_n\} \subset A) \wedge (x_n \rightarrow x_0 \in X)$.
 $x_0 \in A^C \implies \exists \varepsilon > 0 \ B(x_0, \varepsilon) \subset A^C \implies x_n \notin B(x_0, \varepsilon) \not\rightarrow x_n \rightarrow x_0$
 $\implies x_0 \in A$

Sps A is \neg closed $\implies (\because \text{Proposition 37}) \ \exists x_0 \in \text{Lim}(A) \setminus A$
 $\implies (\because \text{Item 1}) \ \exists \{x_n\} \subset A \ (x_n \neq x_0) \wedge (x_n \rightarrow x_0 \notin A)$, showing that RHS is false \dashv

Example 16.1.3

Let X be a set and d a discrete metric. Then

$$x_n \rightarrow x_0 \iff \exists N \in \mathbb{N} \ \forall n \geq N \ x_n = x_0.$$



Example 16.1.4

Let $c_0 = \{\{x_n\} \mid \lim_{n \rightarrow \infty} x_n = 0\} \subset \ell_\infty$.

Claim : c_0 is closed in ℓ_∞ .

Assume $\vec{x}_k = \{x_{k,j}\}_{j=1}^\infty \in c_0$, and let

$$\vec{x}_k \xrightarrow{\|\cdot\|_\infty} \vec{x}_0 = \{x_{0,j}\}_{j=1}^\infty \in \ell_\infty,$$

i.e.



$$\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall k \geq N_0 \quad \|\vec{x}_k - \vec{x}_0\|_\infty < \frac{\varepsilon}{2}$$

Let $k_0 \geq N_0$. $\because \vec{x}_{k_0} \in c_0, \exists J_0 \in \mathbb{N}$ such that $\forall j \geq J_0$, we have $|x_{k_0,j}| < \frac{\varepsilon}{2}$, and so

$$|x_{0,j}| \leq |x_{k_0,j} - x_{0,j}| + |x_{k_0,j}| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

Thus we have that

$$\lim_{j \rightarrow \infty} x_{0,j} = 0$$

and so $\vec{x}_0 \in c_0$. Therefore, by  Theorem 4.1 Item 3, c_0 is closed in ℓ_∞ . 

Note, however, that $c_{00} \subset \ell_1 \subset c_0$ is not closed. Also ℓ_p is not closed in c_0 .

17 Lecture 17 Oct 19th

17.1 Induced Metric and Topologies

Definition 44 (Induced Metric & Induced Topology)

Given (X, d) and $A \subset X$, we define the *induced metric* d_A on A by

$$d_A : A \times A \rightarrow \mathbb{R}$$

where $d_A(x, y) = d(x, y)$, for all $x, y \in A$, i.e. $d_A = d \upharpoonright_{A \times A}$.

We define τ_A , the *induced topology* on A by

$$\tau_A = \{W \subset A \mid W = U \cap A, U \subset X \text{ is open} \}$$

Note 17.1.1


Note that τ_A is indeed a topology: it is clear that $\emptyset \in \tau_A$. Also, $A \in \tau_A$, since X is open and $A = X \cap A$.

For an arbitrary collection $\{U_\alpha\}_{\alpha \in I} \subset \tau_A$, we know that each $U_\alpha \subset A$, and so $\bigcup_{\alpha \in I} U_\alpha \subset A$. Since each $U_\alpha \in \tau_A$, $\exists F_\alpha \subset X$ that is an open set such that $U_\alpha = F_\alpha \cap A$. Then

$$\bigcup_{\alpha \in I} U_\alpha = \bigcup_{\alpha \in I} F_\alpha \cap A.$$

Thus $\bigcup_{\alpha \in I} U_\alpha \in \tau_A$.

For a finite collection $\{U_1, U_2, \dots, U_n\} \subset \tau_A$, we have that for each U_i ,

$\exists F_i \subset X$ that is open such that $U_i = F_i \cap A$. By  Proposition 32, we have that

$$\bigcap_{i=1}^n F_i \subset X$$

is open, and so

$$\bigcap_{i=1}^n U_i = \bigcap_{i=1}^n F_i \cap A \subset A.$$

Therefore, $\bigcap_{i=1}^n U_i \in \tau_A$.

Theorem 42 (The Metric Topology of a Subset is Its Induced Topology)

We have

$$\tau_A = \tau_{d_A}.$$

Proof

$$\begin{aligned} \subseteq : W \in \tau_A &\implies \exists U \subset X \text{ open such that } W = U \cap A \\ &\implies \forall x_0 \in W \exists \varepsilon > 0 B_X(x_0, \varepsilon) \subset U \\ &\implies B_A(x_0, \varepsilon) = B_X(x_0, \varepsilon) \cap A \subset W \implies W \in \tau_{d_A} \quad \dashv \end{aligned}$$

$$\begin{aligned} \supseteq : W \in \tau_{d_A} &\implies \forall x_0 \in W \exists \varepsilon_x > 0 B_A(x_0, \varepsilon_x) \subset W \\ &\implies W = \bigcup_{x_0 \in W} B_A(x_0, \varepsilon_x) \\ \text{Let } U &= \bigcup_{x_0 \in W} B_X(x_0, \varepsilon_x), \text{ which is open} \\ &\implies W = \bigcup_{x_0 \in W} B_X(x_0, \varepsilon_x) \cap A = U \cap A \\ &\implies W \in \tau_A \quad \dashv \end{aligned}$$

17.2 Continuity on Metric Spaces

Definition 45 (Continuity)

Given metric spaces (X, d_X) , (Y, d_Y) , and $f : X \rightarrow Y$, we say that f is

continuous at x_0 if

$$\forall \varepsilon > 0 \exists \delta > 0 \forall x \in X \ d_X(x, x_0) < \delta \implies d_Y(f(x), f(x_0)) < \varepsilon.$$

Theorem 43 (Continuity and Neighbourhoods)

Given metric spaces (X, d_X) and (Y, d_Y) , and $f : X \rightarrow Y$, then TFAE:

1. f is continuous at $x_0 \in X$;
2. if W is a neighbourhood of $f(x_0) \in Y$, then $f^{-1}(W)$ is a neighbourhood of $x_0 \in X$, where

$$f^{-1}(W) = \{x \in X : f(x) \in W\}.$$





Figure 17.1: Visual representation of  Theorem 43

Proof

(1) \implies (2) : Sps f is continuous at $x_0 \in X$ and W a neighbourhood of $y_0 = f(x_0)$

$$\implies f(x_0) = y_0 \in W^\circ$$

$$\implies \exists \varepsilon > 0 \ B(f(x_0), \varepsilon) \subset W$$

$\because f$ is continuous,

$$\exists \delta > 0 \ \forall x \in X \ x \in B_X(x_0, \delta) \implies d_Y(f(x), f(x_0)) < \varepsilon$$

$$\implies f(x) \in B_Y(y_0, \varepsilon) \subset W$$

$$\implies x \in f^{-1}(W) \implies x_0 \in f^{-1}(W)^\circ \dashv$$

(2) \implies (1) : Sps $f^{-1}(W)$ is a neighbourhood of $x \in X$ for each neighbourhood W of $y_0 = f(x_0)$

$$\implies \forall \varepsilon > 0 \ W = B_Y(f(x_0), \varepsilon) \text{ is a neighbourhood of } f(x_0)$$

$$\implies U = f^{-1}(W) \text{ is a neighbourhood of } x_0 \in X$$

$$\implies x_0 \in U$$

$$\implies \exists \delta > 0 \ B(x_0, \delta) \subset U = f^{-1}(W)$$

$$\implies (d_X(x, x_0) < \delta \implies d_Y(f(x), f(x_0)) < \varepsilon) \dashv$$

Theorem 44 (★ Sequential Characterization of Continuity)

For metric spaces (X, d_X) and (Y, d_Y) , and $f : X \rightarrow Y$, TFAE

1. f is continuous at $x_0 \in X$;
2. $\{x_n\} \subset X \ x_n \xrightarrow{X} x_0 \implies f(x_n) \xrightarrow{Y} f(x_0)$

Proof

(1) \implies (2) : Sps f is continuous at $x_0 \in X$.

$$x_n \rightarrow x_0 \iff$$

$$\forall \varepsilon > 0 \exists \delta > 0 \ x \in B_X(x_0, \delta) \implies f(x) \in B_Y(f(x_0), \varepsilon)$$

$$x_n \rightarrow x_0 \implies \exists N_0 \in \mathbb{N} \ \forall n \geq N_0$$

$$d_X(x_0, x) < \delta \implies x_n \in B_X(x_0, \delta) \implies f(x) \in B_Y(f(x_0), \varepsilon) \dashv$$

(2) \implies (1) (Prove by Contrapositive) : Sps f is \neg continuous at $x_0 \in X$

$$\implies \exists \varepsilon_0 > 0 \ \forall \delta > 0 \ (x_\delta \in B_X(x_0, \delta)) \wedge (f(x_\delta) \notin B_Y(f(x_0), \varepsilon_0))$$

$$\implies \forall n \in \mathbb{N} \ \exists x_n \in B_X\left(x_0, \frac{1}{n}\right) \wedge f(x_n) \notin B_Y(f(x_0), \varepsilon_0)$$

$$\implies x_n \rightarrow x_0 \wedge f(x_n) \not\rightarrow f(x_0) \dashv$$

18 Lecture 18 Oct 22nd

18.1 Continuity on Metric Spaces (Continued)

Definition 46 (Continuity on a Space)

We say that

$$f : (X, d_X) \rightarrow (Y, d_Y)$$

is *continuous* on X if f is continuous at each $x_0 \in X$.

We let

$$C(X, Y) := \{f : X \rightarrow Y \mid f \text{ is continuous on } X\},$$

be the set of all continuous functions on X .

Note 18.1.1

In the case where $Y = \mathbb{R}$, we will simply write $C(X, X)$ as $C(X)$.

Remark 18.1.1

We can also define the following set

$$C_b(X) = \{f \in C(X) \mid f \text{ is bounded}\}.$$

We can define $\|\cdot\|_\infty$ on $C_b(X)$ by

$$\|f\|_\infty = \sup\{|f(x)| \mid x \in X\}.$$

Then we have that $C_b(X) \subseteq \ell_\infty(X)$.

Theorem 45 (Analogue of Sequential Characterization of Continuity on a Space, and Continuity and Neighbourhoods)

Let $f : (X, d_X) \rightarrow (Y, d_Y)$. TFAE

1. f is continuous;
 2. $f^{-1}(W)$ is open for every open set $W \subset Y$;
 3. $x_n \rightarrow x_0 \in X \implies f(x_n) \rightarrow f(x_0) \in Y$.
-

Proof

(1) \implies (2) : Let $W \subset Y$ be open, and $V = f^{-1}(W)$.

$x_0 \in V \implies f(x_0) = y_0 \in W \implies W$ is a neighbourhood of y_0
 $\implies (\because \text{Theorem 43}) V$ is a neighbourhood of x_0
 $\implies x_0 \in V^\circ \implies V$ is open \dashv

(2) \implies (3) : $x_n \rightarrow x_0 \in X$

$\implies \forall \varepsilon > 0 (\because B_Y(f(x_0), \varepsilon)$ open)
 $\implies x_0 \in V = f^{-1}(B_Y(f(x_0), \varepsilon))$, which is open
 $\implies \exists \delta > 0 B_X(x_0, \delta) \subset V$
 $x_n \rightarrow x_0 \implies \exists N_0 \in \mathbb{N} \forall n \geq N_0 x_n \in B_X(x_0, \delta)$
 $\implies f(x_n) \in B_Y(f(x_0), \varepsilon) \implies f(x_n) \rightarrow f(x_0) \dashv$

(3) \implies (1) : Sps $f \dashv$ continuous, i.e.

$\exists \varepsilon_0 > 0 \forall \delta \geq 0 \exists x_\delta \in X \quad d_X(x_\delta, x_0) < \delta \wedge d_Y(f(x_\delta), f(x_0)) > \varepsilon_0$
 $\implies \forall n \in \mathbb{N} \exists x_n \in d_X(x_0, x_n) < \frac{1}{n} \wedge d_Y(f(x_0), f(x_n)) > \varepsilon_0 \dashv$

Remark 18.1.2

Note that if $f : X \rightarrow Y$ and $B \subset Y$, then


$$(f^{-1}(B))^C = f^{-1}(B^C).$$

Thus we have that $f : (X, d_X) \rightarrow (Y, d_Y)$ is continuous iff $f^{-1}(F)$ is closed for each closed $F \subset Y$.


QUESTION: For the forward direction¹, if $f : (X, d_X) \rightarrow (Y, d_Y)$ is continuous, and if $U \subset X$ is open, is $f(U)$ open? **No**.

¹ instead of talking about the pullback


Example 18.1.1

Consider $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $\forall x \in X, f(x) = 1$. Then $f(\mathbb{R})$ is not open. 

This motivates us to consider such “nice” functions that allow us to bring open sets to open sets, and closed to their closed counterpart.


 **Definition 47 (Homeomorphism)**

A function $\varphi : (X, d_X) \rightarrow (Y, d_Y)$ is a *homeomorphism* if φ is bijective and if both φ and φ^{-1} are continuous.

 **Note 18.1.2**

If φ is a homeomorphism, then we have

- $\varphi(W) \subset Y$ is open $\iff W \subset X$ is open;
- $\varphi(F) \subset Y$ is closed $\iff F \subset X$ is closed.

 **Definition 48 (Equivalent Metric Spaces)**


We say that (X, d_X) and (Y, d_Y) are *equivalent metric spaces* if there exists a bijective $\varphi : X \rightarrow Y$, and $c_1, c_2 \geq 0$ such that

$$c + 1d_X(x_1, x_2) \leq d_Y(\varphi(x_1), \varphi(x_2)) \leq c_2d_X(x_1, x_2).$$

Exercise 18.1.1

Show that the φ in  Definition 48 is a *homeomorphism*.

Example 18.1.2

Let (X, d) be a metric space, where X is any set and d is the discrete metric. Let $f : (X, d) \rightarrow (Y, d_Y)$, where (Y, d_Y) is another metric space that is arbitrary. Since (X, d) is discrete, it is clear that if $W \subset Y$ is open, then $f^{-1}(W)$ is open. 

QUESTION: Suppose that $f : (\mathbb{R}, |\cdot|) \rightarrow (Y, d)$. When is f continuous?

Let $y_0 \in Y$. We know that $\{y_0\}$ is both open and closed. Then if f is continuous, we must have that $f^{-1}(\{y_0\})$ is both open and closed. Therefore, f must be a constant function.


Exercise 18.1.2

Use the *Intermediate Value Theorem* to prove that the only open and closed sets in \mathbb{R} are \emptyset and \mathbb{R} .

 **Definition 49 (Continuity on a set)**

Let $A \subset (X, d)$ and $f : X \rightarrow (Y, d_Y)$. We say that f is **continuous** on A iff $f \upharpoonright_A$ is continuous on (A, d_A) , where d_A is the induced metric, and $f \upharpoonright_A$ is the restriction of f on A .

Remark 18.1.3

From the sequential characterization of continuity, we have that (A, d_A) is the induced metric iff whenever $\{x_n\} \subset A$ is a sequence with $x_n \rightarrow x_0$, then $f(x_n) \rightarrow f(x_0)$. 

19 Lecture 19 Oct 24th

19.1 Completeness of Metric Spaces

QUESTION: Is there an intrinsic way for us to tell if a sequence $\{x_n\} \subset (X, d)$ converges?

OBSERVATION Assume that $x_n \rightarrow x_0$. Then

$$\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall n \geq N_0 \ d(x_0, x_n) < \frac{\varepsilon}{2}.$$

Thus if $m, n \geq N_0$, we have

$$d(x_m, x_n) < d(x_m, x_0) + d(x_0, x_n) < \varepsilon.$$

Definition 50 (Cauchy)

We say that a sequence $\{x_n\} \subset (X, d)$ is *Cauchy* if

$$\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall m, n \geq N_0 \ d(x_m, x_n) < \varepsilon.$$

Theorem 46 (Convergent Sequences are Cauchy)

Every convergent sequence is Cauchy.

We proved this in our observation.

QUESTION: Is the converse true? **No.**

Example 19.1.1

Let $X = (0, 1)$ with the usual metric. Let $x_n = \frac{1}{n}$. It is clear that $\{x_n\}$ is Cauchy in (X, d) , but the sequence does not converge.¹

¹ The flaw here lies in the fact that X is open. Should we have chosen $X = [0, 1]$, then the limit point 0 would have been included, allowing the sequence to actually converge.

Definition 51 (Complete Metric Spaces)

A metric space (X, d) is **complete** if each Cauchy sequence $\{x_n\} \subset X$ converges in (X, d) .

19.1.1 Basic Properties of Cauchy Sequences

OBSERVATION Given a sequence $\{x_n\} \subset (X, d)$, it is possible that $\{x_n\}$ diverges but $\{x_n\}$ has a subsequence $\{x_{n_k}\}$ that converges.

Example 19.1.2

The sequence $\{x_n\}$ defined by $x_n = (-1)^{n-1}$, i.e.

$$\{x_n\} = \{1, -1, 1, -1, \dots\},$$

is divergent. However, $x_{2k} \rightarrow -1$ and $x_{2k+1} \rightarrow 1$.

Theorem 47 (★ ★ ★ Convergent Cauchy Subsequences)

Let $\{x_n\} \subset (X, d)$ be Cauchy and assume $x_{n_k} \rightarrow x_0$ for some subsequence $\{x_{n_k}\}_{k=1}^{\infty}$. Then $x_n \rightarrow x_0$.

Proof (★ ★ ★)

$$\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall m, n \in N_0 \quad d(x_n, x_m) < \frac{\varepsilon}{2}$$

$$x_n \rightarrow x_0 \implies \exists k_0 \in \mathbb{N} \quad n_{k_0} \geq N_0 \quad d(x_0, x_{k_0}) < \frac{\varepsilon}{2}$$

$\therefore n \geq N_0 \implies$


$$d(x_n, x_0) \leq d(x_n, x_{k_0}) + d(x_{k_0}, x_0) < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

$\therefore x_n \rightarrow x_0$

 **Definition 52 (Boundedness)**

Let $A \subset (X, d)$. A is *bounded* if

$$\exists M > 0 \exists x_0 \in X \ A \subset B[x_0, M].$$

 **Proposition 48 (Cauchy Sequences are Bounded)**

If $\{x_n\} \subset (X, d)$ is Cauchy, then $\{x_n\}$ is bounded.

 **Proof**


Let $\varepsilon = 1$. $\exists N_0 \in \mathbb{N} \forall m, n \geq N_0 \ d(x_n, x_m) < \varepsilon$. In particular, if $n \geq N_0$, then $d(x_n, x_{N_0}) < 1$. Then, let

$$M = \max\{d(x_1, x_{N_0}), d(x_2, x_{N_0}), \dots, d(x_{N_0-1}, x_{N_0}), 1\}$$

Then it is clear that $\{x_n\} \subset B[x_{N_0}, M]$.

19.1.2 Examples of Complete Spaces

19.1.2.1 Completeness of \mathbb{R}

 **Theorem 49 (Bolzano-Weierstrass)**

Every bounded sequence $\{x_n\} \subset \mathbb{R}$ has a convergent subsequence.

Be sure to review a proof of this and add it here.

Theorem 50 (\mathbb{R} is complete) \mathbb{R} is complete.**Proof**

If $\{x_n\} \subset \mathbb{R}$ is Cauchy, then it is bounded by Proposition 48, and so by Bolzano-Weierstrass, $\{x_n\}$ has a convergent subsequence $\{x_{n_k}\}$ such that $x_{n_k} \rightarrow x_0$. Since $\{x_n\}$ is Cauchy, by Theorem 47, $x_n \rightarrow x_0$.

Example 19.1.3


Consider $(\mathbb{R}^n, \|\cdot\|_p)$, with $1 \leq p \leq \infty$. Let $\{\vec{x}_k\} = \{(x_{k,1}, x_{k,2}, \dots, x_{k,n})\}$ be Cauchy in $(\mathbb{R}^n, \|\cdot\|_p)$.

$$\because |x_{k,j} - x_{m,j}| \leq \|\vec{x}_k - \vec{x}_m\|_p$$


$\implies \{x_{k,j}\}$ is Cauchy for each $j = 1, \dots, n$

$\implies x_{k,j} \rightarrow x_{0,j}$ for each $j = 1, \dots, n$ \because Theorem 47

$\implies \vec{x}_k \rightarrow \vec{x}_0 = (x_{0,1}, x_{0,2}, \dots, x_{0,n})$

$\implies (\mathbb{R}^n, \|\cdot\|_p)$ is complete. 

Example 19.1.4

Let (X, d) be discrete². If $\{x_n\}$ is Cauchy, then $\exists N_0 \in \mathbb{N}$ such that $\forall m, n \geq N_0$, we have $x_n = x_m$, i.e. $\{x_n\}$ converges. Therefore, (X, d) is complete. 


² By discrete, we mean a discrete metric space, i.e. d is a discrete metric.

Example 19.1.5 (★)

Let $X = \left\{1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}, \dots\right\} \subset \mathbb{R}$ with the induced standard metric.

Recall that each of the singleton $\left\{\frac{1}{n}\right\}$ is open.

Note that given $Y = \{1, 2, \dots, n, \dots\} = \mathbb{N}$ with the discrete metric, if we define $\varphi : \mathbb{N} \rightarrow \left\{1, \frac{1}{2}, \dots, \frac{1}{n}, \dots\right\}$ by $\varphi(n) = \frac{1}{n}$, then φ is a homeomorphism, and so (Y, d) , where d is the discrete metric, is complete.

However, as shown before, since $\left\{\frac{1}{n}\right\}$ is Cauchy but not convergent, $X = \left\{1, \frac{1}{2}, \dots, \frac{1}{n}, \dots\right\}$ is not complete. 

20 Lecture 20 Oct 26th

20.1 Completeness of Metric Spaces (Continued)

20.1.1 Examples of Complete Spaces (Continued)

20.1.1.1 Completeness of ℓ_p

Theorem 51 (★ Completeness of ℓ_p)

ℓ_p is complete for every $1 \leq p \leq \infty$.

Proof

$p = \infty$: Let $\{\vec{x}_k\} \subset \ell_\infty$ be Cauchy in $\|\cdot\|_\infty$. We have

$$\vec{x}_k = \{x_{k,1}, x_{k,2}, \dots, x_{k,j}, \dots\}$$

$$\implies \forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall m, n \geq N_0 \|\vec{x}_n - \vec{x}_m\|_\infty < \frac{\varepsilon}{2}$$

$$\because |x_{n,j} - x_{m,j}| \leq \|\vec{x}_n - \vec{x}_m\|_\infty < \frac{\varepsilon}{2},$$

each of the \vec{x}_k , for $k \geq N_0$, is Cauchy in \mathbb{R} .

$$\implies \exists x_{0,j} \in \mathbb{R} \quad x_{k,j} \rightarrow x_{0,j} \quad \because \mathbb{R} \text{ is complete}$$

Let $\vec{x}_0 = \{x_{0,1}, x_{0,2}, \dots, x_{0,j}, \dots\}$ and $x_{0,j} = \lim_{k \rightarrow \infty} x_{k,j}$.

By our argument on Line 4, we have that

$$|x_{n,j} - x_{0,j}| = \lim_{m \rightarrow \infty} |x_{n,j} - x_{m,j}| \leq \frac{\varepsilon}{2} < \varepsilon \quad (20.1)$$

$$\implies \{x_{n,j} - x_{0,j}\}_{j=1}^\infty \in \ell_\infty$$

$$\implies \{x_{0,j}\}_{j=1}^\infty \in \ell_\infty$$

Also, by Equation (20.1), we have

$$\|\vec{x}_n - \vec{x}_0\|_\infty \leq \frac{\varepsilon}{2} < \varepsilon,$$

so $\vec{x}_k \rightarrow \vec{x}_0$. \dashv .

$1 \leq p < \infty$: Let $\{\vec{x}_k\} \subset \ell_p$ be Cauchy. By the same argument as above, $|x_{n,j} - x_{m,j}| \leq \|\vec{x}_n - \vec{x}_m\|_p \implies \{x_{k,j}\}_{j=1}^\infty$ is Cauchy for each j . Since \mathbb{R} is complete, let $x_{0,j} = \lim_{k \rightarrow \infty} x_{k,j}$, and

$$\vec{x}_0 = \{x_{0,1}, x_{0,2}, \dots, x_{0,j}, \dots\}.$$

Now $\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall n, m \geq N_0 \|\vec{x}_n - \vec{x}_m\| < \frac{\varepsilon}{2}$. Thus for $j \in \mathbb{N}$,

$$\left(\sum_{i=1}^j |x_{n,i} - x_{m,i}|^p \right)^{\frac{1}{p}} \leq \|\vec{x}_n - \vec{x}_m\|_p < \frac{\varepsilon}{2}.$$

Then for $n \geq N_0$,

$$\left(\sum_{i=1}^j |x_{k,i} - x_{0,i}|^p \right)^{\frac{1}{p}} = \lim_{m \rightarrow \infty} \left(\sum_{i=1}^j |x_{n,i} - x_{m,i}|^p \right)^{\frac{1}{p}} \leq \frac{\varepsilon}{2}$$

for each j , and so

$$\lim_{j \rightarrow \infty} \left(\sum_{i=1}^j |x_{n,i} - x_{0,i}|^p \right)^{\frac{1}{p}} \leq \frac{\varepsilon}{2}$$

$$\implies \vec{x}_0 \in \ell_p \text{ and } \|\vec{x}_n - \vec{x}_0\|_p \leq \frac{\varepsilon}{2} < \varepsilon.$$

20.1.1.2 Completeness of $(C_b(X), \|\cdot\|_\infty)$

Definition 53 (Convergence of Functions)


A sequence of functions $f_n : (X, d_X) \rightarrow (Y, d_Y)$ is said to **converge pointwise** to some function $f_0 : (X, d_X) \rightarrow (Y, d_Y)$ if for each $x_0 \in X$,

$$\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall n \geq N_0 \ d_Y(f_n(x_0) - f_0(x_0)) < \varepsilon.$$

The sequence f_n is said to **converge uniformly** if

$$\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall n \geq N_0 \forall x \in X \ d_Y(f_n(x) - f_0(x)) < \varepsilon.$$


Remark 20.1.1


It is clear that uniform convergence implies pointwise convergence. 

Example 20.1.1 (Pointwise Convergent but not Uniformly Convergent)

Let $X = [0, 1]$, $Y = \mathbb{R}$, $f_n(x) = x^n$ for each $n \in \mathbb{N}$. It is quite clear that

$$f_n(x) \rightarrow f_0(x) = \begin{cases} 0 & x \in [0, 1) \\ 1 & x = 1 \end{cases}.$$

f_n is pointwise convergent but not uniformly convergent; just take $\varepsilon = \frac{1}{2}$. 

 **Theorem 52 (★ ★ ★ Uniformly Convergent Pointwise Continuous Functions have a Pointwise Continuous Limit)**

Assume that $f_n : (X, d_X) \rightarrow (Y, d_Y)$ converges uniformly to $f_0 : (X, d_X) \rightarrow (Y, d_Y)$. If each f_n is continuous at $x_0 \in X$, then f_0 is continuous at x_0 .

This is a classic $\frac{\varepsilon}{3}$ argument.

 **Proof**

$$\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall n \geq N_0 \forall x \in X \ d_Y(f_n(x) - f_0(x)) < \frac{\varepsilon}{3}$$

$$f_n \text{ is continuous at } x_0 \implies \exists \delta > 0 \forall x \in X \ x \in B(x_0, \delta)$$

$$\implies \forall n_0 \geq N_0 \ d_Y(f_{n_0}(x) - f_{n_0}(x_0)) < \frac{\varepsilon}{3}$$

$$\implies d_Y(f_0(x), f_0(x_0))$$

$$\leq d_Y(f_0(x), f_{n_0}(x)) + d_Y(f_{n_0}(x), f_{n_0}(x_0)) + d_Y(f_{n_0}(x_0), f_0(x_0))$$

$$< \varepsilon$$

21 Lecture 21 Oct 31st

21.1 Completeness of Metric Spaces (Continued 2)

21.1.1 Examples of Complete Spaces (Continued 2)

21.1.1.1 Completeness of $(C_b(X), \|\cdot\|_\infty)$ (Continued)

Note 21.1.1

A normed linear space V is called a **Banach space** if $(V, \|\cdot\|)$ is complete with respect to d_V .

Theorem 53 (★ ★ ★ Completeness for $C_b(X)$)

The space $(C_b(X), \|\cdot\|_\infty)$ is Banach (i.e. complete).

This will come out in the final.

Proof

Let $\{f_n\} \subset C_b(X)$ be Cauchy.

$$\implies \forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall n, m \geq N_0 \quad \|f_n - f_m\|_\infty < \frac{\varepsilon}{2}, \quad (*)$$

and

$$\forall x \in X \quad |f_n(x) - f_m(x)| \leq \|f_n - f_m\|_\infty < \frac{\varepsilon}{2}.$$

$\therefore \{f_n(x)\}$, for every $x \in X$ is Cauchy, and so $\{f_n(x)\}$ is complete.

Let $f_0(x) = \lim_{n \rightarrow \infty} f_n(x)$, and in particular, $\forall n \geq N_0, \forall x \in X$, we

have

$$|f_n(x) - f_0(x)| = \lim_{m \rightarrow \infty} |f_n(x) - f_m(x)| \leq \frac{\varepsilon}{2} < \varepsilon.$$

So $f_n \rightarrow f_0$ uniformly. By  Theorem 52, f_0 is continuous.

It remains to show that f_0 is bounded: we have that $\{f_n\}$ is bounded.

Let $M > 0$ such that $\|f_n\|_\infty \leq M$ for all $n \in \mathbb{N}$. Let $x \in X$.

From (*), we can find $n_0 \in \mathbb{N}$ such that $|f_{n_0}(x) - f_0(x)| \leq 1$.

$$\implies |f_0(x)| \leq |f_0(x) - f_{n_0}(x)| + |f_{n_0}(x)| \leq 1 + M$$

$\therefore f_0(x) \in C_b(X)$.

Note 21.1.2

Given any set X , if (X, d) is a metric space with the discrete metric, then

$$(C_b(X), \|\cdot\|_\infty) = (\ell_\infty, \|\cdot\|_\infty).$$

21.1.2 Characterizations of Completeness

We shall state the following without proving it, although the proof is straightforward: view $\{a_n\}$ and $\{b_n\}$ as increasing and decreasing sequences respectively and use the monotone convergence theorem.

Theorem 54 (Nested Interval Theorem)

If $\{[a_n, b_n]\}$ with $[a_{n+1}, b_{n+1}] \subset [a_n, b_n]$, then

$$\bigcap_{n=1}^{\infty} [a_n, b_n] \neq \emptyset.$$

We know that this works for \mathbb{R} , but does this work for (X, d) ? In particular, we conjecture that:

If $\{F_n\}$ is a sequence of non-empty closed sets in (X, d) , with

$F_{n+1} \subseteq F_n$, then

$$\bigcap_{n=1}^{\infty} F_n \neq \emptyset.$$

However, this is not true, as shown in the following example.

Example 21.1.1

Let $X = \mathbb{R}$, and $F_n = [n, \infty)$, and $F_{n+1} \subsetneq F_n$. Note that F_n is indeed closed since its complement, $(-\infty, n)$, is open. We notice that

$$\bigcap_{n=1}^{\infty} F_n = \emptyset.$$

Example 21.1.2

Let $X = (0, 1)$, and $F_n = (0, \frac{1}{n}]$, which is closed in X , and that $F_{n+1} \subsetneq F_n$. However, once again, we notice that

$$\bigcap_{n=1}^{\infty} F_n = \emptyset.$$

Of course, one would ask the question as to why does such a property not hold. The following notion will explain why.

Definition 54 (Diameter of a Set)

Given a subset $A \subset (X, d)$, we define the *diameter* of A as

$$\text{diam}(A) = \sup\{d(x, y) \mid x, y \in A\}.$$

Proposition 55 (Diameters of Subsets)

Let $A \subseteq B \subset (X, d)$. Then

1. $\text{diam}(A) \leq \text{diam}(B)$;
2. $\text{diam}(A) = \text{diam}(\overline{A})$.




Figure 21.1: Intuitive illustration of Definition 54. Red lines are the diameters, as captured by the sup function. Blue lines are other possible candidates, but none of them can be a supremum.

 **Proof**

1. If $A = B$, then there is nothing to prove. Suppose $A \subsetneq B$. Suppose to the contrary that $\text{diam}(A) > \text{diam}(B)$. Let $x_A, y_A \in A$ such that $d(x_A, y_A) = \text{diam}(A)$ and $x_B, y_B \in B$ such that $d(x_B, y_B) = \text{diam}(B)$. By our assumption, we have

$$d(x_A, y_A) > d(x_B, y_B).$$

However, $x_A, y_A \in A \subseteq B$, and by definition of a diameter, we have

$$d(x_B, y_B) \geq d(x_A, y_A),$$

which is a contradiction. This proves the statement.

2. If $\text{diam}(A) = \infty$, then we must have $\text{diam}(\bar{A}) = \infty$ since $A \subseteq \bar{A}$. Thus WMA $\text{diam}(A) = d < \infty$. Let $x_0, y_0 \in \bar{A}$. Then given any $\varepsilon > 0$, by definition of limits, we can find $x_1, y_1 \in A$ such that

$$d(x_0, x_1) < \frac{\varepsilon}{2} \text{ and } d(y_0, y_1) < \frac{\varepsilon}{2}.$$

Hence

$$\begin{aligned} d(x_0, y_0) &\leq d(x_0, x_1) + d(x_1, y_1) + d(y_1, y_0) \\ &< \frac{\varepsilon}{2} + d + \frac{\varepsilon}{2} = d + \varepsilon. \end{aligned}$$

Thus $\text{diam}(\bar{A}) \leq d + \varepsilon$, for any $\varepsilon > 0$. Therefore by the earlier part,

$$\text{diam}(\bar{A}) \leq d = \text{diam}(A) \leq \text{diam}(\bar{A}).$$

With this notion, we have a partial equivalence to the nested interval theorem, of which we shall prove in the next lecture.

22 Lecture 22 Nov 02nd

22.1 Completeness of Metric Spaces (Continued 3)

22.1.1 Characterizations of Completeness (Continued)

We are now ready to prove the following statement.

Theorem 56 (Cantor's Intersection Principle)

Let (X, d) be a metric space. TFAE:

1. (X, d) is complete.
2. If $\{F_n\}$ is a sequence of non-empty closed subsets such that $F_{n+1} \subset F_n$ for all $n \in \mathbb{N}$, and if $\lim_{n \rightarrow \infty} \text{diam}(F_n) = 0$, then

$$\bigcap_{n=1}^{\infty} F_n \neq \emptyset.$$

Proof

(1) \implies (2): ¹For each $n \in \mathbb{N}$, pick $x_n \in F_n$. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence formed from these x_n 's.

By the assumption that $\lim_{n \rightarrow \infty} \text{diam}(F_n) = 0$, we have that

$$\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \text{ diam}(F_{N_0}) < \varepsilon.$$

In particular, for $n, m \geq N_0$, we have that $x_n, x_m \in F_{N_0}$, as $F_n, F_m \subset F_{N_0}$, and so

$$d(x_n, x_m) \leq \text{diam}(F_{N_0}) < \varepsilon.$$

¹ Since we have a sequence of non-empty closed subsets, we can, by using * Axiom 2, form a sequence of elements in X from each of the F_n 's. By proving that this sequence of elements is Cauchy, we obtain a limit point from the assumption that X is complete. From there, it remains to show that the limit point lives in all of the F_n 's.

Thus $\{x_n\}$ is Cauchy. By assumption that (X, d) is complete, $x_n \rightarrow x_0 \in X$. Thus $\exists N_1 \in \mathbb{N}$ such that $\forall n \geq N_1, d(x_n, x_0) < \varepsilon$. Thus, for any such n , since $F_{n+1} \subset F_n$, $\{x_n, x_{n+1}, x_{n+2}, \dots\} \subset F_n$, and the sequence converges to x_0 . Since F_n is closed, we must have $x_0 \in F_n$. This forces $x_0 \in F_n$ for every $n \in \mathbb{N}$. This completes (\implies).

(2) \implies (1): Let $\{x_n\} \subset X$ be Cauchy. Let $F_n = \{x_n, x_{n+1}, x_{n+2}, \dots\}$. We have that F_n is closed: given any $y \notin F_n$, we can pick $\delta = \frac{1}{2} \min\{d(x_i, x_j) : n \leq i < j\}$ and we would have that $B(y, \delta) \cap F_n = \emptyset$.

Note that $F_{n+1} \subset F_n$.

$\therefore \{x_n\}$ is Cauchy, $\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall n, m \geq N_0 d(x_n, x_m) < \frac{\varepsilon}{2}$.

Consequently,

$$\text{diam}(\{x_{N_0}, x_{N_0+1}, \dots\}) = \text{diam}(F_{N_0}) \leq \frac{\varepsilon}{2} < \varepsilon.$$

$\therefore \text{diam}(F_n) \rightarrow 0$, which, along with assumption, implies that²

$$\bigcap_{n=1}^{\infty} F_n = \{x_0\}.$$

² Note that the intersection can only contain one element, since $\text{diam}(F_n) \rightarrow 0$.

Also, since $\text{diam}(F_n) \rightarrow 0$, we have that for any $k > 0$, $F_{i_k} \subseteq B(x_0, \frac{1}{k})$ ³. This implies that for each k , $B(x_0, \frac{1}{k})$ contains the tail of the sequence $\{x_n\}$. Then, inductively so, we have

³ Otherwise, x_0 cannot be a limit point.

$$\begin{aligned} k = 1 &\implies \exists n_1 > 0 \quad x_{n_1} \in B(x_0, 1) \\ k = 2 &\implies \exists n_2 > 0 \quad x_{n_2} \in B\left(x_0, \frac{1}{2}\right) \\ &\vdots \\ k = m &\implies \exists n_m > 0 \quad x_{n_m} \in B\left(x_0, \frac{1}{m}\right) \\ &\vdots \end{aligned}$$

$\therefore x_{n_m} \rightarrow x_0$.

Then since $\{x_n\}$ is Cauchy, and $\{x_{n_m}\}$ is a subsequence of $\{x_n\}$, we have $x_n \rightarrow x_0$.

Let $(X, \|\cdot\|)$ be a normed linear space. A series in X is called a **formal sum**, expressed as

$$\sum_{n=1}^{\infty} x_n = x_1 + x_2 + \dots + x_n + \dots, \quad (22.1)$$

where $\{x_n\} \subseteq X$. For each $k \in \mathbb{N}$, the k^{th} **partial sum** of Equation (22.1) is

$$S_k = \sum_{n=1}^k x_n = x_1 + x_2 + \dots + x_k.$$

We say that $\sum_{n=1}^{\infty} x_n$ converges in $(X, \|\cdot\|)$ if $\{S_k\}_{k=1}^{\infty}$ converges. In this case, we write

$$\sum_{n=1}^{\infty} x_n = \lim_{k \rightarrow \infty} S_k.$$

Otherwise, $\sum_{n=1}^{\infty} x_n$ is said to diverge.

Theorem 57 (★ ★ Weierstrass M-test)

Let $(X, \|\cdot\|)$ be a normed linear space. TFAE:

1. $(X, \|\cdot\|)$ is complete, i.e. $(X, \|\cdot\|)$ is a Banach space.
2. If $\sum_{n=1}^{\infty} x_n$ is such that $\sum_{n=1}^{\infty} \|x_n\|$ converges, then $\sum_{n=1}^{\infty} x_n$ converges.

Proof

(1) \implies (2): Given $\sum_{n=1}^{\infty} x_n$, let

$$S_k = \sum_{n=1}^k x_n \text{ and } T_k = \sum_{n=1}^k \|x_n\|.$$

Suppose T_k converges. Then in particular, $\{T_k\}$ is Cauchy. Thus

$$\forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall n > m \geq N_0$$

$$T_n - T_m = \sum_{k=1}^n \|x_k\| - \sum_{k=1}^m \|x_k\| = \sum_{k=m+1}^n \|x_k\| < \varepsilon.$$

$\therefore N_0 \leq m < n \implies$

$$\begin{aligned} \|S_n - S_m\| &= \left\| \sum_{k=1}^n x_k - \sum_{k=1}^m x_k \right\| = \left\| \sum_{k=m+1}^n x_k \right\| \\ &\leq \sum_{k=m+1}^n \|x_k\| \quad \because \text{Triangle Ineq.} \\ &< \varepsilon. \end{aligned}$$

$\therefore \{S_k\}$ is Cauchy, and since $(X, \|\cdot\|)$ is complete, $\{S_k\}$ is convergent.

(2) \implies (1): Suppose $\{x_n\}$ is Cauchy in $(X, \|\cdot\|)$. We can find an increasing sequence

$$N_0 < n_1 < n_2 < \dots < n_j < \dots \in \mathbb{N},$$

for some $N_0 \in \mathbb{N}$ such that

$$\|x_{n_j} - x_{n_{j+1}}\| < \frac{1}{2^j}.$$

Then by the **infinite geometric series**,

$$\sum_{j=1}^{\infty} \|x_{n_j} - x_{n_{j+1}}\| \leq \sum_{j=1}^{\infty} \frac{1}{2^j} < \infty.$$

$\therefore \sum_{j=1}^{\infty} (x_{n_j} - x_{n_{j+1}})$ converges to some $x_0 \in X$. In particular, notice that the partial sums are **telescoping series**:

$$S_k = \sum_{j=1}^k (x_{n_j} - x_{n_{j+1}}) = x_{n_1} - x_{n_{k+1}} \rightarrow x_0.$$

It follows that as $k \rightarrow \infty$,

$$x_{n_{k+1}} \rightarrow x_{n_1} - x_0.$$

We have that the subsequence $\{x_{n_k}\}$ of our Cauchy sequence $\{x_n\}$ has a limit point.

23 Lecture 23 Nov 05th

23.1 Completeness of Metric Spaces (Continued 4)

23.1.1 Characterizations of Completeness (Continued 2)

Example 23.1.1

Let

$$\varphi(x) = \begin{cases} x & x \in [0, 1] \\ 2 - x & x \in [1, 2] \end{cases}.$$

Extend φ to \mathbb{R} by

$$\varphi(x + 2) = \varphi(x) \quad \text{for all } x \in \mathbb{R}.$$

Define

$$f(x) = \sum_{n=1}^{\infty} \left(\frac{3}{4}\right)^n \varphi(4^n x).$$




Figure 23.1: Sawtooth-like graph from φ

Figure 23.2 is a simplified graph of f , drawn using the online tool Desmos.


It is clear that $\varphi \in C_b(\mathbb{R})$, and $\|\varphi\|_{\infty} = 1$. Thus

$$\sum_{n=1}^{\infty} \left\| \left(\frac{3}{4}\right)^n \varphi(4^n x) \right\|_{\infty} = \sum_{n=1}^{\infty} \left(\frac{3}{4}\right)^n < \infty,$$

and so

$$f(x) = \lim_{L \rightarrow \infty} \sum_{n=1}^L \left(\frac{3}{4}\right)^n \varphi(4^n x) = \lim_{L \rightarrow \infty} S_L(x),$$

uniformly so. Since the partial sums are continuous, $f \in C_b(\mathbb{R})$.

Figure 23.2: Function of f as generated on Desmos. See it live.

However, f is not **differentiable**. Let $x \in \mathbb{R}$. For each $m \in \mathbb{N}$, we can find $k \in \mathbb{Z}$ such that

$$k \leq 4^m x \leq k + 1.$$

Let

$$p_m = \frac{k}{4^m} \text{ and } q_m = \frac{k+1}{4^m},$$

and for any $n \in \mathbb{N}$,

$$\alpha = 4^n p_m = 4^{n-m} k \text{ and } \beta = 4^n q_m = 4^{n-m} (k+1).$$

Now

- if $n > m$, then since α and β differ by an even integer, $|\varphi(\alpha) - \varphi(\beta)| = 0$;
- if $n = m$, then α and β differs by 1, and so $|\varphi(\alpha) - \varphi(\beta)| = 1$;
- if $n < m$, then there are no integers between α and β , and so

$$|\varphi(\alpha) - \varphi(\beta)| = |4^n p_m - 4^n q_m|^1 = |4^{n-m} k - 4^{n-m} (k+1)| = 4^{n-m}.$$

¹ Note that if we have $1 \leq \alpha, \beta \leq 2$, we still get the same formula.

For large enough m , consider

$$|f(p_m) - f(q_m)| = \left| \sum_{n=1}^{\infty} \left(\frac{3}{4}\right)^n (\varphi(4^n p_m) - \varphi(4^n q_m)) \right|$$

$$= \left| \sum_{n=1}^m \left(\frac{3}{4}\right)^n (\varphi(4^n p_m) - \varphi(4^n q_m)) \right| \quad (23.1)$$

$$\geq \left| \left(\frac{3}{4}\right)^m - \sum_{n=1}^{m-1} \left(\frac{3}{4}\right)^n |\varphi(4^n p_m) - \varphi(4^n q_m)| \right| \quad (23.2)$$

$$= \left| \left(\frac{3}{4}\right)^m - \sum_{n=1}^{m-1} \left(\frac{3}{4}\right)^n 4^{n-m} \right| \quad (23.3)$$

$$= \left| \left(\frac{3}{4}\right)^m - \frac{1}{4^m} \sum_{n=1}^{m-1} 3^n \right|$$

$$= \left| \left(\frac{3}{4}\right)^m - \frac{1}{4^m} \left[\frac{3^m - 1}{2} \right] \right| \quad (23.4)$$

$$= \frac{1}{4^m} \left[\frac{3^m + 1}{2} \right] > \frac{1}{2} \cdot \left(\frac{3}{4}\right)^m$$

where we note that

(23.1) terms after m are eliminated as they are 0 as argued previously;

(23.2) by the reverse Triangle ineq. and the case where $n = m$;

(23.3) using the argument for when $n < m$;

(23.4) using the formula for a finite geometric sum.

Hence we observe that

$$\frac{|f(p_m) - f(q_m)|}{|p_m - q_m|} > 4^m \cdot \frac{3^m}{2 \cdot 4^m} = \frac{3^m}{2}.$$

Now if $p_m = x$, then

$$\frac{|f(x) - f(q_m)|}{|x - q_m|} > \frac{3^m}{2}.$$

If $p_m \neq x$, then

$$\begin{aligned} \frac{3^m}{2} &< \frac{|f(p_m) - f(q_m)|}{|p_m - q_m|} \leq \frac{|f(p_m) - f(x)| + |f(x) - f(q_m)|}{|p_m - q_m|} \\ &\leq \frac{|f(p_m) - f(x)|}{|p_m - x|} + \frac{|f(x) - f(q_m)|}{|x - q_m|}, \end{aligned}$$

which implies that either


$$\frac{|f(x) - f(q_m)|}{|x - q_m|} > \frac{3^m}{2},$$

or

$$\frac{|f(p_m) - f(x)|}{|p_m - x|} > \frac{3^m}{2}.$$

Then for any sequence $\{t_m\}$ such that $t_m \rightarrow x$, and $t_m \neq x$, we have that

$$\frac{|f(x) - f(t_m)|}{|x - t_m|} \geq \frac{3^m}{4} \rightarrow \infty$$

as $m \rightarrow \infty$. Thus the function f is not differentiable at any x . 

24 Lecture 24 Nov 07th

24.1 Completions of Metric Spaces

Definition 56 (Isometry)

A map $\varphi : (X, d_X) \rightarrow (Y, d_Y)$ is called an *isometry* if

$$d_Y(\varphi(x_1), \varphi(x_2)) = d_X(x_1, x_2).$$

Definition 57 (Completion)

A *completion* of a metric space (X, d) is a pair $((Y, d_Y), \varphi)$ where (Y, d_Y) is a complete metric space, $\varphi : X \rightarrow Y$ is an isometry, and $\overline{\varphi(X)} = Y$.

Proposition 58 (Subsets of Complete Spaces are Complete if they are Closed)

Let (X, d) be a complete metric space. Let $A \subset X$. Then (A, d_A) is complete iff A is closed.

Proof

(\implies) : (A, d_A) is complete

$\implies \{x_n\} \subset A \text{ Cauchy} \implies x_n \rightarrow x_0 \implies x_0 \in A \implies$
 $\text{Lim}(A) \subseteq A$
 $\implies A \text{ is closed.}$

(\longleftarrow) Let $\{x_n\} \subset A$ be Cauchy in (A, d_A)
 $\implies \{x_n\}$ is Cauchy in (X, d)
 $\implies x_n \rightarrow x_0 \in X$
 $\implies (\because A \text{ is closed}) x_0 \in A$
 $\implies (A, d_A)$ is complete.

A natural question arises: does every space have a completion?

To answer this, we need the following concept:

Definition 58 (Uniformly Continuous Functions)

We say that a function $f : (X, d_X) \rightarrow (Y, d_Y)$ is *uniformly continuous* if

$$\forall \varepsilon > 0 \exists \delta > 0 \forall x_1, x_2 \in X \\ d_X(x_1, x_2) < \delta \implies d_Y(f(x_1), f(x_2)) < \varepsilon.$$

Example 24.1.1

Given (X, d) , and $x_0 \in X$, define

$$g_{x_0}(x) = d(x, x_0).$$

Note that $|d(x_0, x) - d(x_0, y)| \leq d(x, y)$.¹ Thus


¹ Proved in A3

$$|g_{x_0}(x_1) - g_{x_0}(x_2)| \leq d(x_1, x_2).$$


Then $\forall \varepsilon > 0 \exists \delta = \varepsilon > 0$, we have

$$d(x_1, x_2) < \delta \implies |g_{x_0}(x_1) - g_{x_0}(x_2)| < \varepsilon.$$

Thus g_{x_0} is uniformly continuous. 

 Theorem 59 (Completion Theorem)

Every metric space (X, d) has a completion.

 Proof

Let $a \in X$. Define $\varphi : X \rightarrow C_b(X)$ by

$$(\varphi(u))(x) = f_u(x) = d(u, x) - d(x, a).$$

By our earlier example, $\varphi(u)$ is continuous. Notice that we have

$$|f_u(x)| = |d(u, x) - d(x, a)| \leq d(u, a).$$

Thus $\varphi(u) \in C_b(X)$, proving that φ is well-defined.

WTS φ is an isometry. Let $u, v \in X$. Then

$$\begin{aligned} |f_u(x) - f_v(x)| &= |d(u, x) - d(x, a) - d(v, x) + d(x, a)| \\ &= |d(u, x) - d(v, x)| \\ &\leq d(u, v). \end{aligned}$$

Thus $\|f_u - f_v\|_\infty \leq d(u, v)$ by definition of $\|\cdot\|_\infty$. Notice that


$$|f_u(v) - f_v(v)| = d(u, v),$$

which gives us the greatest possible value. Thus

$$\|\varphi(u) - \varphi(v)\|_\infty = \|f_u - f_v\|_\infty = d(u, v).$$

Thus φ is an isometry.

Since $(C_b(X), \|\cdot\|_\infty)$ is a complete metric space, let $Y = \overline{\varphi(X)}$.

The proof is complete by  Proposition 58.

QUESTION: If (X, d) has 2 completions, how are they related?

Suppose (X, d) is a metric space that has 2 completions through

the functions φ and ψ .




Figure 24.1: Relation of the 2 completions of a metric space.

Since we have that φ is bijective from X to $\varphi(X)$, we can take its inverse. Consequently, we have that the function $\Gamma = \psi \circ \varphi^{-1}$ is an isometry.

Now for some $\{x_n\} \subset X$ that is Cauchy, we know that in $\varphi(X)$, $\varphi(x_n) \rightarrow y_0 \in \varphi(X)$. Note that y_0 is a limit point of $\varphi(X)$. Through Γ , we have that

$$\Gamma(\varphi(x_n)) = \psi(x_n).$$

If $\psi(x_n) \rightarrow z_0 \in \psi(X)$, then we must have

$$\Gamma(y_0) = z_0,$$

and in particular z_0 is a limit point of $\psi(X)$. This forces limit points of $\varphi(X)$ to also be limit points of $\psi(X)$, and interior to interior. Thus the two completions are isomorphic.

24.2 Banach Contractive Mapping Theorem

QUESTION: Does there exist a function $f \in C[0, 1]$ such that


$$f(x) = e^x + \int_0^x \frac{\sin t}{2} f(t) dt \quad ? \quad (24.1)$$

Let $\Gamma : C[0, 1] \rightarrow C[0, 1]$ such that

$$\Gamma(f)(x) = e^x + \int_0^x \frac{\sin t}{2} f(t) dt.$$

Then f_0 is a solution to Equation (24.1) iff $\Gamma(f_0) = f_0$.

This is known as an **integral transform**.

 **Definition 59 (Fixed Point)**

Given (X, d) , $\Gamma : X \rightarrow X$, we say that x_0 is a fixed point of Γ if $\Gamma(x_0) = x_0$.

25 Lecture 25 Nov 09th

25.1 Banach Contractive Mapping Theorem (Continued)

Definition 60 (Lipschitz)

A function $f : (X, d_X) \rightarrow (Y, d_Y)$ is said to be Lipschitz if there exists $\alpha \geq 0$ such that $\forall x_1, x_2 \in X$,

$$d_Y(f(x_1), f(x_2)) \leq \alpha d_X(x_1, x_2)$$

Definition 61 (Contraction)

A function $f : X \rightarrow Y$ is called a **contraction** if there exists $0 \leq k < 1$ with

$$d_Y(f(x_1), f(x_2)) \leq k d_X(x_1, x_2)$$


for all $x_1, x_2 \in X$.

Note 25.1.1


Notice that a Lipschitz function is uniformly continuous: choose $\delta = \frac{\epsilon}{\alpha}$.

Exercise 25.1.1

Prove that if $f : [a, b] \rightarrow \mathbb{R}$ and f' is continuous, then by the **Extreme Value Theorem** and the **Mean Value Theorem**, f is Lipschitz.

 Theorem 60 (Banach Contractive Mapping Theorem)

Assume that (X, d) is complete. If $\Gamma : X \rightarrow X$ is contractive, then there exists a unique $x_0 \in X$ such that $\Gamma(x_0) = x_0$.

 Proof

Pick $x_1 \in X$. Then, let

$$x_2 = \Gamma(x_1), x_3 = \Gamma(x_2), \dots, x_{n+1} = \Gamma(x_n), \dots$$

Claim: $\{x_n\}$ is Cauchy¹

Let $k \in \mathbb{R}$ such that $0 < k < 1$, so that we have

$$d(\Gamma(x), \Gamma(y)) \leq kd(x, y)$$

for any $x, y \in X$. Then

$$\begin{aligned} d(x_3, x_2) &= d(\Gamma(x_2), \Gamma(x_1)) \leq kd(x_2, x_1) \\ d(x_4, x_3) &= d(\Gamma(x_3), \Gamma(x_2)) \leq kd(x_3, x_2) \leq k^2d(x_2, x_1) \\ &\vdots \\ d(x_{n+1}, x_n) &= d(\Gamma(x_n), \Gamma(x_{n-1})) \leq k^{n-1}d(x_2, x_1) \\ &\vdots \end{aligned}$$

Also, notice that if $m > n$, then

$$\begin{aligned} d(x_m, x_n) &\leq d(x_m, x_{m-1}) + d(x_{m-1}, x_{m-2}) + \dots + d(x_{n+1}, x_n) \\ &\leq k^{m-2}d(x_2, x_1) + k^{m-3}d(x_2, x_1) + \dots + k^{n-1}d(x_2, x_1) \\ &= \sum_{j=n-1}^{m-2} k^j d(x_2, x_1) = \frac{k^{n-1}}{1-k} d(x_2, x_1). \end{aligned}$$

Since $k^{n-1} \rightarrow 0$, we have that $\{x_n\}$ is Cauchy. Since (X, d) is complete, $\exists x_0 \in X$ such that $x_n \rightarrow x_0$.

In particular, we have that $x_{n+1} \rightarrow x_0$, i.e. $\Gamma(x_n) \rightarrow x_0$. Since Γ is continuous, we must have that $\Gamma(x_n) \rightarrow \Gamma(x_0)$. Therefore $\Gamma(x_0) = x_0$

¹ This will CTP since (X, d) is complete, i.e. it will give us a limit point at which Γ must converge to, and thus forcing its iteration to be terminated at the limit point due to Γ being contractive.

as required.

(Uniqueness) Suppose there exists another point $y_0 \in X$ such that $\Gamma(y_0) = y_0$. Then


$$d(x_0, y_0) = d(\Gamma(x_0), \Gamma(y_0)) \leq kd(x_0, y_0),$$

which implies that $d(x_0, y_0) = 0$.

Example 25.1.1

Show that the equation

$$f_0(x) = e^x + \int_0^x \frac{\sin t}{2} f_0(t) dt$$

has a unique solution in $C[0, 1]$. 

Solution


Define $\Gamma : C[0, 1] \rightarrow C[0, 1]$ by

$$\Gamma(f)(x) = e^x + \int_0^x \frac{\sin t}{2} f(t) dt.$$

Let $f, g \in C[0, 1]$. We have that

$$\begin{aligned} |\Gamma(f)(x) - \Gamma(g)(x)| &= \left| \int_0^x \frac{\sin t}{2} f(t) dt - \int_0^x \frac{\sin t}{2} g(t) dt \right| \\ &= \left| \int_0^x \frac{\sin t}{2} (f(t) - g(t)) dt \right| \\ &\leq \int_0^x \left| \frac{\sin t}{2} \right| |f(t) - g(t)| dt \\ &\leq \|f - g\|_\infty \int_0^1 \frac{1}{2} dt \\ &= \frac{1}{2} \|f - g\|_\infty \end{aligned}$$


Thus $\|\Gamma(f) - \Gamma(g)\|_\infty \leq \frac{1}{2} \|f - g\|_\infty$. Thus Γ is contractive. By

 Theorem 60, the unique fixed point is the solution.

Example 25.1.2

Show that the equation

$$f(x) = x + \int_0^x t^2 f(t) dt \tag{25.1}$$

has a unique solution. 

 **Solution**

Let $\Gamma(f)(x) = x + \int_0^x t^2 f(t) dt$. Then

$$\begin{aligned} |\Gamma(f)(x) - \Gamma(g)(x)| &\leq \int_0^1 t^2 \|f - g\|_\infty dt \\ &= \frac{1}{3} \|f - g\|_\infty. \end{aligned}$$

By the Banach Contractive Mapping Theorem, Equation (25.1) has a unique solution. In particular,

$$\begin{aligned} f_1(x) &= x \\ f_2(x) &= \Gamma(f_1)(x) = x + \int_0^x t^2 t_1(t) dt \\ &= x + \int_0^x t^3 dt = x + \frac{1}{4}x^4 \\ f_3(x) &= \Gamma(f_2)(x) = x + \int_0^x t^2 \left(t + \frac{1}{4}t^4 \right) dt \\ &= x + \int_0^x t^3 + \frac{1}{4}t^6 dt = x + \frac{1}{4}x^4 + \frac{1}{4 \cdot 7}x^7 \\ &\vdots \\ f_n(x) &= \frac{x}{1} + \frac{x^4}{4} + \frac{x^7}{4 \cdot 7} + \dots + \frac{x^{3n-2}}{4 \cdot 7 \cdot \dots \cdot (3n-2)} \end{aligned}$$

and so the limit is

$$f_0(x) = \sum_{k=1}^{\infty} \frac{x^{3k-2}}{4 \cdot 7 \cdot \dots \cdot (3k-2)}.$$


Example 25.1.3 (Other Applications)

1. Newton's Method.
2. **(Picard's Theorem)** Let $f : [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$ be Lipschitz in \mathbb{R} , i.e. $\exists \alpha \geq 0$ such that

$$|f(t, y_1) - f(t, y_2)| \leq \alpha |y_1 - y_2|$$

for any $y_1, y_2 \in \mathbb{R}$. If $y_0 \in \mathbb{R}$, then there exists a unique $\varphi \in C[a, b]$ such that

$$\varphi'(t) = f(t, \varphi(t))$$


for all $t \in (a, b)$ with $\varphi(a) = y_0$. 

25.2 Baire Category Theorem

Example 25.2.1 (Dirichlet Function)

Consider the function

$$f(x) = \begin{cases} 0 & x \in \mathbb{R} \setminus \mathbb{Q} \\ 1 & x = 0 \\ \frac{1}{m} & x \in \mathbb{Q} \end{cases}$$

The function g is continuous at each $x \in \mathbb{R} \setminus \mathbb{Q}$, and discontinuous otherwise. 

QUESTION: Does there exist a function f such that f is continuous on \mathbb{Q} but not on $\mathbb{R} \setminus \mathbb{Q}$? **No!**

However, to prove that there is need no such function, we need more machinery. In particular, the set of discontinuities of a function $f : (X, d) \rightarrow \mathbb{R}$ has a particular topological nature.

Definition 62 (Points of Discontinuity)

Let $f : X \rightarrow \mathbb{R}$. For each $n \in \mathbb{N}$, the *points of discontinuity* is a set defined as

$$D_N(f) = \left\{ x_0 \in X : \forall \delta > 0 \exists x_1, y_1 \in B(x_0, \delta) \ |f(x_1) - f(y_1)| \geq \frac{1}{n} \right\}.$$

Note 25.2.1

1. For each $n \in \mathbb{N}$, D_n is closed.
2. f is continuous at $x_0 \iff x_0 \notin \bigcap_{n=1}^{\infty} D_n$.

Remark 25.2.1

Recall the definition of an F_σ -set from the midterm (definition also provided in next lecture).

The set

$$D(f) = \{x_0 \in X \mid f \text{ is discontinuous at } x_0\} = \bigcap_{n=1}^{\infty} D_n(f)$$

is an F_σ -set.



A natural question to ask is:

QUESTION: Is $\mathbb{R} \setminus \mathbb{Q}$ an F_σ -set?

26 Lecture 26 Nov 12th

26.1 Baire Category Theorem (Continued)

Definition 63 (F_σ Sets)

Let (X, d) be a metric space. We say that $A \subseteq X$ is F_σ if there exists a sequence $\{F_n\}_{n=1}^\infty$ of closed sets with

$$A = \bigcup_{n=1}^{\infty} F_n.$$

Definition 64 (G_δ Sets)

Let (X, d) be a metric space. We say that $A \subseteq X$ is G_δ if there exists a sequence $\{U_n\}_{n=1}^\infty$ of open sets such that

$$A = \bigcap_{n=1}^{\infty} U_n.$$

Example 26.1.1

The interval $[0, 1) \subset \mathbb{R}$ is G_δ , since

$$[0, 1) = \bigcap_{n=1}^{\infty} \left(\frac{1}{n}, 1 \right)$$



Remark 26.1.1

A is F_σ iff A^c is G_δ .



Recall the definition of a dense set. We have the following complementary definition.

Definition 65 (Nowhere Dense)

Given a metric space (X, d) , we say that $A \subseteq X$ is *nowhere dense* if $\overline{A}^\circ = \emptyset$.

Remark 26.1.2

The above definition is equivalent to saying that \overline{A}^c is dense. 

Definition 66 (First Category)

We say that a set A is of *first category* if


$$A = \bigcup_{n=1}^{\infty} A_n$$

where each A_n is nowhere dense.

Definition 67 (Second Category)

We say that A is of *second category* if A is not of first category.

Remark 26.1.3

We colloquially refer to a set of first category as being *topologically thin*, and a set of second category as being *topologically thick*. 

Definition 68 (Residual)

We say that $A \subseteq (X, d)$ is a *residual* in X if A^c is of first category.

Theorem 61 (Set of Points of Discontinuity is F_σ)

Let $f : (X, d_X) \rightarrow (Y, d_Y)$. Then for each $n \in \mathbb{N}$, $D_N(f)$ is closed in X .

Moreover,

$$D(f) = \bigcup_{n=1}^{\infty} D_N(f).$$

In particular, $D(f)$ is F_σ .

Exercise 26.1.1

Prove **Theorem 61**.

Example 26.1.2

If $F \subset (X, d)$ is closed, then f is G_δ . In particular, notice that

$$F = \bigcap_{n=1}^{\infty} \left(\bigcup_{x \in F} B\left(x, \frac{1}{n}\right) \right),$$

where we note that each of the $B\left(x, \frac{1}{n}\right)$ is F_δ . ➔

Theorem 62 (Baire Category Theorem I)

Let (X, d) be complete. Let $\{U_n\}_{n=1}^{\infty}$ be a countable collection of dense open sets. Then¹

$$\bigcap_{n=1}^{\infty} U_n \text{ is dense in } X.$$

In particular, it is not empty.

¹Note that we have ourselves a dense G_δ set.

Proof

Assume that $\{U_n\}_{n=1}^{\infty}$ is a sequence of open and dense sets. Let $W \subset X$ be open and non-empty. Since U_1 is dense, we have that $W \cap U_1 \neq \emptyset$. Then $\exists x_1 \in W \cap U_1$ such that $\exists 0 < r_1 \leq 1$ so that

$$B(x_1, r_1) \subset B[x_1, r_1] \subset W \cap U_1.$$

Similarly,




Figure 26.1: Visualization of proof for Baire Category Theorem I

we can find $x_2 \in X$ such that for some $0 < r_2 \leq \frac{1}{2}$,

$$B(x_2, r_2) \subset B[x_2, r] \subset B(x_1, r_1) \cap U_2.$$

We can proceed recursively and find, for $n \in \mathbb{N}$, an $x_n \in X$ with $0 < r_n \leq \frac{1}{n}$ such that

$$B(x_n, r_n) \subset B[x_n, r_n] \subset B(x_{n-1}, r_{n-1}) \cap U_n.$$

Now since (X, d) is complete, $\{\text{diam}(B[x_n, r_n])\} = \{r_n\}$ is a decreasing sequence such that $r_n \rightarrow 0$, by Cantor's Intersection Principle,

$$\exists x_0 \in \bigcap_{n=1}^{\infty} B[x_n, r_n].$$

Then by this construction, we must have $x_0 \in B[x_1, r_1] \subset W \cap U_1$, and $x_0 \in B[x_n, r_n] \subset U_n$ for each $n \in \mathbb{N}$. Thus

$$x_0 \in W \cap \left(\bigcap_{n=1}^{\infty} U_n \right).$$



Note that the statement does not hold if we have an uncountable collection of dense open sets.

Example 26.1.3

Consider $U_x = \mathbb{R} \setminus \{x\}$, where $x \in \mathbb{R}$. This is clearly a dense and open set. Notice, however, that

$$\bigcap_{x \in \mathbb{R}} U_x = \emptyset.$$

Remark 26.1.4

 Theorem 62 shows that given a countable sequence $\{U_n\}_{n=1}^{\infty}$ of open dense sets of X , the countable intersection of these sets, $\bigcap_{n=1}^{\infty} U_n$, is a dense G_{δ} . 

If (X, d) is complete, then X is of second category.

 **Proof**

Suppose to the contrary that $X = \bigcup_{n=1}^{\infty} A_n$ where each A_n is nowhere dense. Since each A_n is nowhere dense, we have that

$$X = \bigcup_{n=1}^{\infty} A_n = \bigcup_{n=1}^{\infty} \overline{A_n}.$$

Let $U_n = \overline{A_n}^c$, which would then be open and dense, as X is complete. However, by De Morgan's Laws, we have that

$$\left(\bigcap_{n=1}^{\infty} U_n \right)^c = \bigcup_{n=1}^{\infty} U_n^c = \bigcup_{n=1}^{\infty} \overline{A_n} = X$$

and so


$$\bigcap_{n=1}^{\infty} U_n = \emptyset,$$

which is impossible by  Theorem 62.


Example 26.1.4

There are set that are neither F_σ or G_δ . For instance, consider the union of positive rationals and negative irrationals, i.e. a set

$$S = \mathbb{Q}_{>0} \cup \mathbb{Q}_{<0}^c.$$

If S is a G_δ , then by the Baire Category Theorem, $S \cap (0, \infty)$ is also G_δ , but that's the set of positive rationals, which cannot be G_δ . Similarly, if S were F_σ , then its intersection with $(-\infty, 0)$ is also F_σ , but the set of negative irrationals cannot be F_σ . Thus S is neither F_σ nor G_δ . 

Example 26.1.5

\mathbb{R} and $\mathbb{R} \setminus \mathbb{Q}$ are of second category. In fact, $\mathbb{R} \setminus \mathbb{Q}$ is a residual, since \mathbb{Q} is of first category. 

QUESTION: Is

$$Q = \bigcap_{k=1}^{\infty} \bigcup_{n=1}^{\infty} \left(r_n - \frac{1}{2^{k+n}}, r_n + \frac{1}{2^{k+n}} \right),$$

where $Q = \{r_1, r_2, \dots\}$, \mathbb{Q} ? No. Notice that this is fairly close, but it is not.²

² It should be \mathbb{R} ?

 **Corollary 64 (Q is not G_δ)**

Q is not a G_δ set.

 **Proof**

Suppose to the contrary that Q is G_δ , i.e. there exists a countable sequence of open sets $\{U_n\}$ such that

$$Q = \bigcap_{n=1}^{\infty} U_n.$$


Let $F_n = U_n^c$. Since Q is dense, it follows that each of the U_n 's is also dense. Thus F_n is nowhere dense and closed.

Let $Q = \{r_1, r_2, \dots\}$, an enumeration on Q , and $S_n = F_n \cup \{r_n\}$. Then S_n is closed and nowhere dense. However, we would then have

$$\mathbb{R} = \bigcup_{n=1}^{\infty} S_n,$$

which contradicts the fact that \mathbb{R} is of second category.

Consequently:

 **Corollary 65 (There are no Functions Discontinuous on all Irrational Numbers)**

There is no function $f : \mathbb{R} \rightarrow \mathbb{R}$ for which $D(f) = \mathbb{R} \setminus \mathbb{Q}$.

We are now able to show that for a sequence $\{f_n\} \subset C[a, b]$ that converges pointwise, the limit function must be continuous at each point on a residual set. We require the following notion:

Definition 69 (Uniformly Convergent Sequence of Functions on a Point)

We say that a sequence of functions $\{f_n\}$ where,

$$f_n : (X, d_X) \rightarrow (Y, d_Y),$$

converges uniformly at $x_0 \in X$ if

$$\forall \varepsilon > 0 \exists \delta > 0 \exists N \in \mathbb{N} \forall n, m \geq N \\ x \in B(x_0, \delta) \implies d_Y(f_n(x), f_m(x)) < \varepsilon.$$

The proof of the following theorem is left as an exercise.

Theorem 66 (Limit of Sequence of Continuous Functions that Converges Pointwise is Continuous)

Let (X, d_X) and (Y, d_Y) be metric spaces. Let $\{f_n : X \rightarrow Y\}$ be a sequence of functions that converges pointwise on X to f_0 . Assume that $\{f_n\}$ converges uniformly at $x_0 \in X$. If each f_n is continuous at x_0 , then so is f_0 .

27 Lecture 27 Nov 14th

27.1 Baire Category Theorem (Continued 2)

Theorem 67 (Uniform Convergence of A Sequence of Continuous Functions that Converges Pointwise)

Let $f_n : (a, b) \rightarrow \mathbb{R}$ be a sequence of continuous functions that converges pointwise to $f(x)$. Then there exists an $x_0 \in (a, b)$ such that $f_n \rightarrow f$ uniformly at x_0 .

Proof

Assume that $f_n \rightarrow f_0$ on (a, b) , pointwise.

Claim There exists $[\alpha_1, \beta_1] \subset (a, b)$ and $N_1 \in \mathbb{N}$ such that if $x \in [\alpha_1, \beta_1]$ and $n, m \geq N_1$, then $|f_n(x) - f_m(x)| \leq 1$.

Suppose not. Then $\exists t_1 \in (a, b)$ and $n_1, m_1 \in \mathbb{N}$ such that $|f_{n_1}(t_1) - f_{m_1}(t_1)| > 1$. Since $f_{n_1} - f_{m_1}$ is continuous, there exists an open interval $I_1 \subsetneq \bar{I}_1 \subsetneq (a, b)$ such that $|f_{n_1}(x) - f_{m_1}(x)| > 1$ for all $x \in I_1$.

Similarly, $\exists t_2 \in I_1$ and $n_2, m_2 \geq \max\{n_1, m_1\}$ such that $|f_{n_2}(t_2) - f_{m_2}(t_2)| > 1$. Again, since $f_{n_2} - f_{m_2}$ is continuous, there exists an open interval $I_2 \subsetneq \bar{I}_2 \subsetneq I_1$ such that $|f_{n_2}(x) - f_{m_2}(x)| > 1$ for all $x \in I_2$.

Recursively so, we get a sequence $\{I_n\}$ of open interval with $I_{n+1} \subset \bar{I}_{n+1} \subset \bar{I}_n$, and two sequence of integers $\{n_k\}$ and

$\{m_k\}$, with $n_{k+1}, m_{k+1} \geq \max\{n_k, m_k\}$ and if $x \in I_k$, we have $|f_{n_k}(x) - f_{m_k}(x)| > 1$.

Then, by the Nested Interval Theorem, we have

$$\bigcap_{k=1}^{\infty} \bar{I}_k \neq \emptyset.$$

Let $x^* \in \bigcap_{k=1}^{\infty} \bar{I}_k$. Then by construction, we have that for any k , $|f_{n_k}(x^*) - f_{m_k}(x^*)| > 1$. However, since $\{f_n\}$ converges pointwise, $\{f_n(x^*)\}$ is Cauchy and hence we have a contradiction. This proves the claim \neg .

In a similar manner, we can find a sequence $\{[\alpha_k, \beta_k]\}$ of closed sets, where $\alpha_k < \beta_k$, such that

$$(\alpha_{k+1}, \beta_{k+1}) \subseteq [\alpha_{k+1}, \beta_{k+1}] \subseteq (\alpha_k, \beta_k) \subseteq \dots \subseteq (a, b),$$

and a sequence

$$N_1 < N_2 < \dots < N_k < \dots,$$

such that if $x \in [\alpha_k, \beta_k]$ and $n, m \geq N_k$, then $|f_n(x) - f_m(x)| \leq \frac{1}{k}$. Then, once again, by the Nested Interval Theorem, let $x_0 \in \bigcap_{k=1}^{\infty} [\alpha_k, \beta_k]$. Let $\varepsilon > 0$. Now if $\frac{1}{k} < \varepsilon$, then if $n, m \geq N_k$, then we have

$$|f_n(x) - f_m(x)| \leq \frac{1}{k} < \varepsilon.$$

Since $x_0 \in \bigcap_{k=1}^{\infty} [\alpha_k, \beta_k]$ and $\alpha_k < \beta_k$, we can choose $\delta = \min\{\beta_k - \alpha_k : k \in \mathbb{N} \setminus \{0\}\} > 0$, so that $(x_0 - \delta, x_0 + \delta) \subset (\alpha_k, \beta_k)$, then for any $x \in (x_0 - \delta, x_0 + \delta)$, we have


$$|f_n(x) - f_m(x)| < \varepsilon.$$

► Corollary 68 (Continuity of the Limit of a Sequence of Pointwise Convergent Functions on a Residual Set)


Let $\{f_n\} \subset C[a, b]$ be such that $f_n \rightarrow f_0$ pointwise on $[a, b]$. Then there exists a residual set $A \subset [a, b]$ such that $f_0(x)$ is continuous at each


$x \in A$.

 **Proof**

 Theorem 67 shows that the set A of which f_0 is continuous on is dense in $[a, b]$. However, from XXX that $D(f_0)$ is F_σ , and so A is a dense G_δ . □

Remark 27.1.1

Thus we have that $D(f_0)$ is a nowhere dense F_σ , i.e. it is of first category. 

 **Corollary 69 (Derivative of a Function is Continuous on a dense G_δ set in \mathbb{R})**

Assume that $f : \mathbb{R} \rightarrow \mathbb{R}$ is differentiable. Then $f'(x)$ is continuous for every point on a dense G_δ -subset of \mathbb{R} .

 **Proof**

Using notions from the first principles of calculus, notice that $f'(x)$ is a pointwise limit of the sequence of continuous functions

$$\left\{ \frac{f\left(x + \frac{1}{n}\right) - f(x)}{\frac{1}{n}} \right\}.$$

27.2 Compactness

In this section, we study 3 important properties of a topological space, namely:

- compactness;

- sequential compactness; and
- the Bolzano-Weierstrass Property.

We shall see that, in fact, the three properties are equivalent.

Definition 70 (Cover)

Given (X, d) a metric space, an (open) **cover** of X is a collection $\{U_\alpha\}_{\alpha \in I}$ of open sets with

$$X = \bigcup_{\alpha \in I} U_\alpha.$$

A **subcover** is a subset (or subcollection) $\{U_\alpha\}_{\alpha \in J \subset I}$ such that

$$X = \bigcup_{\alpha \in J} U_\alpha.$$

If $A \subset X$, then we say that $\{U_\alpha\}_{\alpha \in I}$ **covers** A if $A \subset \bigcup_{\alpha \in I} U_\alpha$, or, equivalently, if $\{U_\alpha \cap A\}_{\alpha \in I}$ is a cover of (A, d_A) .

Definition 71 (Compact)

We say that (X, d) is **compact** iff each cover of X , $\{U_\alpha\}_{\alpha \in I}$, has a finite subcover.

We say that $A \subset (X, d)$ is **compact** if every cover $\{U_\alpha\}_{\alpha \in I}$ of A has a finite subcover (or, equivalently, if (A, d_A) is compact).

From earlier courses in Calculus, recall:

Theorem 70 (Heine-Borel Theorem)

$A \subset \mathbb{R}^n$ is compact iff A is closed and bounded.

Example 27.2.1

$[0, 1] \subset \mathbb{R}$ is compact, but $(0, 1) \subset \mathbb{R}$ is not compact.



However, the Heine-Borel Theorem is not true for arbitrary metric spaces.


Example 27.2.2 (★)

Let

$$A = \{\{x_n\} \in \ell_\infty \mid \|x_n\|_\infty \leq 1\}.$$

It is clear that A is closed and bounded. However, consider $U_{\{x_n\}} = B\left(\{x_n\}, \frac{1}{2}\right)$. It is then clear that

$$A \subset \bigcup_{\{x_n\} \in A} U_{\{x_n\}}.$$

Let $S = \{\{x_n\} \mid x_n = 1 \vee x_n = 0\}$, which is infinite. Then we notice that $|S \cap B\left(\{x_n\}, \frac{1}{2}\right)| \leq 1$, showing to us that we cannot find a finite subcover for S itself is infinite. 

However, we do have the following implication.

 **Proposition 71 (Compact Spaces are Closed and Bounded)**

If $A \subset (X, d)$ is compact, then A is closed and bounded.

 **Proof**

Suppose A is not closed. Then $\exists x_0 \in \text{bdy}(A) \setminus A$. Let

$$U_n = \left(B \left[x_0, \frac{1}{n} \right] \right)^c.$$

Since $x_0 \notin A$, we have that $A \subset \bigcup_{n=1}^{\infty} U_n$. However, $\{U_n\}_{n=1}^{\infty}$ has no finite subcover. Otherwise, if it does have some finite subcover, say $\{U_n\}_{n=1}^N$, then for any $n_0 > N$, we would have that

$$\left(B \left[x_0, \frac{1}{n_0} \right] \right) \not\supseteq \bigcup_{n=1}^N U_n,$$

and so $\exists x_1 \in B \left[x_0, \frac{1}{n_0} \right]$ such that $x_1 \in A$ but $x_0 \notin \bigcup_{n=1}^N U_n$. This contradicts the assumption that a subcover exists. But A must have some subcover for we assumed that A is compact. Therefore A

must be closed.

For boundedness, let $x_0 \in X$. Then $\{B(x_0, n)\}_{n=1}^{\infty}$ is an open cover of A . Since A is compact, $\{B(x_0, n)\}_{n=1}^{\infty}$ must have some finite subcover $\{B(x_0, n_1), B(x_0, n_2), \dots, B(x_0, n_k)\}$. WMA $n_1 < n_2 < \dots < n_k$, for we may rearrange the radii. It follows that $A \subset B(x_0, n_k)$, and so A is bounded as required.

28 Lecture 28 Nov 16th

28.1 Compactness (Continued)

We also have the following relation between compact sets and their closed subsets.

Proposition 72 (Closed Subsets of Compact Sets are Compact)

If (X, d) is compact and A is closed, then A is compact.

Proof

Let $\{U_\alpha\}_{\alpha \in I}$ be a cover of A . Then

$$\{U_\alpha\}_{\alpha \in I} \cup A^C \tag{*}$$

is a cover of X . Since X is compact, Equation (*) has a finite subcover $\{U_{\alpha_1}, U_{\alpha_2}, \dots, U_{\alpha_k}, A^C\}$ such that

$$\left(\bigcup_{i=1}^k U_{\alpha_i} \right) \cup A^C = X.$$

Since $A \subset X$ and $A \cap A^C = \emptyset$, we must have

$$A \subset \bigcup_{i=1}^k U_{\alpha_i}.$$

We have the following 2 variants of compactness:

Definition 72 (Sequential Compactness)

A set $A \subset (X, d)$ is said to be **sequentially compact** if every sequence¹ $\{x_n\} \subset A$ has a subsequence $\{x_{n_k}\}$ such that $x_{n_k} \rightarrow x_0 \in A$.

¹ Beware that this is not the same as completeness.

Definition 73 (Bolzano-Weierstrass Property (BWP))

Let (X, d) be a metric space. We say that X has the **Bolzano-Weierstrass Property (BWP)** if every infinite subset of X has a limit point in the subset.

Exercise 28.1.1

Show that for $A \subset \mathbb{R}^n$, A is compact iff A is sequentially compact.

Proof

(\implies) Suppose A is not sequentially compact. Then

$$\exists \{x_n\} \subset A \forall \{x_{n_k}\} \subset \{x_n\} \forall x_0 \in A \quad x_{n_k} \not\rightarrow x_0.$$

Let this $\{x_n\} = \{x_1, x_2, \dots, x_n, \dots\}$. Let

$$U_n = A \setminus \{x_j \mid j \geq n\}.$$

Then it is clear that

$$\bigcup_{n=1}^{\infty} U_n = A,$$

i.e. $\{U_n\}$ is a cover of A . Since A is compact, $\{U_n\}$ has a finite subcover, say $\{U_{n_1}, U_{n_2}, \dots, U_{n_k}\}$. WMA $n_1 < n_2 < \dots < n_k$. Then

$$A = \bigcup_{m=1}^k U_{n_m} = A \setminus \{x_j \mid j \geq n_k\}.$$


But that is impossible since $x_{n_k+1} \notin \bigcup_{m=1}^k U_{n_m}$. Thus A must be

sequentially compact.

(\Leftarrow) Suppose A is sequentially compact. Then

$$\forall \{x_n\} \subset A \exists \{x_{n_k}\} \subset \{x_n\} \exists x_0 \in A \ x_{n_k} \rightarrow x_0.$$

Let $\{U_\alpha\}_{\alpha \in I}$ be a cover of A . **Yet to figure out where to go from here.** Tried looking into trying to construct a finite subcover using the convergent subsequence, but that actually leads to nowhere. \square

 **Theorem 73 (Sequential Compactness is Equivalent to BWP)**

Let (X, d) be a metric space. TFAE:

1. (X, d) is sequentially compact.
2. (X, d) has the BWP.

 **Proof**

(\Rightarrow) Let (X, d) be sequentially compact. Let $A \subset (X, d)$ be infinite. By sequential compactness, every sequence $\{x_n\} \subset A$ has a convergent subsequence $\{x_{n_k}\}$, such that $x_{n_k} \rightarrow x_0 \in A$. \dashv

(\Leftarrow) Suppose (X, d) has the BWP. Let $\{x_n\}$ be a sequence in X . If $\{x_n\}$ is not infinite (as a set), then it has a subsequence $\{x_{n_k}\}$ such that $x_{n_{k_1}} = x_{n_{k_2}}$ for all k_1, k_2 , which is convergent. WMA $\{x_n\}$ is infinite (as a set). By the BWP, $\{x_n\}$ (as a set) has a limit point $x_0 \in \{x_n\}$. Then for $k \in \mathbb{N} \setminus \{0\}$, let

$$x_{n_k} \in B\left(x_0, \frac{1}{k}\right).$$

Clearly then $x_{n_k} \rightarrow x_0$, and $\{x_{n_k}\}$ is a subsequence of $\{x_n\}$.


 **Definition 74 (Finite Intersection Property (FIP))**

A collection $\{A_\alpha\}_{\alpha \in I}$ of subsets of X is said to have the *finite intersection property (FIP)* if

$$\bigcap_{i=1}^n A_n \neq \emptyset$$

for all finite subcollections $\{A_1, \dots, A_n\}$.

Example 28.1.1

Let $F_n = [n, \infty)$. Then $\{F_n\}_{n=1}^\infty$ has the FIP, but $\bigcap_{n=1}^\infty F_n = \emptyset$. 

The following theorem can be seen as an upgrade to Cantor's Intersection Principle for compact metric spaces: instead of allowing only a countably infinite intersection, we can now take an arbitrary number of intersections.


Theorem 74 (FIP and Compactness)

Let (X, d) be a metric space. TFAE:

1. (X, d) is compact.
2. If $\{F_\alpha\}_{\alpha \in I}$ is a non-empty collection of closed sets with the FIP, then

$$\bigcap_{\alpha \in I} F_\alpha \neq \emptyset.$$

Remark 28.1.1

As compared to Cantor's Intersection Principle, we do not need the notion of a diameter of a set to achieve this result in a compact set. 

Proof

(1) \implies (2) Suppose to the contrary that for a non-empty collection $\{F_\alpha\}_{\alpha \in I}$ of closed sets with the FIP, we have

$$\bigcap_{\alpha \in I} F_\alpha = \emptyset.$$

Let $U_\alpha = F_\alpha^C$. Then by De Morgan's Laws, we have $X = \bigcup_{\alpha \in I} U_\alpha$. Since (X, d) is compact, $\exists \{U_{\alpha_1}, \dots, U_{\alpha_n}\}$ such that

$$\bigcup_{i=1}^n U_{\alpha_i} = X.$$

But that implies that

$$\emptyset = X^C = \left(\bigcup_{i=1}^n U_{\alpha_i} \right)^C = \bigcap_{i=1}^n F_{\alpha_i},$$

contradicting FIP.

(2) \implies (1) Suppose to the contrary that $\{U_\alpha\}_{\alpha \in I}$, a cover of X , has no finite subcover. Then $\forall \{U_{\alpha_1}, \dots, U_{\alpha_n}\}$, we must have

$$X \setminus \bigcup_{i=1}^n U_{\alpha_i} \neq \emptyset,$$

i.e., by De Morgan's Laws, $\bigcap_{i=1}^n U_{\alpha_i}^C \neq \emptyset$. Then $\{F_\alpha\}_{\alpha \in I}$, where $F_\alpha = U_\alpha^C$, is a non-empty collection of closed sets with the FIP (by our argument), but via De Morgan's Laws, we have

$$\bigcap_{\alpha \in I} F_\alpha = \emptyset,$$

contradicting our assumption.

► Corollary 75 (Generalized Nested Interval Theorem for Compact Metric Spaces)

Let (X, d) be compact and $\{F_N\}_{N=1}^\infty$ be a sequence of non-empty closed sets such that $F_{n+1} \subset F_n$. Then

$$\bigcap_{n=1}^\infty F_n \neq \emptyset.$$

► Corollary 76 (Compact Metric Spaces are Complete)

If (X, d) is compact, then (X, d) is complete.

Note 28.1.1

RECALL the definition for compactness, in which we may then have the following notion: for a compact set (X, d) , for $\varepsilon > 0$, since $\{B(x, \varepsilon)\}_{x \in X}$ is an open cover of X , we know that there exists $x_1, \dots, x_n \in X$ such that they form a finite subcover on X .

$$X = \bigcup_{i=1}^n B(x_i, \varepsilon).$$

We use the same idea and make the following definition:

Definition 75 (ε -net)

Given $A \subset (X, d)$ and $\varepsilon > 0$. An ε -net for A is a set $\{x_\alpha\}_{\alpha \in I} \subset X$ such that

$$A \subset \bigcup_{\alpha \in I} B(x_\alpha, \varepsilon).$$

Definition 76 (Totally Bounded)

We say that a subset $A \subset (X, d)$ is **totally bounded** if A has a finite ε -net for every $\varepsilon > 0$.

Theorem 77 (Compact Sets are Totally Bounded)

If (X, d) is compact, then (X, d) is totally compact.

Proof


The proof immediately follows from the definition of compactness, as discussed in Note 28.1.1. □


Note that bounded and totally bounded are **not equivalent**.

Example 28.1.2


Let

$$S = \{\{x_n\} \in \ell_\infty \mid \|\{x_n\}\|_\infty \leq 1\}.$$

We have that S is bounded, but it does not have a $\frac{1}{2}$ -net. 


 **Proposition 78 (A Set is Totally Bounded iff Its Closure is Totally Bounded)**

$A \subset (X, d)$ is totally bounded iff \bar{A} is totally bounded.

 **Proof**

The (\Leftarrow) direction is immediate, since $A \subset \bar{A}$. It suffices to show for (\Rightarrow). Suppose A is totally bounded. If A is closed, then we are done, so WMA A is open. Then $\text{Lim}(A) \not\subset A$. Let $x_0 \in \text{Lim}(A) \setminus A$. Since x_0 is a limit point, for any $\varepsilon > 0$, $B(x_0, \varepsilon) \cap A \neq \emptyset$. **Need to verify definition of an ε -net.** □

Exercise 28.1.2

Prove  Proposition 78.

29 Lecture 29 Nov 19th

29.1 Compactness (Continued 2)

Theorem 79 (Compact Sets have BWP)

If (X, d) is compact, then (X, d) has the BWP.

Proof

Suppose $S \subset X$ is infinite. Then we can obtain a sequence $\{x_n\} \subset S$ such that for $n \neq m$, $x_n \neq x_m$. Then, consider

$$F_n = \{x_n, x_{n+1}, \dots\}.$$


We have that $F_{n+1} \subseteq F_n$ and we observe that $\{F_n\}$ has the FIP, i.e.

$$\exists x_0 \in \bigcap_{n=1}^{\infty} F_n.$$

Then for any $\varepsilon > 0$, for any $n \in \mathbb{N}$, we have that


$$B(x_0, \varepsilon) \subset F_n.$$

In fact, $B(x_0, \varepsilon) \cap \{x_n\} \neq \emptyset$ is also infinite. Thus $x_0 \in \text{Lim}(S)$.

 Proposition 80 (Sequential Compactness \implies Completeness and Total Boundedness)

If (X, d) is sequentially compact, then (X, d) is both complete and totally bounded.

Proof

Completeness Let $\{x_n\} \subset X$ be Cauchy. Then by the assumption that X is sequentially compact, $\{x_n\}$ has a subsequence $\{x_{n_k}\}$ such that $x_{n_k} \rightarrow x_0 \in X$. Then by  Theorem 47, $x_n \rightarrow x_0$. \dashv

Totally Bounded Suppose to the contrary that X is not totally bounded, i.e. $\exists \varepsilon_0 > 0$ such that X has no finite ε_0 -net. Then we can find $x_1 \in X$ such that $B(x_1, \varepsilon_0) \neq X$, an $x_2 \in X \setminus B(x_1, \varepsilon_0)$, $x_3 \in X \setminus (B(x_1, \varepsilon_0) \cup B(x_2, \varepsilon_2))$, and so on. In other words, we can construct a sequence $\{x_n\} \subset X$ such that $d(x_n, x_m) > \varepsilon$ for all $n \neq m$. Then by construction, $\{x_n\}$ has no convergent subsequences, i.e. X is not sequentially compact.

Theorem 81 (Continuity Preserves Sequential Compactness)

If (X, d) is sequentially compact and if $f : (X, d_X) \rightarrow (Y, d_Y)$ is continuous, then $f(X)$ is sequentially compact.

Proof

Let $\{y_n\} \subset f(X)$. Consider $\{x_n\}$ such that $f(x_n) = y_n$. Since X is sequentially compact, $\{x_n\}$ has a convergent subsequence $\{x_{n_k}\}$ with $x_{n_k} \rightarrow x_0$. Then by continuity,

$$y_{n_k} = f(x_{n_k}) \rightarrow f(x_0) = y_0.$$



Corollary 82 (Extreme Value Theorem)

If (X, d) is sequentially compact and $f : X \rightarrow \mathbb{R}$ is continuous, then
 $\exists c, d \in X$ such that

$$f(c) \leq f(x) \leq f(d)$$

for all $x \in X$.

 **Proof**

By  Theorem 81, $f(X)$ is sequentially compact in \mathbb{R} , and by
 Proposition 80, $f(X)$ is complete, and so by Heine-Borel, $f(X)$
 is closed and bounded. Thus

$$\sup(f(X)), \inf(f(X)) \in f(X).$$

 **Theorem 83 (Lesbesgue)**

Let (X, d) be sequentially compact. Let $\{U_\alpha\}_{\alpha \in I}$ be an open cover of X .
 Then $\exists \varepsilon > 0$ such that for every $0 < \delta < \varepsilon$, and every $x \in X$ such that
 for some $\alpha_0 \in I$

$$B(x_0, \delta) \subset U_{\alpha_0}.$$

 **Proof**

If $U_{\alpha_0} = X$, then any $\varepsilon > 0$ will work. WMA $U_\alpha \neq X$ for any $\alpha \in I$.
 Let $\varphi : X \rightarrow \mathbb{R}$ be defined by

$$\varphi(x) = \sup \{ \delta > 0 : B(x, \delta) \subseteq U_{\alpha_0}, \alpha_0 \in I \}.$$

Since $\{U_\alpha\}_{\alpha \in I}$ is an open cover of X , every x must be in one of the
 U_α 's, and so the set

$$\{ \delta > 0 : B(x, \delta) \subseteq U_{\alpha_0}, \alpha_0 \in I \}$$

is non-empty and $\varphi(x) > 0$. Also, $\varphi(x) < \infty$, since X is bounded

(as X is sequentially compact) and $U_\alpha \neq X$ for any $\alpha \in I$.

Now for any $x, y \in X$,¹ we have that

$$\varphi(x) \leq \varphi(y) + d(x, y)$$

by the Triangle Inequality. Thus

$$\varphi(x) - \varphi(y) \leq d(x, y)$$

and by symmetry we have

$$|\varphi(x) - \varphi(y)| \leq d(x, y).$$

Thus φ is Lipschitz, and so φ is uniformly continuous². Then by the Extreme Value Theorem, $\exists \varepsilon > 0$ such that $\exists \delta > 0$ such that $\varphi(x) \geq \varepsilon$ for all $x \in X$.

¹ I should check in with the professor on how to show this

² see note on definition of Lipschitz.

🗨 Note 29.1.1

The ε in Lesbesgue's Theorem is also called a *Lesbesgue Number*.

📖 Theorem 84 (Lesbesgue-Borel)

Let (X, d) be a metric space. TFAE:

1. (X, d) is compact.
2. (X, d) has BWP.
3. (X, d) is sequentially compact.

✎ Proof

We already have $(1) \implies (2)$ and $(2) \iff (3)$. It suffices to prove $(3) \implies (1)$. Let $\{U_\alpha\}_{\alpha \in I}$ be a cover of X . By Lesbesgue's Theorem, let $\varepsilon_0 > 0$, and fix $0 < \delta < \varepsilon_0$. Since (X, d) is totally


bounded (as it sequentially compact), there exists $\{x_1, \dots, x_n\}$ with

$$X = \bigcup_{i=1}^n B(x_i, \delta).$$

Then for each i , we have that $B(x_i, \delta) \subset U_{\alpha_i}$ for some $\alpha_i \in I$. Then

$$X = \bigcup_{i=1}^n U_{\alpha_i}$$



is a finite subcover of the cover $\{U_\alpha\}_{\alpha \in I}$.

 **Theorem 85 (Compactness \iff Completeness + Totally Bounded)**

Let (X, d) be a metric space. TFAE:

1. (X, d) is compact.
2. (X, d) is complete and totally bounded.

 **Proof**

By  Theorem 84 and  Proposition 80, we have $(1) \implies (2)$. Thus it suffices to show for $(2) \implies (1)$. Notice that we only need to show that (X, d) is sequentially compact. Let $\{x_n\} \subset (X, d)$.

Since (X, d) is totally bounded, X can be covered by finitely many open balls of radius 1. Thus one such ball $S_1 = B(y_1, 1)$, for some $y_1 \in X$, contains infinitely many terms in $\{x_n\}$ ³.

Similarly, X can be covered by finitely many open balls of radius $\frac{1}{2}$, and we can pick one of these open balls $S_2 = B\left(y_2, \frac{1}{2}\right)$ which contains infinitely many terms in $\{x_n\} \cap S_1$.

Recursively, we may construct a sequence of open balls

$$\left\{ S_k = B\left(y_k, \frac{1}{k}\right) \right\}$$

³ Note that sequences are infinitary by nature in our context.

with the property that each S_{k+1} contains infinitely many terms in

$$\{x_n\} \cap \left(\bigcap_{i=1}^k S_i \right).$$

Note that

$$\text{diam}(S_k) = \frac{2}{k} \rightarrow 0$$

as $k \rightarrow \infty$, and since can pick

$$n_1 < n_2 < \dots < n_k < \dots$$

such that

$$x_{n_k} \in \bigcap_{i=1}^k S_i.$$

WMA for some $N \in \mathbb{N}$, for any $k, m \geq N$, we have that $x_{n_k}, x_{n_m} \in S_N$, i.e.

$$d(x_{n_k}, x_{n_m}) \leq \text{diam}(S_N).$$

Thus $\{x_{n_k}\} \subset \{x_n\}$ is Cauchy. Since (X, d) is complete, $x_{n_k} \rightarrow x_0$, and therefore X is sequentially compact by definition.

30 Lecture 30 Nov 21st

30.1 Compactness (Continued 3)

The proof of the following theorem was left as an exercise:

Theorem 86 (Continuity Preserves Compactness)

If (X, d_X) is compact and $f : (X, d_X) \rightarrow (Y, d_Y)$ is continuous, then $f(X)$ is compact in Y .

Proof

The proof easily follows from  Theorem 84 and  Theorem 81.

30.2 Finite Dimensional Normed Linear Spaces


Definition 77 (Bounded Linear Map)

A linear map $T : (V, \|\cdot\|_V) \rightarrow (W, \|\cdot\|_W)$ is said to be **bounded** if

$$\|T\|_T = \sup\{\|T(v)\|_W \mid \|v\|_V \leq 1\} < \infty.$$


In assignment 3, we proved the following important result about

linear maps in finite dimensional normed linear spaces.


 **Theorem 87 (Boundedness is Equivalent to Continuity in Finite Dimensional Normed Linear Spaces)**

Let $T : (V, \|\cdot\|_V) \rightarrow (W, \|\cdot\|_W)$ be a linear map. TFAE:

1. T is bounded.
2. T is continuous.
3. T is continuous at 0.

 **Lemma 88 (Continuity of the Norm)**

The function $f : (V, \|\cdot\|) \rightarrow \mathbb{R}$ given by $f(x) = \|x\|$ is continuous.

 **Proposition 89 (Linear Map Between Spaces of Different Dimensions is Bounded)**

Let $T : (\mathbb{R}^n, \|\cdot\|_2) \rightarrow (\mathbb{R}^m, \|\cdot\|_2)$ be linear. Then T is bounded.

 **Proof**

Since T is a linear map, we may represent T using a matrix A such that


$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \vec{a}_1 \\ \vec{a}_2 \\ \vdots \\ \vec{a}_m \end{bmatrix}.$$

If $\|x\| \leq 1$, then

$$\|T(x)\|_2 = \left\| \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \right\| = \left\| \begin{bmatrix} \vec{a}_1 \cdot \vec{x} \\ \vec{a}_2 \cdot \vec{x} \\ \vdots \\ \vec{a}_m \cdot \vec{x} \end{bmatrix} \right\|$$


$$\begin{aligned}
 &= \left(\sum_{i=1}^m (\vec{a}_i \cdot \vec{x})^2 \right)^{\frac{1}{2}} \leq \left(\sum_{i=1}^m \|\vec{a}_i\|^2 \|\vec{x}\|^2 \right)^{\frac{1}{2}} \\
 &\leq \left(\sum_{i=1}^m \|\vec{a}_i\|^2 \right)^{\frac{1}{2}}.
 \end{aligned}$$


This completes the proof.

 **Theorem 90 (Boundedness of Functions between n -dimensional Vector Spaces and n -dimensional Normed Linear Spaces)**

Let $(V, \|\cdot\|_V)$ be an n -dimensional normed linear space with basis $\{v_1, \dots, v_n\}$. Let $\Gamma_n : \mathbb{R}^n \rightarrow V$ be given by

$$\Gamma_n(\alpha_1, \dots, \alpha_n) = \alpha_1 v_1 + \dots + \alpha_n v_n.$$

Then Γ_n and Γ_n^{-1} are both bounded. Furthermore, they are both continuous by  Theorem 87.


 **Proof**

Γ_n is bounded Suppose $\|(\alpha_1, \dots, \alpha_n)\|_2 \leq 1$. Then

$$\begin{aligned}
 \|\Gamma_n(\alpha_1, \dots, \alpha_n)\|_V &= \|\alpha_1 v_1 + \dots + \alpha_n v_n\|_V \\
 &\leq |\alpha_1| \|v_1\|_V + \dots + |\alpha_n| \|v_n\|_V \\
 &\leq \sum_{i=1}^n \|v_i\|_V.
 \end{aligned}$$

Γ_n^{-1} is bounded Note that since Γ_n is bounded, it is continuous. Consider

$$S = \{(\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n \mid \|(\alpha_1, \dots, \alpha_n) = 1\|_2 = 1\}.$$

Since S is closed and bounded, and is a subset of \mathbb{R}^n , S is compact by the Heine-Borel Theorem, and so $\Gamma(S)$ is compact in V by  Theorem 86. Since the mapping $v \rightarrow \|v\|_V$ is continuous, by the

Extreme Value Theorem,

$$\min\{\|\Gamma_n(\alpha_1, \dots, \alpha_n)\|_V \mid (\alpha_1, \dots, \alpha_n) \in S\} = \alpha > 0.$$

It follows by continuity that if $\|v\|_V \leq \alpha$, then $\|\Gamma_n^{-1}(v)\|_2 \leq 1$.

Therefore, we have that $\|\Gamma_n^{-1}\| \leq \frac{1}{\alpha}$.


Note 30.2.1

1. Γ_n is a homeomorphism.
2. As a consequence of Γ being continuous, we have that $\{x_n\}$ is Cauchy in \mathbb{R}^n iff $\{\Gamma(x_n)\}$ is Cauchy in $(V, \|\cdot\|_V)$.
3. As a result, $(V, \|\cdot\|_V)$ is complete by the Heine-Borel Theorem. Since V is arbitrary, we have that *all finite dimensional normed linear spaces are complete*.

Theorem 91 (The Basis of a Infinite Dimensional Banach Spaces is Uncountable)

Suppose $(W, \|\cdot\|)$ is a infinite dimensional Banach Space. If $\{w_\alpha\}_{\alpha \in I}$ is a basis of W , then I is uncountable.

Exercise 30.2.1

Prove  Theorem 91 (see also in A3).

Theorem 92 (All Linear Maps Between Finite Dimensional Normed Linear Spaces are Bounded)


If $(V, \|\cdot\|_V)$ and $(W, \|\cdot\|_W)$ are finite dimensional normed linear spaces, and $T : V \rightarrow W$ is linear, then T is bounded.


Proof

Consider the following diagram that illustrates the relationship between each of the spaces: Then, we define $S : (\mathbb{R}^n, \|\cdot\|_2) \rightarrow$

$$\begin{array}{ccc}
 (V, \|\cdot\|_V) & \xrightarrow{T} & (W, \|\cdot\|_W) \\
 \Gamma_n \downarrow \uparrow \Gamma_n^{-1} & & \Gamma_m \downarrow \uparrow \Gamma_m^{-1} \\
 (\mathbb{R}^n, \|\cdot\|_2) & & (\mathbb{R}^m, \|\cdot\|_2)
 \end{array}$$

Figure 30.1: Relationship between the finite dimensional normed linear spaces.

$(\mathbb{R}^m, \|\cdot\|_2)$ such that $S = \Gamma_m \circ T \circ \Gamma_n^{-1}$. By  Proposition 89, S is continuous. Consequently, we have that $T = \Gamma_m^{-1} \circ S \circ \Gamma_n$, which is a composition of continuous functions. Thus T is continuous, and hence bounded.

 **Corollary 93 (All Linear Maps from A Finite Dimensional Normed Linear Space to Any Normed Linear Space is Bounded)**

If $(V, \|\cdot\|_V)$ is a finite dimensional normed linear space, and $T : (V, \|\cdot\|_V) \rightarrow (W, \|\cdot\|_W)$ is linear, then T is bounded.

31.1 Finite Dimensional Normed Linear Space (Continued)

In the last lecture, we discovered that if $(V, \|\cdot\|_V)$ is an n -dimensional normed linear space, then

$$(V, \|\cdot\|_V) \simeq (\mathbb{R}^n, \|\cdot\|_2).$$

Notice that if $v \in V$, then $v = \Gamma_n(\Gamma_n^{-1}(v))$, and so

$$\|v\| = \|\Gamma_n(\Gamma_n^{-1}(v))\| \leq \|\Gamma_n\| \|\Gamma_n^{-1}(v)\|_2.$$

By applying Γ_n^{-1} once more, we have

$$\|\Gamma_n^{-1}(v)\|_2 \leq \|\Gamma_n^{-1}\| \|v\|_V.$$

It follows that if we let $\alpha = \frac{1}{\|\Gamma_n^{-1}\|}$ and $\beta = \|\Gamma_n\|$, then

$$\alpha \|\Gamma_n^{-1}(v)\|_2 \leq \|v\|_V \leq \beta \|\Gamma_n^{-1}(v)\|_2$$

for every $v \in V$.

We can deduce the following from the above:

1. A set $A \subset V$ is open/closed/compact in V iff $\Gamma_n^{-1}(A)$ is open/closed/compact in \mathbb{R}^n .
2. $A \subset (V, \|\cdot\|)$ is compact iff A is closed and bounded¹.
3. A sequence $\{v_n\}$ is Cauchy/converges to v_0 in $(V, \|\cdot\|_V)$ iff $\{\Gamma_n(v_n)\}$ is Cauchy/converges to $\Gamma_n(v_0)$ in $(\mathbb{R}^n, \|\cdot\|_2)$.

¹ This is also known as the **Heine-Borel Property**.

The following result follows from our observations above:

Theorem 94 (Completeness of Finite Dimensional Normed Linear Spaces)

Let $(V, \|\cdot\|_V)$ be a finite dimensional normed linear space. Then $(V, \|\cdot\|_V)$ is complete. In particular, if $(W, \|\cdot\|_W)$ is any normed linear space, and V is a finite dimensional subspace of W , then V is closed in W .


Example 31.1.1 (Unbounded Linear Function)

Let $(W, \|\cdot\|_W)$ be infinite dimensional, with basis $\{v_\alpha\}_{\alpha \in I}$. WMA that $\{v_\alpha\}_W = 1$. Choose a countable collection $\{v_1, v_2, \dots\} \subset \{v_\alpha\}_{\alpha \in I}$, and define

$$\varphi(v_\alpha) = \begin{cases} n & v_\alpha = v_n \\ 0 & \text{otherwise} \end{cases}$$

Then if $w = \alpha_1 v_1 + \dots + \alpha_n v_n$, we have

$$\varphi(w) = \sum_{i=1}^n \alpha_i \varphi(v_{\alpha_i}).$$

Then $\varphi : W \rightarrow \mathbb{R}$ is linear. 

QUESTION: Is φ bounded? No.

31.2 Uniform Continuity

We will finish on compactness with a few more results about uniform continuity.

Theorem 95 (Sequential Characterization of Uniform Continuity)

Let $f : (X, d_X) \rightarrow (Z, d_Z)$. TFAE:

- f is uniformly continuous.
- if $\{x_n\}, \{y_n\} \subset X$ with $d(x_n, y_n) \rightarrow 0$, then $d(f(x_n), f(y_n)) \rightarrow 0$.


 **Proof**

(1) \implies (2) f is uniformly continuous

$$\begin{aligned} &\implies \forall \varepsilon > 0 \exists \delta > 0 \forall x, y \in X \ d_X(x, y) < \delta \implies \\ &d_Z(f(x), f(y)) < \varepsilon \\ &\implies \exists N_0 \in \mathbb{N} \forall n \geq N_0 \ d_X(x_n, y_n) < \delta \implies d_Z(f(x_n), f(y_n)) < \\ &\varepsilon \dashv \end{aligned}$$

(2) \implies (1) f is not uniformly continuous

$$\begin{aligned} &\implies \exists \varepsilon_0 > 0 \forall \delta > 0 \exists x_0, y_0 \in X \\ &d_X(x_0, y_0) < \delta \wedge d_Z(f(x_0), f(y_0)) > \varepsilon_0 \\ &\implies \forall N \in \mathbb{N} \exists n_0 \geq N \\ &d_X(x_{n_0}, y_{n_0}) < \frac{1}{n} \wedge d_Z(f(x_{n_0}), f(y_{n_0})) > \varepsilon_0 \dashv \end{aligned}$$

 **Theorem 96 (Continuous Functions from a Compact Set Is Uniformly Continuous)**

If (X, d_X) is compact and if $f : (X, d_X) \rightarrow (Z, d_Z)$ is continuous, then f is uniformly continuous.


 **Proof**

Suppose to the contrary that f is not uniformly continuous


$$\begin{aligned} &\implies (\because \text{Theorem 95}) \forall \{x_n\}, \{y_n\} \subset X \\ &d_X(x_n, y_n) \rightarrow 0 \wedge d_Z(f(x_n), f(y_n)) \geq \varepsilon_0 > 0 \end{aligned}$$

But compactness $\implies \exists \{x_{n_k}\} \subset \{x_n\}, \{y_{n_k}\} \subset \{y_n\}$ such that

$$\begin{aligned} &x_{n_k} \rightarrow x_0 \in X \wedge y_{n_k} \rightarrow y_0 \in X \\ &\implies (\because \text{continuity}) f(x_{n_k}) \rightarrow f(x_0) \wedge f(y_{n_k}) \rightarrow f(y_0) \\ &\implies d_Z(f(x_{n_k}), f(y_{n_k})) \rightarrow 0 \not\geq \end{aligned}$$

 **Theorem 97 (Continuous Bijections from a Compact Space is a Homeomorphism)**

Assume (X, d_X) is compact and that $f : (X, d_X) \rightarrow (Y, d_Y)$ is continuous and bijective. Then $f^{-1} : Y \rightarrow X$ is continuous. In particular, f is a homeomorphism.

 **Proof**

Notice that $(f^{-1})^{-1} = f$. Thus it suffices to show that if $U \subset X$ is open, then $f(U)$ is open in Y . Also, note that $Y = f(X)$ is compact as X is compact.

$U \subset X$ is open $\implies F = U^C$ is closed
 $\implies F$ is compact ($\because f$ is continuous)
 $\implies f(F)$ is compact in Y
 $\implies f(F)$ is closed
 $\implies f(U) = (f(F))^C$ is open ($\because f$ is bijective)

31.3 The Space $(C(X), \|\cdot\|_\infty)$

31.3.1 Weierstrass Approximation Theorem


Example 31.3.1

Note that by **Taylor's Expansion**, we have that

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \dots$$

Consider the partial sum

$$S_k(x) = \sum_{n=0}^k \frac{x^n}{n!}.$$

Then we have that $S_k(x) \rightarrow e^x$ pointwise on \mathbb{R} . In fact, $S_k(x) \rightarrow e^x$ uniformly on $[-M, M]$. 

QUESTION: Given a function $f \in C[a, b]$, can f be uniformly approximated by polynomials?

Before going further, notice that if, e.g. we let

$$\varphi(x) = \frac{x-a}{b-a},$$

then $\varphi : [a, b] \rightarrow [0, 1]$ bijectively so. Also, φ is continuous. Its inverse,

$$\varphi^{-1}(x) = x(b-a) + a$$

is also continuous. We can then define $\Gamma : C[0, 1] \rightarrow C[a, b]$ by

$$\Gamma(f)(x) = f \circ \varphi^{-1}(x),$$

whose inverse is

$$\Gamma^{-1} : C[a, b] \rightarrow C[0, 1] \text{ given by } \Gamma^{-1}(f)(x) = f \circ \varphi(x).$$

Notice that Γ is an isometry: we have

$$\|\Gamma(f) - \Gamma(g)\|_{\infty} = \|f - g\|_{\infty}$$

for any $f, g \in C[0, 1]$. Moreover, $\Gamma(p)$ is a polynomial iff p is a polynomial.²

Thus every continuous function in $C[a, b]$ can be uniformly approximated by polynomials iff the same is true in $C[0, 1]$, i.e. we only need to consider continuous functions on $[0, 1]$ for approximations.

² Basically, this part shows us that we can use φ , which is also a continuous function, to scale the domain of f so as to shrink it down to only at $[0, 1]$ instead of $[a, b]$.

NEXT, observe that if $f \in C[0, 1]$, and if we can approximate

$$g(x) = f(x) - ([f(1) - f(0)]x + f(0)),$$

uniformly to within $\varepsilon > 0$ ³, i.e.

$$\|g - p\|_{\infty} < \varepsilon,$$

we may do the same for $f(x)$ with polynomials

$$\|g - p\|_{\infty} < \varepsilon \iff \|f - [p - q]\| < \varepsilon,$$

where $q(x) = f(1) - f(0)$. Notice that here, we have

$$g(0) = 0 = g(1).$$

³ Notice that if we rearrange the equation, we have

$$f(x) = g(x) + f(0) + x(f(1) - f(0))$$

which tells us that if we can approximate g by a polynomial, then we can do so for f cause the later term is also a polynomial.

32 Lecture 32 Nov 26th

32.1 The Space $(C(X), \|\cdot\|_\infty)$ (Continued)

32.1.1 Weierstrass Approximation Theorem (Continued)

Before proving Weierstrass' Approximation Theorem, we require the following lemma:

Lemma 98 (Lemma for Weierstrass Approximation)

Let $x \in [0, 1]$, then if $n \in \mathbb{N}$, we have

$$(1 - x^2)^n \geq 1 - nx^2.$$

Proof

Let $f(x) = (1 - x^2)^n - [1 - nx^2]$. Notice that $f(0) = 0$. Then

$$f'(x) = 2nx \left(1 - (1 - x^2)^{n-1} \right) \geq 0.$$

Thus f is increasing from $x = 0$. It follows that

$$(1 - x^2)^n \geq 1 - nx^2,$$

as required.

Theorem 99 (★★★ Weierstrass Approximation Theorem)




Figure 32.1: Graph of $(1 - x^2)^n$ for large n , where $x \in [0, 1]$.

If $f \in C[a, b]$, then for each $\varepsilon > 0$, there exists a polynomial $p(x)$ such that

$$\|f - p\|_\infty < \varepsilon.$$

Proof

By our discussion by the end of Section 31.3.1, we may assume that $[a, b] = [0, 1]$, and that $f(0) = 0 = f(1)$. Consequently, we may extend f to a uniformly continuous function on \mathbb{R} by defining $f(x) = 0$ if $x \in (-\infty, 0] \cup [1, \infty)$.

Now, let $Q_n(x) = c_n (1 - x^2)^n$, where c_n is chosen such that

$$\int_{-1}^1 Q_n(t) dt = 1.$$

Notice that

$$\begin{aligned} \int_{-1}^1 (1 - x^2)^n dx &= 2 \int_0^1 (1 - x^2)^n dx \geq 2 \int_0^{\frac{1}{\sqrt{n}}} (1 - nx^2) dx \\ &= \frac{4}{3\sqrt{n}} > \frac{1}{\sqrt{n}}, \end{aligned}$$

and so we have

$$c_n < \sqrt{n}.$$

For each n , define

$$\begin{aligned} p_n(x) &= \int_{-1}^1 f(x+t) Q_n(t) dt = \int_{-x}^{1-x} f(x+t) Q_n(t) dt \\ &= \int_0^1 f(u) Q_n(u-x) du. \end{aligned}$$

Notice that by Leibniz's Integral Rule, we have

$$\frac{d^{2n+1}}{dx^{2n+1}} p_n(x) = \int_0^1 f(u) \frac{\partial^{2n+1}}{\partial x^{2n+1}} Q_n(u-x) du = 0$$

by the construction of $Q_n(t)$. Thus p_n is a polynomial of degree at most $2n$.

Now, note that since $\int_{-1}^1 Q_n(t) dt = 1$, we have that

$$f(x) = \int_{-1}^1 f(x) Q_n(t) dt.$$

¹ How did we arrive at this new limit of $\frac{1}{\sqrt{n}}$? **There is** no deep meaning behind the choice of $\frac{1}{\sqrt{n}}$. It's simply because it works.

² Here, we can shrink the limits of integration, for anything below $-x$ or above $1-x$ are 0 as per our assumption that f is zero at $(-\infty, 0] \cup [1, \infty)$.

Also, in the first integral, we used $Q_n(t)$ to average over the transformation $f(x+t)$, and in the last integral, we see that we can "massage" the first integral into one where we have, instead, f as an **averaging function** over $Q_n(u-x)$.

Let $\varepsilon > 0$. By continuity of f , we may find $0 < \delta < 1$ such that

$$|x - y| < \delta \implies |f(x) - f(y)| < \frac{\varepsilon}{2}.$$

Then for $x \in [0, 1]$, we have

$$\begin{aligned} |p_n(x) - f(x)| &= \left| \int_{-1}^1 (f(x+t) - f(x)) Q_n(t) dt \right| \\ &\leq \int_{-1}^1 |f(x+t) - f(x)| Q_n(t) dt \\ &= \int_{-1}^{-\delta} |f(x+t) - f(x)| Q_n(t) dt \\ &\quad + \int_{-\delta}^{\delta} |f(x+t) - f(x)| Q_n(t) dt \\ &\quad + \int_{\delta}^1 |f(x+t) - f(x)| Q_n(t) dt \\ &\leq 2 \|f\|_{\infty} \sqrt{n} (1 - \delta^2)^n + \frac{\varepsilon}{2} + 2 \|f\|_{\infty} \sqrt{n} (1 - \delta^2)^n \\ &= 4 \|f\|_{\infty} \sqrt{n} (1 - \delta^2)^n + \frac{\varepsilon}{2}, \end{aligned} \tag{32.1}$$

where Equation (32.1) follows by $\int_{-1}^1 Q_n(t) dt = 1$ and $Q_n(t) \geq 0$ for $x \in [0, 1]$. Then since $0 < \delta < 1$, it follows that for sufficiently large N , we have

$$4 \|f\|_{\infty} \sqrt{N} (1 - \delta^2)^N \leq \frac{\varepsilon}{2},$$

as the $(1 - \delta^2)^N$ term will “decay” much faster than \sqrt{N} .




Figure 32.2: Dirac Sequence




Figure 32.3: One of the Dirac Functions with δ as an inflection point

Proposition 100 (Moments)

Assume that $f \in C[0, 1]$, that

$$\int_0^1 f(t) dt = 0,$$

and

$$\int_0^1 f(t)t^n dt = 0$$

for every $n \in \mathbb{N}$. Then $f(x) = 0$ for $x \in [0, 1]$.

 **Proof**

Since $f \in C[0, 1]$, by the Weierstrass Approximation Theorem, for $\varepsilon > 0$, let $p_n(x)$ be a polynomial such that $\|f - p_n\|_\infty < \varepsilon$. Then by the linearity of integration, and our assumption, we have


$$\int_0^1 f(t)p_n(t) dt = 0.$$


Consequently, we have

$$\int_0^1 f^2(t) dt = 0,$$

and thus $f(x) = 0$ at $[0, 1]$.

Exercise 32.1.1

Prove  Proposition 100.

 **Theorem 101 (Banach-Mazurkiewickz Theorem)**

Let

$$\text{ND}([0, 1]) = \{f \in C[0, 1] : f \text{ is nowhere differentiable}\}.$$

Then $\text{ND}([0, 1])$ is residual³ in $(C[0, 1], \|\cdot\|_\infty)$.

³ For quick reference, a set is residual if its complement is of first category.

 **Proof**

For each n , define

$$\mathcal{F}_n = \left\{ f \in C[0, 1] \mid \exists x_0 \in \left[0, 1 - \frac{1}{n}\right] \forall 0 < h < 1 - x_0 \quad |f(x_0 + h) - f(x_0)| \leq nh \right\}.$$

We notice that each of the \mathcal{F}_n 's is closed. **incomplete proof, require further work** □

Remark 32.1.1

There is nothing special about $[0, 1]$ in the above theorem. In particular, it


works for any closed interval $[a, b]$.



33 Lecture 33 Nov 28th

33.1 The Space $(C(X), \|\cdot\|_\infty)$ (Continued 2)

33.1.1 Weierstrass Approximation Theorem (Continued 2)

 **Corollary 102 (Separability of $(C[a, b], \|\cdot\|_\infty)$)**

$(C[a, b], \|\cdot\|_\infty)$ is separable.

 **Proof**

Let

$$P_n = \{a_0 + a_1x + \dots + a_nx^n : a_i \in \mathbb{R}\}$$

$$Q_n = \{r_0 + r_1x + \dots + r_nx^n : r_i \in \mathbb{Q}\}.$$

Then $\overline{Q_n} = P_n$. Then by the Weierstrass Approximation Theorem, $\bigcup_{n=1}^{\infty} P_n$ is dense, and so is the countable set $\bigcup_{n=1}^{\infty} Q_n$.

33.1.2 Stone-Weierstrass Theorem

QUESTION: Given a compact metric space (X, d) , and a subspace $\Phi \subset C(X)$, how can we tell that Φ is dense?

From here, we shall always assume that (X, d) is a compact metric

space.

Definition 78 (Point-Separating)

We say that $\Phi \subset C(X)$ is *point-separating* if¹

$$\forall x, y \in X (x \neq y \implies \exists f \in \Phi (f(x) \neq f(y))).$$

¹ Note that this definition does mean that every $f \in \Phi$ is injective, as the function may depend on either one or both x and y . Of course, if every $f \in \Phi$ is injective, then Φ is, trivially, point-separating.

Proposition 103 ($C(X)$ is Point-Separating)

$C(X)$ is point-separating.

Proof

Let $a, b \in X$ such that $a \neq b$. Then, define $f_a(x) = d(a, x)$. It is then clear that $f_a \in C(X)$. Since $a \neq b$, we have that $f_a(b) = d(a, b) > 0$.

Note 33.1.1

Suppose that $\Phi \subset C(X)$, and $x_1, x_2 \in X$ with $x_1 \neq x_2$, such that for any $f \in \Phi$, $f(x_1) = f(x_2)$. Then if $g \in \overline{\Phi}$, we must have $g(x_1) = g(x_2)$ ². This shows that if Φ is dense in $C(X)$, then it must separate points.

² For otherwise g would not be continuous.

33.1.2.1 Lattice Version

Definition 79 (Lattice)

A subspace $\Phi \subset C(X)$ is a *lattice* if $f \vee g, f \wedge g \in \Phi$ for each $f, g \in \Phi$, where³

$$(f \vee g)(x) = \max\{f(x), g(x)\}$$

$$(f \wedge g)(x) = \min\{f(x), g(x)\}.$$

³ In words, a lattice is a set of functions closed under **maxima** and **minima**.

“Note 33.1.2

1. Notice that

$$(f \vee g)(x) = \frac{(f(x) + g(x)) + |f(x) - g(x)|}{2} \in C(X)$$

for any $f, g \in C(X)$.

2. For minima, we have

$$(f \wedge g)(x) = -(-f \vee -g) = \frac{(f(x) + g(x)) - |f(x) - g(x)|}{2} \in C(X).$$

It follows that since both $f \vee g$ and $f \wedge g$ are in $C(X)$ that $C(X)$ is a lattice. Moreover, if $\Phi \subset C(X)$ is a linear subspace, then Φ is a lattice if $f \vee g \in \Phi$ for every $f, g \in \Phi$.

Example 33.1.1

A function $f \in C[a, b]$ is said to be a piecewise linear if there exists

$$P = \{a = t_0 < t_1 < \dots < t_n = b\},$$

i.e. a partition of $[a, b]$, such that

$$f \upharpoonright_{[t_{i-1}, t_i]}(x) = m_i x + b_i.$$

The function is **piecewise polynomial** if

$$f \upharpoonright_{[t_{i-1}, t_i]} = c_{0,i} + c_{1,i}x + \dots + c_{n,i}x^n,$$

where $c_{j,i} \in \mathbb{R}$. Let

$$\Phi_1 = \{f \in C[a, b] \mid f \text{ is piecewise linear} \}$$

and

$$\Phi_2 = \{f \in C[a, b] \mid f \text{ is piecewise polynomial} \}.$$

It is clear that both Φ_1 and Φ_2 are lattices.



Theorem 104 (Stone-Weierstrass Theorem — Lattice Version)

Let (X, d) be a compact metric space. Let Φ be a linear subspace of $C(X)$ such that

1. the constant function $1 \in \Phi$ ⁴;
2. Φ is point-separating; and
3. $f \vee g \in \Phi$ for any $f, g \in \Phi$ ⁵

Then $\overline{\Phi}$ is dense in $C(X)$.

⁴ It is okay that we simultaneously have $1 \in \Phi$ and Φ separating points, for all we need to know that Φ separate points is that for any $x, y \in X$ with $x \neq y$, **there exists** some $f \in \Phi$ such that $f(x) \neq f(y)$.

⁵ This implies that Φ is a lattice by note on page 195.

Proof

Note that given $\alpha, \beta \in \mathbb{R}$ with $a \neq b \in X$, since Φ is point-separating (2), we can find $\varphi \in \Phi$ such that $\varphi(a) \neq \varphi(b)$. Then, let

$$g(t) = \alpha \cdot 1(t) + (\beta - \alpha) \frac{\varphi(t) - \varphi(a)}{\varphi(b) - \varphi(a)},$$

where $1(t)$ is the constant function $1 \in \Phi$. We have that $g \in \Phi$ since it uses operations of which Φ is closed under. Notice that

$$g(a) = \alpha \text{ and } g(b) = \beta.$$

Let $f \in C(X)$ and $\varepsilon > 0$. Now for any pair $x, y \in X$, we can find $\varphi_{x,y} \in \Phi$ such that $\varphi_{x,y}(x) = f(x)$ and $\varphi_{x,y}(y) = f(y)$ ⁶. Let $x \in X$. Since $\varphi_{x,y}(y) - f(y) = 0$, and both $\varphi_{x,y}$ and f are continuous, we can find, for each $y \in X$, a $\delta_y > 0$ such that if $t \in B(y, \delta_y)$, then

$$-\varepsilon < \varphi_{x,y}(t) - f(t) < \varepsilon.$$

Now since (X, d) is compact, we can find a finite collection $\{y_1, \dots, y_n\} \subset X$ such that

$$X = \bigcup_{i=1}^n B(y_i, \delta_{y_i}),$$

and within each of the $B(y_i, \delta_{y_i})$, we have

$$-\varepsilon < \varphi_{x,y_i}(t) - f(t) < \varepsilon$$

I need to get a better picture of the motivation of the proof.

This is likely not a proof that one can come up in one sitting, especially when it is a theory that covers over 2 centuries of mathematical work. As it is, it is very difficult to understand how this proof came by, and many of the steps are purely constructive.

What else can we understand from $\varphi_{x,y}$?

⁶ I feel somewhat on edge not having the faintest idea how $\varphi_{x,y}$ works, except that it separates x and y .

for $t \in B(y_i, \delta_{y_i})$. Then, let

$$\varphi_x = \varphi_{x, y_1} \vee \dots \vee \varphi_{x, y_n}.$$

If $z \in X$, then $z \in B(y_{i_0}, \delta_{i_0})$ for some $i_0 \in \{1, \dots, n\}$, and so

$$f(z) - \varepsilon \leq \varphi_{x, y_{i_0}}(z) \leq \varphi_x(z).$$

On the other hand, since $\varphi_x(x) - f(x) = 0$, and both φ_x and f are continuous, for each $x \in X$, we can find a $\delta_x > 0$ such that if $t \in B(x, \delta_x)$, then

$$-\varepsilon < \varphi_x(t) - f(t) < \varepsilon. \quad (33.1)$$

As before, by the compactness of (X, d) , we can find $\{x_1, \dots, x_m\} \subset X$ such that

$$X = \bigcup_{i=1}^m B(x_i, \delta_{x_i}).$$

Then, using a similar argument as in the previous case, by Equation (33.1), we have that

$$\varphi_x(t) < f(t) + \varepsilon.$$

Thus, if $z \in B(x_{i_1}, \delta_{x_{i_1}})$ for some $i_1 \in \{1, \dots, m\}$, we have

$$\varphi(z) := \varphi_{x_1}(z) \wedge \dots \wedge \varphi_{x_m}(z) \leq \varphi_{x_{i_1}}(z) < f(z) + \varepsilon.$$

Consequently, for any $z \in X$, we have that

$$f(z) - \varepsilon < \varphi(z) < f(z) + \varepsilon.$$

This gives us that for any $W \subset C(X)$, since we can construct such a φ that is within ε -distance of f , $W \cap \overline{\Phi} \neq \emptyset$, thus implying that $\overline{\Phi}$ is dense in $C(X)$.

33.1.2.2 Subalgebra Version

Definition 80 (Subalgebra)

A subspace $\Phi \subset C(X)$ is a **subalgebra** if $f \cdot g(x) = f(x)g(x) \in \Phi$ for any $f, g \in \Phi$.

Example 33.1.2

Let

$$P_n = \{a_0 + a_1x + \dots + a_nx^n\}.$$

Then

$$P = \bigcup_{n=1}^{\infty} P_n$$

is a subalgebra of $C[a, b]$.⁷



⁷ See a quick work in notes on PMATH 347.

🌲 Lemma 105 (Closure of a Subalgebra is a Subalgebra)

If $\Phi \subset C(X)$ is a subalgebra, then so is $\overline{\Phi}$.

Proof

Suppose $f_n \rightarrow f$ and $g_n \rightarrow g$, where $\{f_n\}, \{g_n\} \subset \Phi$. Then, we have

$$\alpha f_n \rightarrow \alpha f$$

for $\alpha \in \mathbb{R}$, and

$$f_n + g_n \rightarrow f + g.$$

Note that $\{g_n\}$ is bounded if $g_n \rightarrow g$. Then,

$$\|f_n g_n - f g\|_\infty \leq \|g_n\|_\infty \|f_n - f\|_\infty + \|f\|_\infty \|g_n - g\|_\infty$$

would imply that $f_n g_n \rightarrow f g$, and so $f g \in \Phi$.

We are ready for the subalgebra version of Stone-Weierstrass, which we shall prove in the next lecture.


Theorem (Stone-Weierstrass Theorem — Subalgebra Version)

34 Lecture 34 Nov 30th

34.1 The Space $(C(X), \|\cdot\|_\infty)$ (Continued 3)

34.1.1 Stone-Weierstrass Theorem (Continued)

34.1.1.1 Subalgebra Version (Continued)

 **Theorem 106 (Stone-Weierstrass Theorem — Subalgebra Version)**

If $\Phi \subset C(X)$ is a linear subspace such that

1. $1 \in \Phi$;
2. Φ is point-separating; and
3. $f \cdot g \in \Phi$ for all $f, g \in \Phi$ (which implies Φ is a subalgebra).

Then $\overline{\Phi}$ is dense in $C(X)$.

 **Proof (★ ★ ★)**

By Lemma 105, we may assume that Φ is closed.

Let $f \in \Phi$ and $\varepsilon > 0$. Also, let $M = \|f\|_\infty$. From the Weierstrass Approximation Theorem, we may find some polynomial

$$p(x) = a_0 + a_1x + \dots + a_nx^n$$

such that for an $yt \in [-M, M]$, we get

$$\|t - p(t)\| < \varepsilon.$$

Now consider the composition

$$p \circ f = a_0 \cdot 1 + a_1 \cdot f + \dots + a_n \cdot f^n,$$

which is in Φ . Thus for $x \in X$, we have

$$\|f(x) - p \circ f(x)\| < \varepsilon.$$

This implies that

$$\|f - p \circ f\|_\infty < \varepsilon.$$

Thus by the closure of Φ , we have that $f \in \bar{\Phi} = \Phi$.

Now notice that for $f, g \in \Phi$, since

$$f \vee g = \frac{f + g + |f - g|}{2},$$

we have $f \vee g \in \Phi$. Thus by [Theorem 104](#), $\bar{\Phi}$ is dense in $C(X)$.




Figure 34.1: Visualization of the proof for [Theorem 106](#).

Example 34.1.1

Let

$$\mathcal{P} = \{a_0 + a_1x + \dots + a_nx^n : n \in \mathbb{N}, a_i \in \mathbb{R}\}.$$

Then by [Theorem 106](#), $\bar{\mathcal{P}} \subset C[a, b]$ is dense. 

Let

$$\Phi = \text{span}\{1, x^2, x^4, \dots\} \subseteq C[-1, 1].$$

It is clear that Φ is an algebra. However, it does not separate points, since

$$(-1)^2 = 1 = (1)^2.$$

So in this case Φ is not dense.

But what about $C[0, 1]$? Notice that x^2 separates points on $[0, 1]$, and all other conditions are still met. Thus Φ is dense in $C[0, 1]$.

QUESTION: Then what about

$$\Phi' = \text{span}\{x^2, x^4, \dots\}?$$

Is Φ' dense in $C[0, 1]$? No. ¹If $f \in \Phi'$, then $f(0) = 0$.

But what about the closure

$$\overline{\text{span}\{x^2, x^4, \dots\}} \subset C[0, 1]?$$

Consider the set

$$S := \{f \in C[0, 1] \mid f(0) = 0\},$$

which is a **closed ideal** in $C[0, 1]$. Then, in particular, we have that for any $g \in C[0, 1]$, we have that $gf, fg \in S$ for any $f \in S$. It can be shown² that

$$S = \overline{\text{span}\{x^2, x^4, \dots\}}.$$

Consequently, we see that if $f \in S$, we have that

$$f \in \overline{\text{span}\{1, x^2, x^4, \dots\}} = C[0, 1].$$

Example 34.1.2

Let $X = [0, 2\pi)$ and

$$A = \{\lambda \in \mathbb{C} \mid |\lambda| = 1\}.$$

Consider the function $\varphi : X \rightarrow A$ given by

$$\varphi(\theta) = e^{i\theta}.$$

It is clear that φ is bijective. We may then define $d : X \rightarrow \mathbb{R}$ by

$$d(\theta_1, \theta_2) = \text{shortest arclength between } e^{i\theta_1} \text{ and } e^{i\theta_2}.$$

Then we have

$$([0, 2\pi), d) \simeq A,$$

and the space $([0, 2\pi))$ is compact. Then in particular, we have

$$\{f \in C[0, 2\pi] \mid f(0) = f(2\pi)\} = C[0, 2\pi) \simeq C(A).$$

I should find out about this.

¹This was given as a reason but I don't know what exactly does it entail. That said, it is clear that Φ' separates points, and still a subalgebra, but 1 can we still create the constant function 1 in Φ using only the other generators?

how?



²I should probably work this out on my own.




Example 34.1.3

The set

$$\text{Trig}([0, 2\pi]) := \text{span}\{1, \cos(nx), \sin(mx) \mid n, m \in \mathbb{Z}\}$$

is a subalgebra of $C[0, 2\pi]$ that is point separating, and has 1 in it (and closed). By  Theorem 106, $\text{Trig}([0, 2\pi])$ is dense in $C[0, 2\pi]$. 

 Note 34.1.1

Consider the set

$$C(X, \mathbb{C}) = \{f : X \rightarrow \mathbb{C} \mid f \text{ continuous and bounded}\},$$


with norm

$$\|f\|_\infty = \sup\{|f(x)| \mid x \in X\}.$$

We say that $\Phi \subset C(X, \mathbb{C})$ is **self-adjoint** if

$$f \in \Phi \implies \bar{f} \in \Phi.$$

With this, we have the complex version of the Stone-Weierstrass Theorem.

 Theorem 107 (Stone-Weierstrass Theorem — Complex Version)

If (X, d) is compact and Φ is a linear subspace of $C(X, \mathbb{C})$ that is self-adjoint, with

1. $1 \in \Phi$;
2. Φ separates points; and
3. $f \cdot g \in \Phi$ for any $f, g \in \Phi$.

Then $\bar{\Phi}$ is dense in $C(X, \mathbb{C})$.

Example 34.1.4

Reusing our last example, now

$$\text{Trig}([0, 2\pi)) = \text{span}\{e^{in\theta} \mid n \in \mathbb{Z}\}$$

is dense in $C([0, 2\pi), \mathbb{C})$.



35 Lecture 35 Dec 03rd

35.1 The Space $(C(X), \|\cdot\|_\infty)$ (Continued 4)

35.1.1 Compactness in $C(X)$ and the Ascoli-Arzelà Theorem

QUESTION: If (X, d) is compact and $\mathcal{F} \subset C(X)$, when is \mathcal{F} compact?

We require the following notion:

Definition 81 (Equicontinuity)


Let (X, d) be a metric space with $\mathcal{F} \subset C_b(X)$. We say that \mathcal{F} is *(pointwise) equicontinuous at $x_0 \in X$* if

$$\forall \varepsilon > 0 \exists \delta_{x_0} > 0 \forall f \in \mathcal{F} \forall x \in X \\ d(x, x_0) < \delta_{x_0} \implies |f(x) - f(x_0)| < \varepsilon.$$

We say that \mathcal{F} is *equicontinuous* if it is (pointwise) equicontinuous at each $x_0 \in X$.

We say that \mathcal{F} is *uniformly equicontinuous* if

$$\forall \varepsilon > 0 \exists \delta > 0 \forall f \in \mathcal{F} \forall x, y \in X \\ d(x, y) < \delta \implies |f(x) - f(y)| < \varepsilon.$$

 Note 35.1.1


Notice that in the definition above, as compared to **regular continuity** we have

1. for **continuity**, δ may depend on ε , f and x_0 ;
2. for **uniform continuity**, δ may depend on ε and f ;
3. for **equicontinuity**, δ may depend on ε and x_0 ; while
4. for **uniform equicontinuity**, δ may solely depend on ε .

This was outlined on Wikipedia¹.

¹ So take it with a grain of salt?

Example 35.1.1

A finite collection $\{f_1, \dots, f_n\} \subset C_b(X)$ is equicontinuous. This is a clear result since we may check for each of the functions. 

Proposition 108 (Equicontinuity in a Compact Set is Uniform)

If (X, d) is compact and if $\mathcal{F} \subset C(X)$ is equicontinuous, then \mathcal{F} is uniformly equicontinuous.


Proof

Let $\varepsilon > 0$. Since \mathcal{F} is equicontinuous, for each $x_0 \in X$, we can find $\delta_{x_0} > 0$ if $x \in B(x_0, \delta_{x_0})$, then $|f(x) - f(x_0)| < \frac{\varepsilon}{2}$ for any $f \in \mathcal{F}$. Since (X, d) is compact, the cover $\{B(x_0, \delta_{x_0})\}_{x_0 \in X}$ has a Lebesgue Number $\delta_0 > 0$. Then, let $0 < \delta < \delta_0$. If for $w, z \in X$ we have $d(w, z) < \delta$, then $z \in B(w, \delta) \subset B(x'_0, \delta_{x'_0})$ for some $x'_0 \in X$. Then

$$|f(z) - f(w)| \leq |f(z) - f(x'_0)| + |f(x'_0) - f(w)| < \varepsilon.$$


Definition 82 (Pointwise Bounded Functions)

A family of functions $\mathcal{F} \subset C_b(X)$ is **pointwise bounded** if for each $x_0 \in X$, $\exists M_{x_0} > 0$ such that $|f(x_0)| < M_{x_0}$ for every $f \in \mathcal{F}$. We say that \mathcal{F} is **uniformly bounded** if $\exists M > 0$ such that $\|f\|_\infty \leq M$ for every $f \in \mathcal{F}$.

 **Proposition 109 (Pointwise Bounded Equicontinuous Functions in a Compact Set are Uniformly Bounded)**

Assume that (X, d) is compact and that $\mathcal{F} \subseteq C(X)$ is equicontinuous and pointwise bounded. Then \mathcal{F} is uniformly bounded.

 **Proof**

By  Proposition 108, \mathcal{F} is uniformly equicontinuous. So let $\varepsilon = 1$. Then $\exists \delta > 0$ such that for any $x, y \in X$, if $y \in B(x, \delta)$, then $|f(x) - f(y)| < 1$ for any $f \in \mathcal{F}$. By compactness of (X, d) there exists a finite subset $\{x_1, \dots, x_n\} \subset X$ such that


$$X = \bigcup_{i=1}^n B(x_i, \delta).$$

By assumption, we also know that for each of these x_i 's, there exists $M_1, \dots, M_n > 0$ such that for any $f \in \mathcal{F}$, $|f(x_i)| \leq M_i$. Then let

$$M_0 = \max\{M_1, \dots, M_n\}.$$

Then for any $z \in X$, we have that $z \in B(x_{i_0}, \delta)$ for some i_0 . Therefore, we have that

$$|f(z)| \leq |f(z) - f(x_{i_0})| + |f(x_{i_0})| < 1 + M_0.$$

 **Definition 83 (Relatively Compact Sets)**

Let $A \subset (X, d)$. We say that A is **relatively compact** if \bar{A} is compact.

Note 35.1.2

If (X, d) is complete, then we have that A is relatively compact iff A is totally bounded.

Theorem 110 (Arzelà-Ascoli)

Let (X, d) be a compact metric space, and $\mathcal{F} \subset C(X)$. TFAE:

1. \mathcal{F} is relatively compact.
2. \mathcal{F} is equicontinuous and pointwise-bounded.

Proof

(1) \implies (2) Since (X, d) is compact, it is complete, and so \mathcal{F} being relatively compact implies that \mathcal{F} is totally bounded. Thus \mathcal{F} has a finite $\frac{\varepsilon}{3}$ -net $\{f_1, f_2, \dots, f_n\} \subset \mathcal{F}$. By an earlier example, we have that $\{f_1, f_2, \dots, f_n\}$ is equicontinuous, and hence uniformly equicontinuous by [Proposition 108](#). By that, we can find a $\delta > 0$ such that $\forall x, y \in X$, if $d(x, y) < \delta$, we have

$$|f_i(x) - f_i(y)| < \frac{\varepsilon}{3}$$

for all $i = 1, 2, \dots, n$.

Now let $f \in \mathcal{F}$ be arbitrary, and let $w, z \in X$ such that $d(w, z) < \delta$. Since \mathcal{F} has a finite $\frac{\varepsilon}{3}$ -net, there exists $i_0 = 1, 2, \dots, n$ such that $\|f - f_{i_0}\|_\infty < \frac{\varepsilon}{3}$. Thus

$$\begin{aligned} |f(w) - f(z)| &\leq |f(w) - f_{i_0}(w)| + |f_{i_0}(w) - f_{i_0}(z)| + |f_{i_0}(z) - f(z)| \\ &\leq \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} < \varepsilon. \end{aligned}$$

Therefore, \mathcal{F} is uniformly continuous and uniformly bounded².

² We proved for the stronger version.

(2) \implies (1) By [Proposition 108](#) and [Proposition 109](#), we have that \mathcal{F} is uniformly continuous and uniformly bounded. Let

$\varepsilon > 0$. By uniform boundedness, let $M > 0$ be such that $|f(x)| < M$ for every $x \in X$ and every $f \in \mathcal{F}$. Consider the partition

$$P = \{-M = y_0 < y_1 < y_2 < \dots < y_m = M\},$$

where $y_j - y_{j-1} < \frac{\varepsilon}{3}$ for each $j = 1, \dots, m$.

We may also find, by uniform equicontinuity, a $\delta > 0$ such that $d(w, z) < \delta$ implies that $|f(z) - f(w)| < \frac{\varepsilon}{3}$. Since (X, d) is totally bounded (as it is compact), we may find, in particular, a finite δ -net $\{x_1, \dots, x_n\} \subset X$ such that

$$X = \bigcup_{i=1}^n B(x_i, \delta).$$

Now consider the set functions

$$\Phi = \{\varphi \mid \varphi : \{1, \dots, n\} \rightarrow \{1, \dots, m\}\}.$$

It is clear that Φ is finite, and so we may write

$$\Phi = \{\varphi_1, \dots, \varphi_l\},$$

where $l = m^n$.

Next, for each $k = 1, \dots, l$, let

$$\mathcal{F}_k = \left\{ f \in \mathcal{F} \mid f(x_i) \in [y_{\varphi_k(i)-1}, y_{\varphi_k(i)}] \right\}.$$

Clearly so, by construction, while some of the \mathcal{F}_k 's may be empty, we have that $\{\mathcal{F}_k\}$ partitions \mathcal{F} , i.e.

$$\mathcal{F} = \bigcup_{k=1}^l \mathcal{F}_k.$$

Then for each of the non-empty sets \mathcal{F}_k , pick a $f_k \in \mathcal{F}_k$. From here, since we want to show that \mathcal{F} is relatively compact and (X, d) is compact and hence complete itself, it suffices for us to show that \mathcal{F} is totally bounded. In other words, it suffices for us to show that \mathcal{F} has some finite ε -net.

Let $f \in \mathcal{F}$. Then $f \in \mathcal{F}_k$ for some $k = 1, 2, \dots, l$. Then for




Figure 35.1: Basic Visual Sketch of the Proof of the Arzelà-Ascoli Theorem

A Useful Theorems from Earlier Calculus

Theorem A.1 (Monotone Convergence Theorem)

Let $\{x_k\}$ be a sequence in \mathbb{R} .

1. Suppose $\{x_k\}$ is increasing.
 - If $\{x_k\}$ is bounded above, then $x_k \rightarrow \sup\{x_k\}$ as $k \rightarrow \infty$.
 - If $\{x_k\}$ is not bounded above, then $x_k \rightarrow \infty$ as $k \rightarrow \infty$.
 2. Suppose $\{x_k\}$ is decreasing.
 - If $\{x_k\}$ is bounded below, then $x_k \rightarrow \inf\{x_k\}$ as $k \rightarrow \infty$.
 - If $\{x_k\}$ is not bounded below, then $x_k \rightarrow -\infty$.
-

B Assignment 1

1. *

- (a) How many relations are there on the set $\{1, 2, 3, \dots, n\}$?
(i) n (ii) n^2 (iii) 2^n (iv) 2^{n^2}
- (b) Determine the number of equivalence relations on the set $X = \{1, 2, 3\}$.
(i) 4 (ii) 5 (iii) 6 (iv) None of the above
- (c) Recall that we would say that $A \sim B$ and that A and B have the same *cardinality*, if there is a 1 – 1 and onto function from A to B .

If $X = \{1, 2, 3, 4\}$ and \sim is the equivalence relation on $\mathcal{P}(X)$ as above:

- i. How many different equivalence classes are there in this equivalence relation:

A. 4 B. 2^4 C. 5 D. 2^5

- ii. List all of the elements of $[A]$ if $A = \{1, 2, 3\}$.

- iii. If $X = \{1, 2, 3, \dots, n\}$ and \sim is as in Part 1c, how many elements are there in $[A]$ where $A = \{1, 2, 3, \dots, k\}$?

A. 2^k B. $k!$ C. $\frac{n!}{k!}$ D. $\frac{n!}{k!(n-k)!}$

- 2.(a) Let V be a vector space. Let W be a subspace of V . Show that:

$$v \sim y \iff v - y \in W,$$

defines an equivalence relation on V .

- (b) Show that $[z] + [v] = [z + v]$ and $\alpha[z] = [\alpha z]$ is well defined. That

is, show that if $z_1 \sim z_2$ and $v_1 \sim v_2$, then $z_1 + v_1 \sim z_2 + v_2$ and $\alpha z_1 \sim \alpha z_2$.

Remark The set $V/W = [v] \mid v \in V$ is a vector space under the operations above. It is called the *quotient* of V by W .

3. *

- (a) Use cardinal arithmetic to determine $(\aleph_0)^{\aleph_0}$ and c^{\aleph_0} and c^{\aleph_0} .
- (b) Show that there exists a 1 – 1 map from the power set of \mathbb{R} onto the set of all real-valued functions on \mathbb{R} by showing that $2^c = c^c$.
- (c) Explain why there is a one to one and onto map $\Gamma : \mathbb{Q}^\infty \rightarrow \mathbb{R}^\infty$ where

$$\mathbb{Q}^\infty = \{\{r_n\} \mid r_n \in \mathbb{Q}\}$$

and

$$\mathbb{R}^\infty = \{\{s_n\} \mid s_n \in \mathbb{R}\}.$$

- (d) Let $C(\mathbb{R})$ denote the set of all continuous real-valued functions on \mathbb{R} .
- i. Explain why if $f, g \in C(\mathbb{R})$ and $f(x) = g(x)$ for every $x \in \mathbb{Q}$, then $f = g$.
- ii. Determine $|C(\mathbb{R})|$.

4. A real number $\alpha \in \mathbb{R}$ is called algebraic if there exists a polynomial $p(x)$ with integer coefficients such that $p(\alpha) = 0$. Show that the collection Ψ of all algebraic numbers is countable.

5. A collection $\mathfrak{S} \subseteq \mathcal{P}(X)$ is called a topology on X if

- (a) $\emptyset, X \in \mathfrak{S}$
- (b) $\left\{ \bigcup_{\alpha \in I} U_\alpha \right\} \in \mathfrak{S}$ whenever $\{U_\alpha\}_{\alpha \in I} \subseteq \mathfrak{S}$
- (c) $\bigcap_{i=1}^n U_i \in \mathfrak{S}$ whenever $\{U_1, U_2, \dots, U_n\} \subseteq \mathfrak{S}$

The elements of \mathfrak{S} are called \mathfrak{S} -open sets or simply open sets for short.

- (a) Show that if $\{\mathfrak{S}_\alpha\}_{\alpha \in I}$ is a collection of topologies on X , then

$\mathfrak{S} = \bigcap_{\alpha \in I} \mathfrak{S}_\alpha$ is also a topology on X . In particular, show that if $\Gamma \subseteq \mathcal{P}(X)$, then there is a smallest topology $\mathfrak{S}(\Gamma)$ on X that contains Γ . $\mathfrak{S}(\Gamma)$ is called *the topology generated by Γ* .

(b) * We call a subset U of \mathbb{R} *open* if for every $x \in U$, there exists an $\varepsilon > 0$ such that $(x - \varepsilon, x + \varepsilon) \subseteq U$. Let $\mathfrak{S}_{\mathbb{R}}$ denote the collection of all open subsets of \mathbb{R} .

i. Show that $\mathfrak{S}_{\mathbb{R}}$ is a topology on \mathbb{R} .

ii. Let

$$\Gamma = \{\emptyset\} \cup \{(a, b) \mid a \in \mathbb{R} \cup \{-\infty\}, b \in \mathbb{R} \cup \{\infty\}, a < b\}$$

be the collection of open intervals in \mathbb{R} . Show that $\mathfrak{S}_{\mathbb{R}} = \mathfrak{S}(\Gamma)$.

iii. Let $U \subset \mathbb{R}$ be open and nonempty. Define a relation \sim on U by $x \sim y$ if and only if whenever $x < z < y$ or $y < z < x$, we must have $z \in U$.

Show that \sim is an equivalence relation on U and that if $I_x = \{y \in U \mid x \sim y\}$, then I_x is an open interval. (Recall that a set I is an interval if whenever $x, y \in I$ and $x < z < y$, then we must have $z \in I$.)

Remark: In this case, in fact, $I_x = (\alpha_x, \beta_x)$, where

$$\alpha_x = \inf\{y : (x, y) \subset U\}$$

$$\beta_x = \sup\{y : (y, x) \subset U\}$$

iv. Show that if $U \in \mathfrak{S}_{\mathbb{R}}$, then U is the union of at most countably many pairwise disjoint open intervals.

v. What is $|\mathfrak{S}_{\mathbb{R}}|$? (Hint: Show that every open set is the countable union of open intervals with rational endpoints.)

(c) * Let X be any set. Let $\mathfrak{S}_{cf}(X) = \{\emptyset\} \cup \{A \subseteq X \mid A^c \text{ is finite}\}$. Show that $\mathfrak{S}_{cf}(X)$ is a topology on X . $\mathfrak{S}_{cf}(X)$ is called the *cofinite topology* on X .

(d) Let X be any set. Let $\mathfrak{S}_{cc}(X) = \{\emptyset\} \cup \{A \subseteq X \mid A^c \text{ is countable}\}$. Show that $\mathfrak{S}_{cc}(X)$ is a topology on X . $\mathfrak{S}_{cc}(X)$ is called the *co-*

countable topology on X .

6. Let X be a given set. A σ -algebra on X is a collection Ψ of subsets of X such that

- (i) $X \in \Psi$;
- (ii) If $S \in \Psi$, then so is S^c .
- (iii) If $\{S_n\} \subset \Psi$, then $\bigcup_{n=1}^{\infty} S_n \in \Psi$.

(a) Show that if $\{\Psi_\alpha\}_{\alpha \in I}$ is any collection of σ -algebras on X , then $\bigcap_{\alpha \in I} \Psi_\alpha$ is also a σ -algebra. In particular, show that if $\mathcal{A} \subseteq \mathcal{P}(X)$, then there is a unique smallest σ -algebra containing \mathcal{A} which we call the σ -algebra generated by \mathcal{A} , and denote by $\sigma(\mathcal{A})$.

(b) Let \mathcal{O} denote the collection of all open subsets of \mathbb{R} . The σ -algebra, $\sigma(\mathcal{O})$ is called the Borel σ -algebra of \mathbb{R} , and is denoted by $\mathcal{B}(\mathbb{R})$.

Give an example of a set $A \subset \mathbb{R}$ that is Borel but neither closed or open.

(c) What is $|\mathcal{B}(\mathbb{R})|$? (**Note:** This one is not so easy. Do not spend much time on it and only do so after you have completed the remaining questions)

(d) True or false: Every uncountable subset S of \mathbb{R} contains a subset A which is not Borel. (Explain your answer.)

7. *

(a) Show that if X is infinite and countable, you can find two disjoint infinite subsets S and T such that $S \cup T = X$ and

$$|S| = |T| = |X|.$$

(b) Show that if X is infinite, then you can find two disjoint subsets S and T such that $S \cup T = X$ and $|S| = |T| = |X|$. (**Hint:** Show that X can be written as the union of a collection of pairwise disjoint countable sets.)

Remark: This is actually a formal proof of the statement for an infinite set $|X| + |X| = |X|$.

C Assignment 2

1. *We have seen that the positive rationals can be well ordered via the order \preceq given by $\frac{n}{m} \preceq \frac{j}{k}$ if and only if $2^n 3^m \leq 2^j 3^k$. With respect to this order find the least element in the set $S = \{r \in \mathbb{Q} \mid \sqrt{2} < r\}$. (Note: In defining S the order we use is the usual order on \mathbb{R} .)

↓ for markings and comments

2. *Let d_1, d_2 and d_∞ be the metrics on \mathbb{R}^n given by

$$d_1(x, y) = \sum_{i=1}^n |x_i - y_i|$$
$$d_2(x, y) = \sqrt{\sum_{i=1}^n (x_i - y_i)^2}$$
$$d_\infty(x, y) = \max_{i=1, \dots, n} |x_i - y_i|$$

Let τ_1, τ_2 and τ_∞ be the topologies induced by the above metrics.

Show that $\tau_1 = \tau_2 = \tau_\infty$.

3. *

- (a) For each of the following sets determine if it is open, closed or neither. Indicate the set of limit points, boundary points and interior points of each set.

i. $(0, 1] \subset \mathbb{R}$.

ii. $\mathbb{Q} \subset \mathbb{R}$.

- (b) Let $\mathcal{P}_1 = \{a_0 + a_1 x \mid a_i \in \mathbb{R}\} \subset (C[0, 1], d_\infty)$. Show that \mathcal{P}_1 is closed.

- (c) Let $c_{00} = \{\{a_n\} \in l_\infty \mid a_n = 0 \text{ for all but finitely many } n\} \subset l_\infty$.
Let $c_0 = \{\{a_n\} \in l_\infty \mid \lim_{n \rightarrow \infty} a_n = 0\}$. Show that c_{00} is dense in

c_0 . That is $\overline{c_{00}} = c_0$.

4. Least Upper Bound Property:

We say that α is an upper bound of $S \subset \mathbb{R}$ if $x \leq \alpha$ for all $x \in S$.

We say that S is bounded above if it has an upper bound. We call α the *least upper bound* of S if α is an upper bound of S and if whenever β is an upper bound of S we have $\alpha \leq \beta$. We denote α by $\text{lub}(S)$ (We may define lower bounds and the *greatest lower bound* ($\text{glb}(S)$) in the obvious way). The *Least Upper Bound Property* states that every nonempty subset S of \mathbb{R} that is bounded above has a least upper bound (or equivalently that every nonempty subset S of \mathbb{R} that is bounded below has a greatest lower bound).

- (a) Prove the Monotone Convergence Theorem: Let $\{a_n\}$ be a sequence in \mathbb{R} with $a_n \leq a_{n+1}$ for all $n \in \mathbb{N}$. If $\{a_n\}$ is bounded above, then $\{a_n\}$ converges.
- (b) Prove the Nest Interval Theorem: Let $\{[a_n, b_n]\}$ be sequence of closed intervals with $[a_{n+1}, b_{n+1}] \subseteq [a_n, b_n]$ for each $n \in \mathbb{N}$. Then $\bigcap_{n=1}^{\infty} [a_n, b_n] \neq \emptyset$.
- (c) Show that the statement in Part 4b may fail if we use open intervals.
- (d) Use the Nest Interval Theorem to show that if $S \subset \mathbb{R}$ is infinite and bounded, then it has a limit point. (This is called the Bolzano-Weierstrass Theorem.)
- (e) Given a nonempty set $A \subset (X, d)$ we define the diameter of A to be $\text{diam}(A) = \sup\{d(x, y) \mid x, y \in A\}$. Show that if A_n is a sequence of nonempty closed sets in \mathbb{R} with $A_{n+1} \subseteq A_n$ and $\text{diam}(A_i) < \infty$, then $\bigcap_{n=1}^{\infty} A_n \neq \emptyset$.

5. *Let $\{U_\alpha\}_{\alpha \in I}$ be a collection of open sets in \mathbb{R} such that $[0, 1] \subset \bigcup_{\alpha \in I} U_\alpha$.

- (a) Show that there exists finitely many sets $U_{\alpha_1}, U_{\alpha_2}, \dots, U_{\alpha_n}$ such that $[0, 1] \subset \bigcup_{i=1}^n U_{\alpha_i}$.

(Hint: Let

$$A = \{x \in [0, 1] \mid [0, x] \text{ can be covered by finitely many } U_\alpha\text{'s}\}.$$

Show that $1 = \text{lub}(A)$ and then that $1 \in A$.)

- (b) Show that the statement in Part 5a can fail if we replace $[0, 1]$ with $(0, 1)$.
6. *A map $\varphi : (X, d_X) \rightarrow (Y, d_Y)$ is called an isometry if $d_Y(\varphi(x_1), \varphi(x_2)) = d_X(x_1, x_2)$.
- (a) Determine all possible isometries $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ and $\psi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and show that each such map is surjective.
- (b) Given an example of an isometry $\varphi : (X, d_X) \rightarrow (X, d_X)$ that is not onto.
7. *A topological space (X, τ) is called separable if there exists a countable subset $S \subset X$ such that $\bar{S} = X$. Show that (ℓ_1, d_1) is separable but (ℓ_∞, d_∞) is not.
8. *Let $\vec{x}_n = \{x_{n,1}, x_{n,2}, x_{n,3}, \dots\} \in l_\infty$. Show that if $\vec{x}_n \rightarrow \vec{x}_0$ in l_∞ where $\vec{x}_0 = \{x_{0,1}, x_{0,2}, x_{0,3}, \dots\}$, then for each $k \in \mathbb{N}$, $\lim_{n \rightarrow \infty} x_{n,k} = x_{0,k}$ but that the converse can fail.
9. Let $P_0 = [0, 1]$. Let P_1 be obtained from P_0 by removing the open interval of length $\frac{1}{3}$ from the middle of P_0 . Then construct P_2 from P_1 by removing open intervals of length $\frac{1}{3^2}$ from the two closed subintervals of P_1 . In general, P_{n+1} is obtained from P_n by removing the open interval of length $\frac{1}{3^{n+1}}$ from the middle of each of the 2^n closed subintervals of P_n . Let

$$P = \bigcap_{n=0}^{\infty} P_n.$$

P is called the Cantor set.

- (a) A subset A of a metric space is **nowhere dense** if $\bar{A}^\circ = \emptyset$. Show that P is closed and nowhere dense.
- (b) Show that P is uncountable. (Hint: You may use the fact that $x \in P$ if and only if we can express $x = \sum_{n=1}^{\infty} \frac{a_n}{3^n}$ where $a_n = 0, 2$.)
- (c) A subset A of \mathbb{R} is said to be *perfect* if $A = \text{Lim}(A)$. Show that the Cantor set P is perfect. (Again, you can use the fact that $x \in P$ if and only if we can express $x = \sum_{n=1}^{\infty} \frac{a_n}{3^n}$ where $a_n = 0, 2$.)

D Assignment 3

- 1.(a) Let (X, d) be a metric space. Let $x_0 \in X$ be fixed. Define $F_{x_0} : X \rightarrow \mathbb{R}$ by

$$F_{x_0}(x) = d(x_0, x).$$

Show that F_{x_0} is continuous.

- (b) * Let $(X, \|\cdot\|)$ be a normed linear space. Define $F : X \rightarrow \mathbb{R}$ by

$$F(x) = \|x\|.$$

Show that F is continuous.

2. * Let $f_n : [0, 1] \rightarrow \mathbb{R}$ be defined by

$$f_n(x) = \sin(x^n).$$

- (a) Show that $f_n(x)$ does not converge uniformly on $[0, 1]$.
(b) Show that $f_n(x)$ does converge uniformly on $\left[0, \frac{1}{2}\right]$.

3. Connectedness of \mathbb{R}

Let $A \subseteq (X, d)$. We say that A is *disconnected* if there exists two open sets U and V such that

- i) $U \cap V \cap A = \emptyset$
- ii) $U \cap A \neq \emptyset$ and $V \cap A \neq \emptyset$
- iii) $A \subseteq U \cup V$.

We say that A is *connected* if it is not disconnected.

- (a) Let (X, d) be a metric space and let $A \subset X$. Show that the

characteristic function

$$\chi_A(x) := \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$

is continuous on X if and only if A is both open and closed.

(b) * Show that \mathbb{R} is connected.

(c) Let $A \subseteq (X, d_X)$ be connected. Let $f : A \rightarrow (Y, d_Y)$ be continuous. Show that $f(A)$ is connected.

4. A function $f : (X, d_X) \rightarrow (Y, d_Y)$ is said to be uniformly continuous if for every $\varepsilon > 0$ there exists a $\delta > 0$ such that if $d_X(x_1, x_2) < \delta$, then $d_Y(f(x_1), f(x_2)) < \varepsilon$.

(a) Let $f : (X, d_X) \rightarrow (Y, d_Y)$ be uniformly continuous. Show that if $\{x_n\}$ is Cauchy in X , then $\{f(x_n)\}$ is Cauchy in Y .

(b) Let (X, d) be a metric space and let $A \subset X$. Let $f : A \rightarrow \mathbb{R}$. Show that if f is uniformly continuous on A , then there exists $g : \bar{A} \rightarrow \mathbb{R}$ that is continuous on \bar{A} and for which $g \upharpoonright_A = f$. That is g extends f to \bar{A} .

5. Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be normed linear spaces. Let $T : X \rightarrow Y$ be linear. We say that T is bounded if

$$\sup_{\|x\|_X \leq 1} \{\|T(x)\|_Y\} < \infty.$$

In this case, we write

$$\|T\| = \sup_{\|x\|_X \leq 1} \{\|T(x)\|_Y\}.$$

Otherwise, we say that T is unbounded.

(a) * Prove that the following are equivalent

i. T is continuous.

ii. T is continuous at 0.

iii. T is bounded.

(b) Assume that $L : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is linear and that L is represented by

the matrix A . We let $\|A\| = \|L\|$.

i. Assume that

$$D = \begin{bmatrix} d_1 & & & & \\ & d_2 & & & \\ & & d_3 & & \\ & & & \ddots & \\ & & & & d_n \end{bmatrix}$$

is a diagonal matrix. Show that $\|D\| = \max_{i=1,\dots,n} \{|d_i|\}$.

ii. Show that if

$$D = \begin{bmatrix} d_1 & & & & \\ & d_2 & & & \\ & & d_3 & & \\ & & & \ddots & \\ & & & & d_n \end{bmatrix}$$

is a diagonal matrix, then

$$\sup_{\|x\| \leq 1} \{|\langle Dx, x \rangle|\} = \max_{i=1,\dots,n} \{|d_i|\}.$$

iii. Let U be an orthonormal $n \times n$ matrix. Show that if $x \in \mathbb{R}^n$, then $\|Ux\| = \|x\|$.

iv. * Assume that $L : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is linear and that L is represented by the matrix A . Show that $\|L\| = \|A\| = \sqrt{|\alpha|}$ where α is the largest eigenvalue of the matrix $A^t A$.

v. * Assume that $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is represented by the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 2 & -1 \end{bmatrix}.$$

Find $\|A\|$. (You can use Maple or MATLAB if you like.)

6. * Let $x_0 \in [0, 1]$. Define the linear map $T_{x_0} : C[0, 1] \rightarrow \mathbb{R}$ by

$$T_{x_0}(f) = f(x_0).$$

(a) Show that as a map from $(C[0, 1], \|\cdot\|_\infty) \rightarrow \mathbb{R}$, T_{x_0} is bounded

with $\|T_{x_0}\| = 1$.

(b) Show that as a map from $(C[0,1], \|\cdot\|_1) \rightarrow \mathbb{R}$, T_0 is unbounded.

7. Define the linear map $T : C[0,1] \rightarrow \mathbb{R}$ by

$$T(f) = \int_0^1 xf(x) dx.$$

(a) Show that if $\|f(x)\|_\infty \leq 1$, then $|T(f)| \leq \frac{1}{2}$.

(b) Show that if $T(1) = \frac{1}{2}$ and hence that $\|T\| = \frac{1}{2}$.

8. * Let (X, d) be a metric space and $\{f_n\}$ be a sequence of real valued functions on X which converges pointwise on X to a function $f : X \rightarrow \mathbb{R}$. Let $x_0 \in X$.

We say that $\{f_n\}$ converges *uniformly* at x_0 if for every $\varepsilon > 0$, there exists a $\delta > 0$ and an $N \in \mathbb{N}$ such that if $n > N$ and $d(x, x_0) < \delta$, then

$$|f_n(x) - f(x)| < \varepsilon.$$

Show that if each function f_n is continuous at x_0 and if $f_n \rightarrow f$ uniformly at x_0 then f is also continuous at x_0 . (Hint: This is almost exactly the same as the proof for uniform convergence with one minor change.)

9. Let (X, d) be a metric space. Let $f : X \rightarrow \mathbb{R}$. Let

$$D(f) = \{x_0 \in X \mid f(x) \text{ is discontinuous at } x_0\}.$$

For each $n \in \mathbb{N}$, let

$$D_n(f) = \left\{ x_0 \in X \mid \forall \delta > 0, \exists y, z \in B(x_0, \delta) \text{ for which } |f(y) - f(z)| \geq \frac{1}{n} \right\}.$$

(a) * Show that for each $n \in \mathbb{N}$, $D_n(f)$ is closed. (Hint: Let $\{x_k\} \subseteq D_n(f)$ be such that $x_k \rightarrow x_0$. Show that $x_0 \in D_n(f)$.)

(b) * A subset A of a metric space is said to be an F_σ set if $A = \bigcup_{n=1}^{\infty} F_n$, where each F_n is closed. Show that $D(f)$ is an F_σ set by showing that

$$D(f) = \bigcup_{n=1}^{\infty} D_n(f).$$

- (c) * A subset A of (X, d) is said to be *nowhere dense* if $\bar{A}^\circ = \emptyset$.
 Assume that $F \subset \mathbb{R}$ is closed and nowhere dense. Let

$$f(x) = \chi_F(x) = \begin{cases} 1 & \text{if } x \in F \\ 0 & \text{if } x \in F^c \end{cases}.$$

Find $D(f)$.

- (d) * A subset A of (X, d) is said to be *first category* if $A = \bigcup_{n=1}^{\infty} A_n$ where each A_n is nowhere dense. Show that if $A \subset \mathbb{R}$ is F_σ and of first category, then there exists a function $f(x)$ on \mathbb{R} with $D(f) = A$.
- (e) **Bonus Question 5:** Show that if $A \subset \mathbb{R}$ is F_σ then there exists a function $f(x)$ on \mathbb{R} with $D(f) = A$.
- 10.(a) * Explain why the integral equation

$$f(x) = x + \int_0^x t f(t) dt$$

has a unique solution $\varphi(x)$ in $C[0, 1]$, and then find a power series representation for $\varphi(x)$.

- (b) **Fredholm Equation:** Assume that $K(x, y) \in C([a, b] \times [a, b])$ with $\|K(x, y)\|_\infty = M$. Show that if $|\lambda| M(b - a) < 1$ and if $\varphi(x) \in C[a, b]$, then the map $\Gamma : C[a, b] \rightarrow C[a, b]$ given by

$$\Gamma(f)(x) = \varphi(x) + \lambda \int_a^b K(x, y) f(y) dy$$

is contractive and hence that the integral equation

$$f(x) = \varphi(x) + \lambda \int_a^b K(x, y) f(y) dy \quad (*)$$

has a unique solution in $C[a, b]$.

11. * **Dini's Theorem:** Let (X, d) be a compact metric space. Let $\{f_n(x)\}$ be a sequence of continuous functions on X such that $f_n(x) \leq f_{n+1}(x)$ for each $n \in \mathbb{N}$ and $f(x) = \lim_{n \rightarrow \infty} f_n(x)$.
- (a) Show that $f(x)$ is continuous on X if and only if the sequence converges uniformly. (Hint: Let $\varepsilon > 0$. Let $U_n = \{x \in X \mid f_n(x) > f(x) - \varepsilon\}$ and show that $\{U_n\}$ is an open cover of X .)

- (b) Show that Dini's Theorem fails on $[0, \infty)$ by giving a sequence $\{f_n(x)\}$ of continuous functions on $[0, \infty)$ such that $f_n(x) \leq f_{n+1}(x)$ for each $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} f_n(x) = 1$ for each x but for which the convergence is not uniform.

12. Let $A \subset (X, d)$ be non-empty. For each $x \in X$, define the distance from x to A by

$$\text{dist}(x, A) = \inf\{d(x, y) \mid y \in A\}.$$

- (a) Show that A is closed if and only if the following property holds:

$$x \in A \iff \text{dist}(x, A) = 0.$$

- (b) Let $F \subseteq X$ be closed and non-empty. Show that

$$F = \bigcap_{n \in \mathbb{N}} \left(\bigcup_{x \in F} B\left(x, \frac{1}{n}\right) \right).$$

(Note: This shows that every closed sets is also F_σ .)

- (c) Show that the function $f(x) = \text{dist}(x, A)$ is continuous.

13. Let $(X, \|\cdot\|)$ be a normed linear space.

- (a) * Prove that if $A \subset (X, \|\cdot\|)$ is compact and non-empty, then for each $x_0 \in X$, there exists a $y_0 \in A$ such that

$$\|x_0 - y_0\| = \inf\{\|x_0 - y\| \mid y \in A\}.$$

- (b) * Assume that X is finite dimensional. Prove that if $A \subset (X, \|\cdot\|)$ is closed and non-empty, then for each $x_0 \in X$, there exists a $y_0 \in A$ such that

$$\|x_0 - y_0\| = \inf\{\|x_0 - y\| \mid y \in A\}.$$

- (c) A subset A of a vector space is said to be convex if $\alpha x + (1 - \alpha)y \in A$ whenever $x, y \in A$ and $0 \leq \alpha \leq 1$.
Let $A \subseteq \mathbb{R}^2$ be convex and closed and let $x_0 \in A^c$. Show that if \mathbb{R}^2 is given the norm $\|\cdot\|_2$, then the point y_0 obtained in part 13b above is unique but that this need not be the case if we use

the norm $\|\cdot\|_\infty$.

- (d) Given $A, B \subseteq X$ non-empty sets, define $\text{dist}(A, B) = \inf\{d(a, b) \mid a \in A, b \in B\}$. Show that if A is closed, B is compact with $A \cap B = \emptyset$, then $\text{dist}(A, B) > 0$.
- (e) Show that even in \mathbb{R} , 13d can fail if you only assume that B is closed.
- (f) * Let $f(x) \in C[0, 1]$. Let

$$P_n = \{p(x) = a_0 + a_1x + \dots + a_nx^n \mid a_i \in \mathbb{R}\}.$$

Show that there exists a polynomial $p(x) \in P_n$ such that

$$\|f(x) - p(x)\|_\infty \leq \|f(x) - q(x)\|_\infty$$

for any $q(x) \in P_n$.

- (g) * Show that if $\{p_k(x)\}$ is a sequence of polynomials such that $\{p_k(x)\}$ converges uniformly to $f(x) = e^x$ on $[0, 1]$, then

$$\lim_{k \rightarrow \infty} \deg(p_k(x)) = \infty.$$

14. * Let $(V, \|\cdot\|)$ be an infinite dimensional Banach space.

- (a) Show that if $\mathcal{B} = \{v_\alpha\}_{\alpha \in I}$ is a basis for V , then I is uncountable. (Hint: Assume that $\mathcal{B} = \{v_1, v_2, v_3, \dots\}$ was countable. Let $F_n = \text{span}\{v_1, v_2, v_3, \dots, v_n\}$.)
- (b) Show that there exist a linear function $\varphi : V \rightarrow \mathbb{R}$ that is unbounded. (Hint: You can assume that V has a basis consisting of vectors of norm 1. From here you need only define φ on the basis elements and then extend it linearly.)

15. * Let $f(x)$ be continuous on $[0, 1]$. Assume that

$$\int_0^1 f(x) dx = 0$$

and that

$$\int_0^1 f(x)x^n dx = 0$$

for each $n \in \mathbb{N}$. Show that $f(x) = 0$ for all $x \in [0, 1]$.

16. Let $X = [0, 1] \times [0, 1] \subset (\mathbb{R}^2, \|\cdot\|_2)$. Let $f(x, y) \in C(X)$. For each $y \in [0, 1]$, define $f_y(x) = f(x, y)$ for each $x \in [0, 1]$.

(a) Show that $\mathcal{F} = \{f_y \mid y \in [0, 1]\}$ is equicontinuous.

(b) Show that the map $\Gamma : [0, 1] \rightarrow (C[0, 1], \|\cdot\|_\infty)$ given by

$$\Gamma(y) = f_y$$

is continuous.

(c) Is \mathcal{F} compact in $C(X)$? Explain your answer.

17. Let

$$\Psi = \left\{ F(x, y) \in C([0, 1] \times [0, 1]) \mid F(x, y) = \sum_{i=1}^k f_i(x)g_i(y) \right\}$$

where in the sum above, the functions f_i and g_i are continuous on $[0, 1]$. Show that Ψ is dense in $C([0, 1] \times [0, 1])$.

18. Let $g(x)$ be continuous and strictly increasing on $[a, b]$. Let $f(x) \in C[a, b]$. Let $\varepsilon > 0$. Then there exists constants c_0, c_1, \dots, c_n such that

$$\left| f(x) - \sum_{k=0}^n c_k g^k(x) \right| < \varepsilon$$

for each $x \in [a, b]$.

19. Let I be a closed ideal of $C[0, 1]$. (That is, I is a closed subalgebra of $C[0, 1]$ with the property that if $g(x) \in I$ and if $f(x) \in C[0, 1]$, then $f(x)g(x) \in I$.)

(a) Let $Z(I) = \{x \in [0, 1] \mid \forall f \in I, f(x) = 0\}$. Show that $Z(I)$ is a closed subset of $[0, 1]$.

(b) Show that if $Z(I) = \emptyset$, then $I = C[0, 1]$. (Hint: Show that there exists a function $f(x) \in I$ such that $f(x) > 0$ for every $x \in [0, 1]$.)

(c) Let $A \subseteq [0, 1]$ be closed. Let $I(A) = \{f \in C[0, 1] \mid \forall x \in A, f(x) = 0\}$. Show that I is a maximal closed ideal in $C[0, 1]$ if and only if $I = I(\{x_0\})$ for some $x_0 \in [0, 1]$.

(Recall: A closed ideal I is maximal if $I \neq C[0, 1]$ and if J is any closed ideal containing I , then either $I = J$ or $J = C[0, 1]$.)

Bibliography

Forrest, B. E. (2018). Pmath351, real analysis.

Index

- F_σ Sets, 147
- G_δ Sets, 147
- ε -net, 166
- $\|\cdot\|_p$ -norm, 67
- d -topology, 84
- 1-norm, 64
- 2-metric, 62
- 2-norm, 64

- Anti-symmetric, 29
- Arzelà-Ascoli, 210
- Axiom of Choice, 27

- Baire Category Theorem I, 149
- Baire Category Theorem II, 151
- Banach Contractive Mapping Theorem, 142
- Banach space, 123
- Banach-Mazurkiewickz Theorem, 190
- Bolzano-Weierstrass, 117
- Bolzano-Weierstrass Property, 162
- Boundary Point, 92
- bounded above, 32
- Bounded Linear Map, 175
- Boundedness, 117

- Cantor Set, 89
- Cantor Ternary Set, 89
- Cantor's Diagonal Argument, 49
- Cantor's Intersection Principle, 127
- Cantor-Schröder-Bernstein Theorem, 47
- Cardinality, 41
- Cauchy, 115
- Cauchy-Schwarz Inequality, 68
- Chains, 32
- Choice Function, 25
- closed, 82
- closed ball, 82
- Closure, 91
- cluster points, 97
- Compact, 158
- Completeness, 116
- Completion, 135
- Completion Theorem, 137
- Continuity, 108, 114
- Continuity on a Space, 111
- Continuum Hypothesis, 60
- Contraction, 141
- converge pointwise, 120
- converge uniformly, 121
- Convergence, 101
- Convergent Sequences are Cauchy, 115

Countable, 44
 Cover, 158

 De Morgan's Laws, 22
 Dense, 95
 denumerable, 44
 Diameter, 125
 Discrete Metric, 62
 discrete topology, 85
 diverges, 101
 domain, 24

 Empty Set, 21
 Equicontinuity, 207
 Equivalence Class, 38
 Equivalence Relation, 38
 Equivalent Metric Spaces, 113
 Euclidean Metric, 62
 Euclidean norm, 64
 Exponentiation of Cardinals, 57
 Extreme Value Theorem, 171

 Finite Intersection Property, 163
 Finite Sets, 41
 First Category, 148
 Fixed Point, 139
 Formal Sum, 129
 Function, 24

 Generalized Continuum Hypothesis, 60

 Hölder's Inequality, 68
 Hölder's Inequality v_2 , 77
 Hasse diagram, 31
 Heine-Borel Property, 181
 Heine-Borel Theorem, 158

 Homeomorphism, 113

 indiscrete topology, 85
 Induced Metric, 107
 Induced Topology, 107
 Infinite Sets, 42
 Interior, 91
 Intermediate Value Theorem, 84
 Intersection, 20
 Isometry, 135

 Lattice, 194
 least upper bound, 32
 Least Upper Bound Property of \mathbb{R} , 33
 Lebesgue, 171
 Lebesgue Number, 172
 Lebesgue-Borel, 172
 Limit Points, 97
 Lipschitz, 141

 Maximal Element, 35
 maximum, 32
 Metric, 61
 Metric Space, 61
 metric topology, 84
 Minkowski's Inequality, 69
 Minkowski's Inequality v_2 , 78
 Moments, 189
 Monotone Convergence Theorem, 213
 Multiplication of Cardinals, 56

 Neighbourhood, 91
 Nested Interval Theorem, 124
 Norm, 63
 Normed Linear Space, 63

Nowhere Dense, 148
 open, 82
 open ball, 82
 partial sum, 129
 Partially Ordered Sets, 30
 Partition, 39
 Picard's Theorem, 144
 piecewise linear function, 195
 piecewise polynomial, 195
 Pigeonhole Principle, 41
 Point-Separating, 194
 Points of Discontinuity, 145
 Pointwise Bounded Functions,
 209
 Poset, 30
 Power Set, 21
 Product of Sets, 23
 range, 24
 Reflexive, 29
 Relation, 24
 Relations, 29
 Relatively Compact Sets, 209
 Residual, 148
 Russell's Paradox, 59
 Second Category, 148
 self-adjoint, 204
 Separable, 94
 Sequence Spaces, 72
 Sequential Characterization of
 Continuity, 110
 Sequential Compactness, 162
 Set Complement, 21
 Set Difference, 20
 Standard Metric, 61
 Stone-Weierstrass Theorem —
 Complex Version, 204
 Stone-Weierstrass Theorem —
 Lattice Version, 196
 Stone-Weierstrass Theorem —
 Subalgebra Version, 201
 Subalgebra, 197
 subcover, 158
 Sum of Cardinals, 55
 supremum, 32
 Symmetric, 29
 Symmetric Difference, 20
 topological space, 84
 Topology, 84
 Totally Bounded, 166
 Totally Ordered Sets, 32
 Transitive, 29
 tuples, 23
 Uncountable, 49
 Uniform Continuity, 136
 Uniformly Bounded Functions,
 209
 uniformly equicontinuous, 207
 Union, 20
 Universal Set, 19
 upper bound, 32
 Weierstrass Approximation Theo-
 rem, 188
 Weierstrass M-test, 129
 Well-Ordered, 36
 Well-Ordering Principle, 37
 Zermelo's Axiom of Choice, 27

Todo list

I need to get a better picture of the motivation of the proof. . . .	196
What else can we understand from $\varphi_{x,y}$?	196
I should find out about this.	203
how?	203