

STAT330S18 - Mathematical Statistics

CLASSNOTES FOR SPRING 2018

by

Johnson Ng

BMath (Hons), Pure Mathematics major, Actuarial Science Minor

University of Waterloo

Table of Contents



List of Definitions	8
List of Theorems	11
1 Lecture 1 May 1st 2018	15
1.1 Introduction	15
1.2 Random Variable	20
1.3 Discrete Random Variable	22
2 Lecture 2 May 03rd 2018	27
2.1 Continuous Random Variable	27
2.2 Examples of Discrete RVs	28
2.2.1 Binomial Distribution	29
2.2.2 Geometric Distribution	29
2.2.3 Poisson Distribution	30
2.3 Examples of Continuous RVs	30
2.3.1 Normal/Gaussian Distribution	30
2.3.2 Uniform Distribution	31
2.3.3 Exponential Distribution	31
2.3.4 Gamma Distribution	31
2.4 Functions of Random Variables	32
2.4.1 Discrete X and Discrete Y	33
2.4.2 Continuous X and Discrete Y	34
2.4.3 Continuous X and Continuous Y	35
2.4.4 A Formula for the Continuous Case	36
3 Lecture 3 May 08th 2018	39
3.1 Functions of Random Variables (Continued)	39
3.1.1 Special Cases	39

3.2	Probability Integral Transformation	40
3.3	Location-Scale Families	41
3.4	Expectations	44
3.4.1	Expectations	44
4	Lecture 4 May 10th 2018	47
4.1	Expectations (Continued)	47
4.1.1	Expectations (Continued)	47
4.1.2	Moments and Variance	50
4.2	Inequalities	52
4.2.1	Markov/Chebyshev Style Inequalities	52
5	Lecture 5 May 15th 2018	55
5.1	Inequalities (Continued)	55
5.1.1	Markov/Chebyshev Style Inequalities (Continued)	55
5.2	Moment Generating Function	56
5.2.1	MGF of a Linear Transformation	60
5.2.2	Uniqueness of the MGF	61
6	Lecture 6 May 17th 2018	63
6.1	Joint Distributions	63
6.1.1	Introduction to Joint Distributions	63
6.1.2	Joint and Marginal CDFs	63
6.1.3	Joint Discrete RVs	66
6.1.4	Independence of Discrete RVs	70
7	Lecture 7 May 24th 2018	73
7.1	Joint Distributions (Continued)	73
7.1.1	Independence of Discrete RVs (Continued)	73
7.1.2	Joint Continuous RVs	73
7.1.3	Marginal Distribution (Continuous)	77
7.1.4	Independence of Continuous RVs	78
8	Lecture 8 May 29th 2018	81
8.1	Joint Distributions (Continued 2)	81
8.1.1	Independence of Continuous RVs (Continued)	81
8.1.2	Conditional Distributions	83
8.1.3	Joint Expectations	89
9	Lecture 9 May 31st 2018	91
9.1	Joint Distributions (Continued 3)	91
9.1.1	Joint Expectations (Continued)	91





















9.1.2	Covariance	92
9.1.3	Correlation	98
9.1.4	Conditional Expectation	100
10	Lecture 10 Jun 05th 2018	103
10.1	Joint Distribution (Continued 4)	103
10.1.1	Conditional Expectation (Continued)	103
10.1.2	Joint Moment Generating Functions	107
11	Lecture 11 Jun 07th 2018	111
11.0.1	Working with Multivariate Cases	111
12	Lecture 12 Jun 12th 2018	117
12.1	Functions of Random Variables	117
12.1.1	Transformation of Two or More Random Variables	117
12.1.2	One-to-One Bivariate Transformations	120
13	Lecture 13 Jun 14th 2018	125
13.1	Functions of Random Variables (Continued)	125
13.1.1	One to One Bivariate Transformations (Continued)	125
13.1.2	Moment Generating Function Method	126
14	Lecture 14 Jun 19th 2018	131
14.1	Functions of Random Variables (Continued 2)	131
14.1.1	Moment Generating Function Method (Continued)	131
15	Lecture 15 Jun 21st 2018	135
15.1	Limiting or Asymptotic Distributions	135
15.1.1	Convergence in Distribution	135
15.1.2	Convergence in Probability	139
16	Lecture 16 Jun 26th 2018	141
16.1	Limiting or Asymptotic Distributions (Continued)	141
16.1.1	Convegence in Probability (Continued)	141
16.1.2	Limit Theorems	146
17	Lecture 17 Jun 28th 2018	157
17.1	Estimation	157
17.1.1	Maximum Likelihood Estimation	158
18	Lecture 18 Jul 3rd 2018	163
18.1	Estimation (Continued)	163
18.1.1	Maximum Likelihood Estimation (Continued)	163

19 Lecture 19 July 5th 2018	171
19.1 Estimation (Continued 2)	171
19.1.1 Maximum Likelihood Estimation (Continued 2)	171
20 Lecture 20 Jul 10th 2018	179
20.1 Estimation (Continued 3)	179
20.1.1 Maximum Likelihood Estimation (Continued 3)	179
20.1.2 Asymptotic Properties of ML Estimators	182
21 Lecture 21 Jul 12th 2018	187
21.1 Estimation (Continued 4)	187
21.1.1 Asymptotic Properties of ML Estimators (Continued)	187
21.1.2 Interval Estimators	188
22 Lecture 22 Jul 17th 2018	195
22.1 Estimate (Continued 5)	195
22.1.1 Interval Estimators (Continued)	195
23 Lecture 23 Jul 19th 2018	201
23.1 Hypothesis Testing	201
23.1.1 Introduction	201
23.1.2 Likelihood Ratio Tests for Simple Hypothesis	204
A Regularity Conditions	211
B Useful References	213
B.1 Commonly Used Distributions	213
Bibliography	217
Index	219

List of Definitions

1	 Definition (Sample Space)	15
2	 Definition (σ -field)	15
3	 Definition (Measurable Space)	16
4	 Definition (Probability Measure)	16
5	 Definition (Conditional Probability)	19
6	 Definition (Independent Events)	19
7	 Definition (Random Variable)	20
8	 Definition (Cumulative Distribution Function)	21
9	 Definition (Discrete Random Variable)	22
10	 Definition (Continuous Random Variable)	27
11	 Definition (Binomial RV)	29
12	 Definition (Geometric RV)	29
13	 Definition (Poisson RV)	30
14	 Definition (Normal / Gaussian RV)	30
15	 Definition (Standard Normal Distribution)	31
16	 Definition (Uniform RV)	31
17	 Definition (Exponential RV)	31
18	 Definition (Gamma RV)	32
19	 Definition (Location Parameter and Family)	41
20	 Definition (Scale Parameter and Family)	42
21	 Definition (Location-Scale Family)	42
22	 Definition (Expectation of A Discrete RV)	44
23	 Definition (Expectation of A Continuous RV)	45
24	 Definition (Variance)	50
25	 Definition (Moments)	51
26	 Definition (Moment Generating Function)	56
27	 Definition (Indicator Function)	57
28	 Definition (Joint CDF)	63

29	Definition (Marginal CDF)	66
30	Definition (Joint Discrete RV)	66
31	Definition (Marginal Distribution - Discrete Case)	68
32	Definition (Independence of Discrete RVs)	70
33	Definition (Joint Continuous RVs)	73
34	Definition (Marginal PDF)	77
35	Definition (Independence of Continuous RVs)	78
36	Definition (Conditional Distributions)	83
37	Definition (Beta Distribution)	88
38	Definition (Joint Expectation)	89
39	Definition (Covariance)	93
40	Definition (Correlation Coefficient)	99
41	Definition (Conditional Expectation)	101
42	Definition (Joint Moment Generating Functions)	107
43	Definition (Joint Moments and Marginal MGF)	107
44	Definition (k-variate CDF)	111
45	Definition (k-variate Support Set)	111
46	Definition (k-Variate Marginal Distribution)	112
47	Definition (k-Variate Joint MGF)	112
48	Definition (Multinomial Distribution)	113
49	Definition (Bivariate Normal Distribution)	114
50	Definition (Random Sample (IID))	115
51	Definition (One-to-One Bivariate Transformation)	120
52	Definition (Convergence in Distribution)	135
53	Definition (Degenerate Distribution)	138
54	Definition (Convergence in Probability)	139
55	Definition (Convergence in Probability to a Constant)	143
56	Definition (Double Parameter Exponential Distribution)	144
57	Definition (Statistic)	157
58	Definition (Estimators and Estimates)	158
59	Definition (Likelihood function)	159
60	Definition (Score Function)	163
61	Definition (Information Function)	163

62	 Definition (Fisher Information)	164
63	 Definition (Relative Likelihood)	171
64	 Definition (Likelihood Region & Likelihood Interval)	172
65	 Definition (Log Relative Likelihood)	172
66	 Definition (Score Vector)	175
67	 Definition (Information Matrix)	175
68	 Definition (Fisher Information Matrix)	175
69	 Definition (Likelihood Region - Multivariate)	175
70	 Definition (Interval Estimators)	188
71	 Definition (Pivotal Quantity)	188
72	 Definition (Confidence Interval)	189
73	 Definition (Likelihood Interval)	198
74	 Definition (Hypothesis)	201
75	 Definition (Null Hypothesis and Alternative Hypothesis)	201
76	 Definition (Test of Hypothesis)	202
77	 Definition (Test Statistic)	202
78	 Definition (Significance Level and p -Values)	202
79	 Definition (Type I and Type II Errors)	203
80	 Definition (Simple Hypothesis)	204
81	 Definition (Likelihood Ratio Statistic)	204

List of Theorems

1	 Proposition (Properties of Probability Set Functions)	17
2	 Proposition (Boole's Inequality)	19
3	 Proposition (Bonferroni's Inequality)	20
4	 Proposition (Continuity Property)	20
5	 Proposition (Properties of the cdf)	21
6	 Proposition (Properties of pmf)	22
7	 Proposition (Properties of pdf)	27
8	 Theorem (One-to-One Transformation of a Random Variable)	36
9	 Theorem (Probability Integral Transformation)	40
10	 Theorem (Converse of Probability Integral Transformation)	41
11	 Theorem (Expectation from the cdf)	47
12	 Theorem (Expected Value of a Function of X)	48
13	 Theorem (Linearity of Expectation)	49
14	 Theorem (Variance of a Linear Function)	51
15	 Theorem (Markov's Inequality)	52
16	 Theorem (Markov's Inequality 2)	53
17	 Theorem (Chebyshev's Inequality)	54
18	 Proposition (Properties of the MGF)	58
19	 Theorem (MGF of a Linear Transformation)	60
20	 Theorem (Uniqueness of the MGF)	61
21	 Proposition (Properties of Joint CDF)	64
22	 Proposition (Properties of Joint PMF)	67
23	 Theorem (Independence by PF)	71
25	 Proposition (Properties of Joint PDF)	74
26	 Theorem (Factorization Theorem for Independence)	82
27	 Proposition (Properties of Conditional Distributions)	86
28	 Theorem (Product Rule)	87

29	 Proposition (Independence from Conditionality)	87
30	 Theorem (Linearity of Expectation in Bivariate Case)	91
31	 Theorem (Implication of Independence on Joint Expectation)	91
32	 Theorem (Generalized Implication of Independence on Joint Expectation)	92
33	 Theorem (Variance of Linear Combinations)	96
34	 Theorem (Generalized Variance of Linear Combinations)	97
35	 Proposition (Properties of the Correlation Coefficient)	99
36	 Proposition (Independence on Conditional Expectation)	104
37	 Theorem (Law of Total Expectation)	104
38	 Theorem (Law of Total Variance)	106
39	 Proposition (Independence on Joint MGF)	109
40	 Proposition (Law of Total Probability - Multivariate)	111
41	 Proposition (Independence for Multivariate Cases)	112
42	 Proposition (Properties of Multinomial Distribution)	113
43	 Proposition (Properties of Bivariate Normal Distribution)	114
44	 Theorem (One-to-One Bivariate Transformations)	121
45	 Proposition (Properties of the Jacobian)	122
46	 Theorem (Sums of MGF)	126
47	 Theorem (Gaussian Distribution)	131
48	 Theorem (Properties of the Gaussian Distribution)	131
49	 Theorem (F Distribution)	132
50	 Theorem (Taylor Series with Lagrange's Remainder)	135
51	 Theorem (Generalized Limit Definition of e)	136
52	 Corollary (Limit definition of e)	137
53	 Lemma	141
54	 Proposition (Convergence in Probability Implies Convergence in Distribution)	142
55	 Proposition (Partial Converse of  Proposition 54)	144
56	 Proposition (Convergence in Distribution and MGF)	146
57	 Theorem (Central Limit Theorem)	148
58	 Proposition (Other Limit Theorems)	151
59	 Theorem (Generalized δ -Method)	154
60	 Corollary (δ -Method)	155
61	 Theorem (Invariance Property of the MLE)	169

Foreword

The proofs in this set of notes will be more rigorous compared to the expectations of the course (at least, for the course this term). If you are not the author and are interested in reading the notes, you may skip the proofs should you have little interest in them. The rigour is required almost exclusively for the author himself, for his own practice, and because he transferred his STAT230 course from a class that is clean of proofs.

Also, many of the common mathematical notations will be heavily used both in the author's notes and proofs. The author cannot guarantee that his proofs are absolutely correct, but he tries, to the best of his abilities to minimize and assure of that. Should you be suspicious about the proofs, or should you notice erroneous ones, please do inform the author at https://github.com/japorized/TeX_notes/issues.

You are also warned that the author is rather bummed with how the course is presented in the term that he is/was taking it, and so there may be sarcastic language (towards the lectures) mixed in his notes.

1 Lecture 1 May 1st 2018

1.1 Introduction

Definition 1 (Sample Space)

A *sample space*, S of a random experiment is the set of all possible outcomes of the experiment.

Example 1.1.1

The following are some random experiments and their sample space.

- Flipping a coin
 $S = \{H, T\}$ where H denotes head and T tail.
- Rolling a 6-faced dice twice
 $S = \{(x, y) : x, y \in \mathbb{N}, 1 \leq x, y \leq 6\}$
- Measuring a patient's height
 $S = \mathbb{R}^+ = \{x \in \mathbb{R} : x \geq 0\}$

Definition 2 (σ -field)

Let S be a sample space. The collection of sets $\mathcal{B} \subseteq \mathbb{P}(S)$ ¹, is called a σ -field (or σ -algebra) on S if:

1. $\emptyset \in \mathcal{B}$ and $S \in \mathcal{B}$;
2. $\forall A \in \mathcal{B} \quad A^C \in \mathcal{B}$; ² and
3. $\forall n \in \mathbb{N} \quad \forall \{A_j\}_{j=1}^n \subseteq \mathcal{B} \quad \bigcup_{j=1}^n A_j \in \mathcal{B}$.

¹ The **power set** of S , $\mathbb{P}(S)$, is defined as the set that contains all subsets of S .

² We shall denote the compliment of a set by a superscript C in this set of notes. The supplemental notes provided in the class uses an overhead bar, e.g. \bar{A} , while lecture notes will use A^C and A' interchangeably.

Definition 3 (Measurable Space)

Given that S is a non-empty set, and \mathcal{B} is a σ -field, (S, \mathcal{B}) is a *measurable space*.³

³ A measurable space is a basic object in **measure theory**.

Example 1.1.2

Consider $S = \{1, 2, 3, 4\}$. Check if $\mathcal{B} = \{\emptyset, \{1, 2, 3, 4\}, \{1, 2\}, \{3, 4\}\}$ is a σ -field on S .

1. It is clear that $\emptyset, S \in \mathcal{B}$.
2. Note that $S^C = \emptyset$ and $\{1, 2\}^C = \{3, 4\}$.
3. Note that the largest possible result of any countable union of the elements of \mathcal{B} is $\{1, 2, 3, 4\}$, which is an element of \mathcal{B} .

BECAUSE (S, \mathcal{B}) is a measurable space, we can define a measure on it.

Definition 4 (Probability Measure)

Suppose S is a sample space of a random experiment. Let $\mathcal{B} = \{A_1, A_2, \dots\} \subseteq \mathbb{P}(S)$ be the σ -field on S . The **probability set function** (or **probability measure**), $P : \mathcal{B} \rightarrow [0, 1]$, is a function that satisfies the following:⁴

- $\forall A \in \mathcal{B} \ P(A) \geq 0$;
- $P(S) = 1$;
- $\forall \{A_j\}_{j=1}^{\infty} \subseteq \mathcal{B} \ \forall i \neq j \in \mathbb{N} \ A_i \cap A_j = \emptyset \implies$

$$P\left(\bigcup_{j=1}^{\infty} A_j\right) = \sum_{j=1}^{\infty} P(A_j) \quad (1.1)$$

⁴ These conditions are also known as **Kolmogorov Axioms**, or **probability axioms**.

(S, \mathcal{B}, P) is called a **probability space**.

Example 1.1.3

Consider flipping a coin where $S = \{H, T\}$. Let P be defined as follows

$$P(\{H\}) = \frac{1}{3} \quad P(\{T\}) = \frac{2}{3} \quad P(\emptyset) = 0 \quad P(S) = 1$$

Conditions 1 and 2 of [Definition 4](#) are met. Notice that

$$P(\{H\} \cup \{T\}) = P(S) = 1 \quad \text{and} \quad P(\{H\}) + P(\{T\}) = \frac{1}{3} + \frac{2}{3} = 1.$$

Hence condition 3 is also fulfilled.

♦ Proposition 1 (Properties of Probability Set Functions)

Let P be a probability set function and A, B be any set in \mathcal{B} . Prove the following:⁵

1. $P(A^C) = 1 - P(A)$
2. $P(\emptyset) = 0$
3. $P(A) \leq 1$
4. $P(A \cap B^C) = P(A) - P(A \cap B)$
5. $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
6. $A \subseteq B \implies P(A) \leq P(B)$

⁵ Many among these properties illustrate that the probability is indeed a **measure**.

Exercise 1.1.1

Prove that $A \subseteq B \iff B^C \subseteq A^C$.

✎ Proof

Let S be the sample space for P .

1. Note that

$$A \in \mathcal{B} \implies A \in \mathbb{P}(S) \iff A \subseteq S$$

$A \in \mathcal{B} \iff A^C \in \mathcal{B} \implies A^C \subseteq S$. Also, since A^C is the complement of A , it is clear that $S = A \cup A^C$.

$$\therefore P(S) = 1 \iff P(A \cup A^C) = 1 \xrightarrow{1} P(A) + P(A^C) = 1$$

where 1 is by condition 3 in [Definition 4](#) since $A \cap A^C = \emptyset$ by definition of a complement of a set.

2. Note that $S \cup \emptyset = S$ and $S \cap \emptyset = \emptyset$. Using a similar argument as

above,

$$1 = P(S) = P(S \cup \emptyset) = P(S) + P(\emptyset) \implies P(\emptyset) = 0$$

3. By 1 from above, $P(A) = 1 - P(A^C)$. Since $0 \leq P(A^C) \leq 1$, we have that $P(A)$ is at most 1, as required.
4. Note that $A = (A \cap B) \cup (A \cap B^C)$. Clearly, $(A \cap B) \cap (A \cap B^C) = \emptyset$.⁶ Hence by condition 3 in [Definition 4](#),

$$P(A) = P(A \cap B) + P(A \cap B^C)$$

⁶ This is an easy proof using the basic way of proving membership.

5. Consider $P(A \cup B) + P(A \cap B)$. By definition,

$$A \cup B = (A \cap B^C) \cup (A \cap B) \cup (A^C \cap B)$$

where each of the sets in brackets are disjoint from each other⁷.

⁷ Again, this is not hard to show

By condition 3 of [Definition 4](#), we would then have

$$\begin{aligned} P(A \cup B) + P(A \cap B) &= P(A \cap B^C) + P(A \cap B) + P(A^C \cap B) + P(A \cap B) \\ &= 2P(A \cap B) + P(A) - P(A \cap B) + P(B) - P(A \cap B) \text{ by 4} \\ &= P(A) + P(B) \end{aligned}$$

6. Note that $B = B \cap S = B \cap (A^C \cup A) = (B \cap A^C) \cup A$. Clearly, $A \cap (B \cap A^C) = \emptyset$. By condition 3 in [Definition 4](#), we thus have that

$$P(B) = P(B \cap A^C) + P(A). \quad (\dagger)$$

Suppose $A \subsetneq B$. Then $B \cap A^C \neq \emptyset$. I shall make the claim that $B \cap A^C \in \mathcal{B}$. Since $A \subseteq B$ we have that

$$\begin{aligned} a \in (B \cap A^C) &\iff a \in B \wedge a \in A^C \\ &\iff a \in B \wedge a \notin A \\ &\iff a \in (B \setminus A). \end{aligned}$$

But $B \setminus A$ is a subset of B from the above steps⁸. Therefore, $(B \cap A^C) \subseteq B \in \mathcal{B}$ as required.

⁸ This is rather obvious from the steps, since $\forall a \in (B \cap A^C), a \in B$.

With that done, by condition 1 in [Definition 4](#), $P(B \cap A^C) \geq 0$.

Hence from Equation (+), we have that

$$\begin{aligned} P(B) &= P(B \cap A^C) + P(A) \\ &\geq P(A) \end{aligned}$$

as required. \square

Definition 5 (Conditional Probability)

Suppose S is a sample space of a random experiment, and $A, B \subseteq S$. The **conditional probability of A given B** is given by

$$P(A|B) = \frac{P(A \cap B)}{P(B)} \quad \text{provided } P(B) > 0. \quad (1.2)$$

Definition 6 (Independent Events)

Suppose S is a sample space of a random experiment, and $A, B \subseteq S$. A and B are said to be **independent of each other** if

$$P(A \cap B) = P(A)P(B)$$

Proposition 2 (Boole's Inequality)

If $\{A_j\}_{j=1}^{\infty}$ is a sequence of events, then

$$P\left(\bigcup_{j=1}^{\infty} A_j\right) \leq \sum_{j=1}^{\infty} P(A_j)$$

Proof

Proof shall be provided later

Proposition 3 (Bonferroni's Inequality)

If $\{A_j\}_{j=1}^k$ is a set of events where $k \in \mathbb{N}$, then

$$P\left(\bigcap_{j=1}^k A_j\right) \geq 1 - \sum_{j=1}^k P(A_j^c)$$

 **Proof**

Proof shall be provided later

 **Proposition 4 (Continuity Property)**

If $A_1 \subset A_2 \subset \dots$ is a sequence where $A = \bigcup_{i=1}^{\infty} A_i$, then

$$\lim_{n \rightarrow \infty} P\left(\bigcup_{i=1}^n A_i\right) = P(A)$$

 **Proof**

Proof shall be provided later

1.2 Random Variable

 **Definition 7 (Random Variable)**

In a given probability space (S, \mathcal{B}, P) , the function $X : S \rightarrow \mathbb{R}$ is called a **random variable**⁹ if

$$P(X \leq x) = P(\{\omega \in S : X(\omega) \leq x\}) \quad (1.3)$$

is defined for all $x \in \mathbb{R}$ ¹⁰.

⁹ We shall use rv as shorthand for random variable in this set of notes.

¹⁰ $X \leq x$ is an abbreviation for $\{\omega \in S : X(\omega) \leq x\} \in \mathcal{B}$.

Example 1.2.1

In a coin flip experiment, we have that $S = \{H, T\}$ where $\mathbb{P}(S) =$

$\{\emptyset, S, \{H\}, \{T\}\}$. Define X : the number of heads in a flip, i.e.

$$X(\{H\}) = 1 \text{ and } X(\{T\}) = 0$$

To prove why X is a random variable given this definition, notice that

$$\begin{aligned} x < 0 &\implies P(X \leq x) = P(\{\omega \in S : X(\omega) < 0\}) = P(\emptyset) = 0 \\ x \geq 1 &\implies P(X \leq x) = P(\{\omega \in S : X(\omega) \leq x\}) = P(\{H, T\}) \\ &= P(\{H\}) + P(\{T\}) = 1 \text{ by Independence} \\ 0 \leq x < 1 &\implies P(X \leq x) = P(\{\omega \in S : X(\omega) \leq x\}) = P(T) \geq 0 \end{aligned}$$

which shows that P is defined for all $x \in \mathbb{R}$. Hence X is a random variable.

Definition 8 (Cumulative Distribution Function)

The **cumulative distribution function (c.d.f)** of a random variable X is defined as

$$\forall x \in \mathbb{R} \quad F(x) = P(X \leq x)$$

“ Note

NOTICE that $F(x)$ is defined for **all** real numbers, and since it is a probability, we have $0 \leq F(x) \leq 1$.

Proposition 5 (Properties of the cdf)


1. $\forall x_1 < x_2 \in \mathbb{R} \quad F(x_1) \leq F(x_2)$
2. $\lim_{x \rightarrow -\infty} F(x) = 0 \wedge \lim_{x \rightarrow \infty} F(x) = 1$
3. $\lim_{x \rightarrow a^+} F(x) = F(a)$ ¹¹
4. $\forall a < b \in \mathbb{R} \quad P(a < X \leq b) = P(X \leq b) - P(X \leq a) = F(b) - F(a)$
5. $P(X = b) = F(b) - \lim_{a \rightarrow b^-} F(a)$ ¹²

¹¹ F is a **right-continuous** function.

¹² This is also called the **magnitude of the jump**.

 **Proof****Proof shall be provided later****“ Note**

The definition and properties of the cdf hold for the rv X regardless of whether S is discrete (finite or countable) or not.


1.3 Discrete Random Variable **Definition 9 (Discrete Random Variable)**

An rv X is a **discrete random variable** when its image is finite or countably infinite, i.e. $X \in \{x_1, x_2, \dots\}$. The function

$$\forall x \in \mathbb{R} \quad f(x) := P(X = x) = F(x) - \lim_{\varepsilon \rightarrow 0^+} F(x - \varepsilon)$$

is its probability function, commonly known as the **probability mass function** (pmf). The set $A := \{x : f(x) > 0\}$ is called the **support set** of X , and

$$\sum_{x \in A} f(x) = \sum_{i=1}^{\infty} f(x_i) = 1. \quad (1.4)$$

 **Proposition 6 (Properties of pmf)**

With the notation from  Definition 9, prove that

1. $\forall x \in \mathbb{R} \quad f(x) \geq 0$
2. $\sum_{x \in A} f(x) = 1$

 **Proof**

1. This result follows from the fact that f is a pdf, a probability, i.e. $\forall x \in \mathbb{R}, f(x) = 0$ is $x \notin S$ where S is the sample space, and $0 \leq f(x) \leq 1$ if $x \in S$.

2. Since $A = \{x : f(x) > 0\}$, we know that

$$\sum_{x \in A} f(x) > 0.$$

If we consider all the elements of A , we have that the events $(X = x_i)$, for $x_i \in A$, constitutes the entire sample space.

Therefore,

$$\sum_{x \in A} f(x) = \sum_{x \in A} P(X = x) = P(S) = 1.$$

□

Exercise 1.3.1

Consider an urn containing r red marbles and b black marbles. Find the pmf of the rv for the following:

1. $X =$ number of red balls in n selections without replacement.
2. $X =$ number of red balls in n selections with replacement.
3. $X =$ number of black balls selected before obtaining the first red ball if sampling is done with replacement.
4. $X =$ number of black balls selected before obtaining the k th red ball if sampling is done with replacement.

Solution

1. Let $d = \max\{n, r + b\}$. The desired pmf is therefore the pmf from the hypergeometric distribution

$$\forall x \in \mathbb{Z}_{\leq r}^+ \quad f(x) = \frac{\binom{r}{x} \binom{b}{d-x}}{\binom{r+b}{d}}.$$

2. $\forall x \in \mathbb{Z}^+ \quad f(x) = \binom{n}{x} \left(\frac{r}{r+b}\right)^x \left(\frac{b}{r+b}\right)^{n-x}$, which is the pmf of the binomial distribution.
3. $\forall x \in \mathbb{Z}^+ \quad f(x) = \left(\frac{b}{r+b}\right)^x \left(\frac{r}{r+b}\right)$
4. $\forall x \in \mathbb{Z}^+ \quad f(x) = \binom{x+k-1}{k-1} \left(\frac{b}{r+b}\right)^x \left(\frac{r}{r+b}\right)^k$

Example 1.3.1

Consider the function

$$f(x) = \begin{cases} \frac{C\mu^x}{x!} & x \in \mathbb{Z}^+, \mu > 0 \\ 0 & \text{otherwise} \end{cases}$$

Find C such that $f(x)$ is a pmf for the rv X .

 **Solution**

We have that

$$\begin{aligned} 1 &= \sum_{x \in \mathbb{Z}^+} \frac{C\mu^x}{x!} \\ &= C \sum_{x \in \mathbb{Z}^+} \frac{\mu^x}{x!} \\ &= Ce^\mu \end{aligned}$$

This gives us that $\forall x \in \mathbb{Z}^+, f(x) = \frac{e^{-\mu}\mu^x}{x!}$, and this is, of course, the pmf of the **Poisson distribution**.

Thus $C = e^{-\mu}$.

Exercise 1.3.2

Prove that the pdf of $X \sim \text{Poi}(\mu)$ sums to 1 over all of its values.

 **Solution**

$$\begin{aligned} \sum_{x \in \mathbb{N}} \frac{\mu^x e^{-\mu}}{x!} &= e^{-\mu} \sum_{x \in \mathbb{N}} \frac{\mu^x}{x!} \\ &= e^{-\mu} e^\mu \quad \because \sum_{x \in \mathbb{N}} \frac{k^x}{x!} = e^k \\ &= 1 \end{aligned}$$

Exercise 1.3.3

If X is a random variable with pmf

$$f(x) = \frac{-(1-p)^x}{x \log p}, \quad x = 1, 2, \dots; \quad 0 < p < 1,$$

show that

$$\sum_{x \in \mathbb{N}} f(x) = 1$$

 Solution

$$\begin{aligned}\sum_{x \in \mathbb{N}} \frac{-(1-p)^x}{x \log p} &= -\frac{1}{\log p} \sum_{x \in \mathbb{N}} \frac{(-1)^x (p-1)^x}{x} \\ &= -\frac{1}{\log p} \underbrace{\left[-(p-1) + \frac{(p-1)^2}{2} - \frac{(p-1)^3}{3} + \dots \right]}_{\text{Taylor expansion of } -\log p} \\ &= 1\end{aligned}$$

2 Lecture 2 May 03rd 2018

2.1 Continuous Random Variable

Definition 10 (Continuous Random Variable)

Suppose X is an rv with cdf F . If F is a continuous function for all $x \in \mathbb{R}$ and F is differentiable except possibly at countably many points, then X is a **continuous rv**. The probability function, or more commonly known as the **probability density function** (pdf), of X is $f(x) = F'(x)$ wherever F is differentiable on x and 0 otherwise.

The set $A = \{x : f(x) > 0\}$ is called the **support set** of X and

$$\int_{x \in A} f(x) dx = 1$$

Proposition 7 (Properties of pdf)

Let X be a random variable and f be its pdf.

1. $\forall x \in \mathbb{R} \quad f(x) \geq 0$
2. $\int_{-\infty}^{\infty} f(x) dx = 1$
3. $f(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{P(x \leq X \leq x+h)}{h}$ (if the limit exists)
4. $\forall x \in \mathbb{R} \quad F(x) = \int_{-\infty}^x f(t) dt$
5. $P(a < X \leq b) = \int_a^b f(x) dx = F(b) - F(a)$
6. $P(X = b) = F(b) - \lim_{a \rightarrow b^-} F(a) = F(b) - F(b) = 0$

✎ Proof

1. The argument of this proof is similar to that provided in **◆ Proposition 6**.
2. Same as above, except that the support set can now have complete intervals.
3. The first equation follows from the first principles of Calculus. The second equation follows by method of calculation using the cdf.
4. $F(x) = P(X \leq x) = \int_{-\infty}^x f(t) dt$.
5. This follows immediately from the above property.
6. The first part of the equation is a way to interpret the above property. The limit equates to $F(b)$ since F is continuous.

□

Example 2.1.1

Consider the function

$$f(x) = \begin{cases} \frac{\theta}{x^{\theta+1}} & x \geq 1 \\ 0 & x < 1 \end{cases}$$

For what values of θ is f a pdf?

✎ Solution

If f is a pdf, then $\theta \geq 0$. In fact, $\theta \neq 0$; otherwise f would be equivalently 0 for all $x \in \mathbb{R}$, which would imply that $\int_{\mathbb{R}} f = 0$, which is impossible. It remains to check if $\theta > 0$ is a safe choice. Now

$$\int_1^{\infty} \frac{\theta}{x^{\theta+1}} dx = -\frac{1}{x^{\theta}} \Big|_1^{\infty} = 1$$

Note that the above integral is valid because $\frac{1}{x^{\theta+1}} \leq \frac{1}{x}$. Therefore the choice of $\theta > 0$ is safe.

2.2.1 Binomial Distribution

Definition 11 (Binomial RV)

Consider X to be the number of successes in a sequence of n experiments where

1. experiments are **independent**;
2. the outcome of each experiment is a **binary** (e.g. success or failure); and
3. has the **probability of success**, p for each singular experiment.

X is called a **Binomial** rv, and we write $X \sim \text{Bin}(n, p)$ and its pmf is

$$P(X = x) = \begin{cases} \binom{n}{x} p^x (1-p)^{n-x} & x = 0, 1, 2, \dots, n \\ 0 & \text{otherwise} \end{cases}$$

2.2.2 Geometric Distribution

Definition 12 (Geometric RV)

Consider a sequence of independent success/failure (binary) experiments, each of which has a success probability of p . Let X be the **number of failures** before the **first success** is reached. We call X a **Geometric** rv, and we write $X \sim \text{Geo}(p)$, and its pmf is

$$P(X = x) = \begin{cases} (1-p)^x p & x = 0, 1, 2, \dots, n \\ 0 & \text{otherwise} \end{cases}$$

“ Note

Some authors would define the Geometric rv as:

Let X be the number of experiments until the first success.

But that really is just a play of words.

2.2.3 Poisson Distribution

Definition 13 (Poisson RV)

Suppose X is defined to be the number of occurrences of an event in a given time period. If the process on which the events occur satisfies the following:

1. The number of occurrences in non-overlapping intervals are independent of each other;
2. The probability of the occurrence of an event in a short interval of length h is proportional to h ;
3. For sufficiently short time periods of length h , the probability of 2 or more events occurring in the interval is negligible, i.e. almost zero;

then X is a **Poisson** rv, and we write $X \sim \text{Poi}(\lambda)$, with $\lambda > 0$, and the pmf is

$$P(X = x) = \begin{cases} \frac{e^{-\lambda} \lambda^x}{x!} & x = 0, 1, \dots \\ 0 & \text{otherwise} \end{cases}$$

2.3 Examples of Continuous RVs

2.3.1 Normal/Gaussian Distribution

Definition 14 (Normal / Gaussian RV)

The **Normal/Gaussian** Distribution is a continuous probability distribution that is symmetric about the mean, showing that data around the mean is more frequent than data far from the mean. If X is a **Normal-Gaussian** rv, we write $X \sim N(\mu, \sigma^2)$, and its pdf is

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \quad \text{for } x \in \mathbb{R}.$$

Definition 15 (Standard Normal Distribution)

The **Standard Normal Distribution** is the simplest case of a Normal Distribution. An rv Z is called the **Standard Normal** rv if $\mu = 0$ and $\sigma = 1$. We write $Z \sim N(0,1)$ and its pdf is

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} \quad \text{for } x \in \mathbb{R}.$$

2.3.2 Uniform Distribution

Definition 16 (Uniform RV)

If X represents the result of drawing a real number from an interval (a, b) , with $a < b$, such that all numbers in between are equally likely to be chosen, then X is called a **Uniform** rv, and we write $X \sim \text{Unif}(a, b)$, and its pdf is

$$f(x) = \begin{cases} \frac{1}{b-a} & x \in (a, b) \\ 0 & \text{otherwise} \end{cases}$$

2.3.3 Exponential Distribution

Definition 17 (Exponential RV)

Let X show the time between two consecutive events in a **Poisson process**, i.e. the 3 conditions in Poisson RV are satisfied. Then X is called an **Exponential** rv, and we write $X \sim \text{Exp}(\theta)$, where $\theta > 0$, with its pdf

$$f(x) = \begin{cases} \frac{1}{\theta} e^{-\frac{x}{\theta}} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$

2.3.4 Gamma Distribution

Definition 18 (Gamma RV)

Let X be the sum of n independent **Exponential** rvs with some fixed θ . Then X is called a **Gamma** rv, in which we write $X \sim \Gamma(n, \theta)$, and its pdf is

$$f(x) = \begin{cases} \frac{x^{n-1} e^{-\frac{x}{\theta}}}{\Gamma(n)\theta^n} & x > 0 \wedge \theta, n > 0 \\ 0 & \text{otherwise} \end{cases}$$

where $\Gamma(n) = \int_0^\infty e^{-y} y^{n-1} dy = (n-1)!$, where the last equality is true when n is an integer.

Note

The Gamma Distribution is usually used for when we are looking for the probability of the occurrence of the n -th event in the desired waiting time.

2.4 Functions of Random Variables

CONSIDER the rv X with pdf/pmf f and cdf F . Given $Y = h(X)$ where h is some real-valued function, we are interested in finding the pdf/pmf of Y .

The following are some possible scenarios:

1. X and Y are both discrete;
2. X is continuous and Y is discrete;
3. X and Y are both continuous

We may also define $Y = h(X)$ for a continuous rv X such that Y is **neither discrete nor continuous** (e.g. discrete for some values of X while continuous for others).

2.4.1 Discrete X and Discrete Y

If X and $Y = h(X)$ are both discrete, we can derive $P(Y = y)$ by mapping values in Y onto their corresponding value through h , i.e.


$$P(Y = y) = \sum_{\{x:h(x)=y\}} P(X = x)$$

Exercise 2.4.1

Let X have the following probability function:

$$f_X(x) = \begin{cases} \frac{e^{-1}}{x!} & x = 0, 1, 2, \dots \\ 0 & \text{otherwise} \end{cases}$$

Find the pmf of $Y = (X - 1)^2$.

 Solution

Note that since

$$\text{Dom } X = \{0, 1, 2, 3, 4, \dots\},$$

we have that

$$\text{Dom } Y = \{1, 0, 1, 4, 9, \dots\}.$$

With that, note that

$$P(Y = 0) = P(X = 1) = \frac{e^{-1}}{1!}$$

$$\begin{aligned} P(Y = 1) &= P(X = 0 \text{ or } 2) = P(X = 0) + P(X = 2) \\ &= \frac{e^{-1}}{0!} + \frac{e^{-1}}{2!} = e^{-1} \left(1 + \frac{1}{2} \right) = \frac{3}{2}e^{-1} \end{aligned}$$

$$P(Y = 4) = P(X = 3) = \frac{e^{-1}}{3!}$$

$$P(Y = 9) = P(X = 4) = \frac{e^{-1}}{4!}.$$

Therefore, the pmf of $Y = (X - 1)^2$ is

$$P(Y = y) = \begin{cases} e^{-1} & y = 0 \\ \frac{3}{2}e^{-1} & y = 1 \\ \frac{e^{-1}}{(1+\sqrt{y})!} & y = 4, 9, 16, \dots \\ 0 & \text{otherwise} \end{cases}$$

2.4.2 Continuous X and Discrete Y

If X is continuous and Y is discrete, we can use the method that we have used in the previous subsection, and replace Σ by the integral sign \int , i.e. define $A := \{x : h(x) = y\}$ such that we have

$$P(Y = y) = \int_A f(x) dx$$

Example 2.4.1 (Example 2.9)

Suppose X is a random variable with the following probability function

$$f_X(x) = \begin{cases} 2e^{2x} & x > 0 \\ 0 & \text{otherwise} \end{cases}.$$

Suppose $Y = h(X)$ is defined as follows:

$$Y = \begin{cases} 1 & X < 1 \\ 2 & 1 \leq X \leq 2 \\ 3 & X > 2 \end{cases}$$

Find the probability function of Y .

Solution

Note that $X \sim \text{Exp}(\frac{1}{2})$. So it is clear that X is a crv and since $Y = 1, 2,$

or 3, we have that Y is discrete. Now

$$\begin{aligned} P(Y = 1) &= P(X < 1) = \int_0^1 2e^{-2x} dx \\ &= -e^{-2x} \Big|_0^1 = 1 - e^{-2} \\ P(Y = 2) &= P(1 \leq X \leq 2) = \int_1^2 2e^{-2x} dx \\ &= -e^{-2x} \Big|_1^2 = e^{-2} - e^{-4} \\ P(Y = 3) &= P(X > 2) = \int_2^\infty 2e^{-2x} dx \\ &= -e^{-2x} \Big|_2^\infty = e^{-4} \end{aligned}$$

Thus the pmf is

$$P(Y = y) = \begin{cases} 1 - e^{-2} & Y = 1 \\ e^{-2} - e^{-4} & Y = 2 \\ e^{-4} & Y = 3 \end{cases}$$

2.4.3 Continuous X and Continuous Y

If X and $Y = h(X)$ are both continuous, start with the definition of the cdf of Y , i.e.

$$F_Y(y) = P(Y \leq y) = P(h(X) \leq y)$$


solve the inequality for X , and then obtain the cdf of Y . We will then only need to differentiate the cdf wrt y to get the pdf that we desire.

Example 2.4.2 (Example 2.10)

Let X have the following pdf:

$$f_X(x) = \begin{cases} 2e^{-2x} & x \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

Find the pdf of $Y = \sqrt{X}$.


 Solution

We have that the range of values where $f_Y(y) \leq 0$ is $y \geq 0$. Now

$$\begin{aligned} F_Y(y) &= P(Y \leq y) = P(\sqrt{X} \leq y) = P(X \leq y^2) \\ &= \int_0^{y^2} 2e^{-2x} dx \\ &= -e^{-2x} \Big|_0^{y^2} = 1 - e^{-2y^2} \end{aligned}$$

Therefore, the pdf of Y is

$$f_Y(y) = \begin{cases} \frac{d}{dy} 1 - e^{-2y^2} = 4ye^{-2y^2} & y \leq 0 \\ 0 & \text{otherwise} \end{cases}.$$

2.4.4 A Formula for the Continuous Case** Theorem 8 (One-to-One Transformation of a Random Variable)**

Suppose X is a continuous random variable with pdf f_X and support set $A = \{x : f_X(x) > 0\}$ and $Y = h(X)$ where h is a real-valued function. Let f_Y be the pdf of the rv Y and let $B = \{y : f_Y(y) > 0\}$. If h is a one-to-one function from A to B and if h' is continuous, then

$$f_Y(y) = f(h^{-1}(y)) \cdot \left| \frac{d}{dy} h^{-1}(y) \right|, \quad y \in B$$

 Proof

Note that since h is one-to-one, it is monotonous. Suppose h is increasing. Then h^{-1} is also an increasing function. Note that the cdf of Y is

$$F_Y(y) = P(Y \leq y) = P(X \leq h^{-1}(y)) = F_X(h^{-1}(y)).$$

Then the cdf of Y is

$$f_Y(y) = \frac{d}{dy} F_X(h^{-1}(y)) = f_X(h^{-1}(y)) \cdot \frac{d}{dy} h^{-1}(y)$$

3 Lecture 3 May 08th 2018

3.1 Functions of Random Variables (Continued)

3.1.1 Special Cases

Example 3.1.1

Recall Example 2.4.1. Suppose X is a rv with the following probability function

$$f_X(x) = \begin{cases} 2e^{-2x} & x > 0 \\ 0 & \text{otherwise} \end{cases}.$$

Define $Y = h(X)$ as follows:

$$Y = \begin{cases} 1 & X < 1 \\ X & 1 \leq X \leq 2 \\ 3 & X > 2 \end{cases}$$

Find the cdf of Y .

Solution

Solution is given differently in the 2 sections. I am not happy with either solutions because some things don't add up. My opinion is that the definition of Y is badly given, along with a badly phrased question. As a result, there are more ways than one to interpret an already confusing information, and thus we have ourselves one hell of a mess.

3.2 Probability Integral Transformation

Theorem 9 (Probability Integral Transformation)

If X is a continuous rv with cdf F , then $Y = F(X) \sim \text{Unif}(0, 1)$.
 $Y = F(X)$ is called the **probability integral transformation**.

Note

The distribution of $Y = F(X)$ can be proven.

Proof

Let X be a continuous rv and $Y = F(X)$. Since $F(X)$ is one-to-one and increasing (i.e. monotonous), there exists $F^{-1}(Y)$ that is a real-valued and increasing function. Then

$$\begin{aligned} F_Y(y) &= P(Y \leq y) = P(F_X(X) \leq y) = P(X \leq F^{-1}(y)) \\ &= F(F^{-1}(y)) = y \end{aligned}$$

Note that $F_Y(y) = y$ is the cdf of a $\text{Unif}(0, 1)$ rv, i.e. the **standard uniform random variable**. Thus $Y \sim \text{Unif}(0, 1)$.

Note

This theorem essentially states that any rv from a continuous distribution can be transformed into a standard uniform distribution.

Example 3.2.1 (Example 2.11)

Suppose $X \sim \text{Exp}(01)$. We know that $F_X(x) = 1 - e^{-10x}$ for all $x \in \mathbb{R}_+$. By Probability Integral Transformation, we have that $Y = F_X(X) = 1 - e^{-10X} \sim \text{Unif}(0, 1)$.

Note that the converse of Probability Integral Transformation is true:

▣ Theorem 10 (Converse of Probability Integral Transformation)

Suppose X is a continuous rv with cdf F such that F^{-1} exists. If $U \sim \text{Unif}(0,1)$, we have that $Y = F^{-1}(U) \sim X$.

✎ Proof

Note that

$$\begin{aligned} F_Y(y) &= P(Y \leq y) = P(F^{-1}(U) \leq y) \\ &= P(U \leq F_X(y)) = F_X(y). \end{aligned}$$

□

Example 3.2.2 (Example 2.12)

Suppose $X \sim \text{Unif}(0,1)$. Find a transformation T such that $T(X) \sim \exp(\theta)$.

✎ Solution

Let $Y = T(X) \sim \text{Exp}(\theta)$. Note that

$$F_Y(y) = 1 - e^{-\frac{y}{\theta}}, \quad y > 0$$

Observe that since

$$x = 1 - e^{-\frac{y}{\theta}} \implies y = -\theta \ln(1 - x)$$

we have that

$$F_Y^{-1}(X) = -\theta \ln(1 - X).$$

By **▣** Theorem 10, we have that $T = F_Y^{-1}$.

3.3 Location-Scale Families

When we look into methods for constructing confidence intervals for an unknown parameter θ . If the parameter θ is either a *scale parameter* or *location parameter*, then a confidence interval is easier to construct.

▣ Definition 19 (Location Parameter and Family)

Suppose X is a continuous rv with pdf $f(x; \mu)$, where μ is a parameter

of the distribution of X . Let $F_0(x) = F_X(x; \mu = 0)$, where F_X is the cdf of X , and $f_0(x) = f(x; \mu = 0)$. The parameter μ is called a **location parameter** of the distribution if

$$F_X(x; \mu) = F_0(x - \mu), \quad \mu \in \mathbb{R}$$

or equivalently,

$$f(x; \mu) = f_0(x - \mu), \quad \mu \in \mathbb{R}.$$

We say that F belongs to a **location family** of distributions.

Definition 20 (Scale Parameter and Family)

Suppose X is a continuous rv with pdf $f(x; \theta)$, where θ is a parameter of the distribution of X . Let $F_1(x) = F_X(x; \theta = 1)$, where F_X is the cdf of X , and $f_1(x) = f(x; \theta = 1)$. The parameter θ is called a **scale parameter** of the distribution if

$$F_X(x; \theta) = F_1\left(\frac{x}{\theta}\right), \quad \theta > 0$$

or equivalently,

$$f(x; \theta) = \frac{1}{\theta} f_0\left(\frac{x}{\theta}\right), \quad \theta > 0.$$

We say that F belongs to a **scale family** of distributions.

Definition 21 (Location-Scale Family)

Suppose X is an rv with cdf $F(x; \mu, \sigma)$ where $\mu \in \mathbb{R}$ and $\sigma > 0$ are the parameters of the distribution. Let $Y = \frac{X - \mu}{\sigma}$. If the distribution of Y does not depend on μ and/or σ , then F is said to belong to a **location-scale family** of distributions, with **location parameter** μ and **scale parameter** σ . In other words, F belongs to a location-scale family of distributions if

$$F(x; \mu, \sigma) = F_0\left(\frac{x - \mu}{\sigma}\right),$$

where $F_0(x) = F(x; \mu = 0, \sigma = 1)$, or equivalently,

$$f(x; \mu, \sigma) = \frac{1}{\sigma} f_0\left(\frac{x - \mu}{\sigma}\right),$$

where $f_0(x) = f(x; \mu = 0, \sigma = 1)$.

Example 3.3.1 (Example 2.13)

Consider $X \sim G(\mu, \sigma)$. Show that F_X belongs to a location-scale family of distributions.

We know that if $\mu = 0$ and $\sigma = 1$, then $Y = \frac{X-\mu}{\sigma} \sim G(0,1)$, and we know that $G(0,1)$ has no dependence on unknowns μ and σ . Therefore, F_X belongs to the location-scale family of distributions, with location parameter μ and scale parameter σ .

Another solution is to show that one of the equations in the definition is fulfilled. Observe that

$$f_x(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

So if we set $\mu = 0$ and $\sigma = 1$ to get f_0 , we have that

$$f_0(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}.$$

Now, note that

$$f(x) = \frac{1}{\sigma} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2} \left(\frac{x-\mu}{\sigma} \right)^2}.$$

Let $y = \frac{x-\mu}{\sigma}$, and we have ourselves

$$f(x) = \frac{1}{\sigma} \frac{1}{\sqrt{2\pi}} e^{-\frac{y^2}{2}} = \frac{1}{\sigma} f_0\left(\frac{x-\mu}{\sigma}\right)$$

Example 3.3.2 (Example 2.14)

Consider $X \in G(\mu, 2)$ where $\mu = E(X)$. Show that μ is a location parameter.

We can use a similar approach as before and define $Y = X - \mu$ which follows $G(0,2)$. It is clear that we then have that F_X , the cdf of X , belongs to a location family of distributions.

Example 3.3.3 (Example 2.15)

Consider $X \sim \text{Exp}(\theta)$. Show that F_X belongs to a scale family of distributions and find the scale parameter.

Note that

$$f(x) = \begin{cases} \frac{1}{\theta} e^{-\frac{x}{\theta}} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$

Let $Y = \frac{X}{\theta}$. Then

$$\begin{aligned} F_Y(y) &= P(Y \leq y) = P\left(\frac{X}{\theta} \leq y\right) \\ &= P(X \leq \theta y) = \int_0^{\theta y} \frac{1}{\theta} e^{-\frac{x}{\theta}} dx \\ &= -e^{-\frac{x}{\theta}} \Big|_0^{\theta y} = 1 - e^{-y} \end{aligned}$$

and we have

$$f_Y(y) = \begin{cases} e^{-y} & y > 0 \\ 0 & \text{otherwise} \end{cases}$$

Note that if we set $\sigma = 1$ to get f_1 , we have

$$f_1(x) = \begin{cases} e^{-x} & x > 0 \\ 0 & \text{otherwise} \end{cases}.$$

Therefore, F_X belongs to a scale family of distributions.

3.4 Expectations

3.4.1 Expectations

Definition 22 (Expectation of A Discrete RV)

If X is a discrete rv with pmf f and support set A , then the **expectation** of X , or the **expected value** of X is defined by

$$E(X) = \sum_{x \in A} xf(x) \tag{3.1}$$

provided that the sum converges absolutely, i.e.

$$E(|X|) = \sum_{x \in A} |x|f(x) < \infty.$$

If $E(|X|)$ does not converge, then we say that $E(X)$ does not exist.

Definition 23 (Expectation of A Continuous RV)

If X is a continuous rv with pdf f and support set A , then the **expectation** of X , or the **expected value** of X is defined by

$$E(X) = \int_{x \in A} xf(x) \quad (3.2)$$

provided that the integral converges absolutely, i.e.

$$E(|X|) = \int_{x \in A} |x| f(x) < \infty.$$

If $E(|X|)$ does not converge, then we say that $E(X)$ does not exist.

Example 3.4.1 (Example 2.16)

Suppose $X \sim \text{Poi}(\lambda)$. Calculate $E(X)$.

Solution

Note

$$f(x) = \begin{cases} \frac{e^{-\lambda} \lambda^x}{x!} & x = 0, 1, 2, \dots \\ 0 & \text{otherwise} \end{cases}.$$

Then

$$\begin{aligned} E(X) &= \sum_{x=0}^{\infty} x \frac{e^{-\lambda} \lambda^x}{x!} \\ &= 0 + \sum_{x=1}^{\infty} \frac{e^{-\lambda} \lambda^x}{(x-1)!} \\ &= e^{-\lambda} \lambda \sum_{x=1}^{\infty} \frac{\lambda^{x-1}}{(x-1)!} \\ &= e^{-\lambda} \lambda e^{\lambda} = \lambda \end{aligned}$$

Example 3.4.2 (Example 2.18)

Suppose X is an rv with

$$f(x) = \begin{cases} \frac{1}{x^2} & 1 < x < \infty \\ 0 & \text{otherwise} \end{cases}.$$

Calculate $E(X)$.

 **Solution**

Observe that $x \cdot \frac{1}{x^2} = \frac{1}{x}$ and the antiderivative of $\frac{1}{x}$ is $\ln x$, which would need to be evaluated at $\ln \infty$. Thus, we should instead immediately check if the integral converges absolutely.

$$\begin{aligned} E(|X|) &= \int_1^{\infty} |x| \frac{1}{x^2} dx \\ &= \int_1^{\infty} |x| \frac{1}{|x||x|} dx \\ &= \int_1^{\infty} \frac{1}{|x|} dx \\ &= \int_1^{\infty} \frac{1}{x} dx, \end{aligned}$$

and we notice that the integral would not converge. Therefore, $E(X)$ does not exist.

4 Lecture 4 May 10th 2018

4.1 Expectations (Continued)

4.1.1 Expectations (Continued)

Theorem 11 (Expectation from the cdf)

Suppose X is a non-negative continuous rv with cdf F , and $E(X) < \infty$.

Then

$$E(X) = \int_0^{\infty} [1 - F(x)] dx = \int_0^{\infty} P(X \geq x) dx \quad (4.1)$$

If X is a discrete rv with cdf F , and $E(X) < \infty$, then

$$E(X) = \sum_{x=0}^{\infty} [1 - F(x)] = \sum_{x=0}^{\infty} P(X \geq x) \quad (4.2)$$

Proof

Note that for a continuous rv X , we have

$$1 - F(x) = P(X \geq x) = \int_x^{\infty} f(t) dt$$

Therefore,


$$\int_0^{\infty} [1 - F(x)] dx = \int_0^{\infty} \int_x^{\infty} f(t) dt dx.$$

Since $1 - F(x)$ is a finite value, so is $\int_0^{\infty} f(t) dt$, and thus we can apply **Fubini's Theorem**¹:

$$\int_0^{\infty} [1 - F(x)] dx = \int_0^{\infty} \int_x^{\infty} f(t) dt dx = \int_0^{\infty} \int_0^t f(t) dx dt$$

Note that the limits of the integral utilizes the following figure:

¹ Condition for Fubini's Theorem to hold is that the integrand of the double integral must be absolutely convergent. See Wikipedia.



With that, note that

$$\int_0^t f(t) dx = xf(t) \Big|_0^t = tf(t)$$

Since t is just a dummy variable, we can indeed let $t = x$, and thus we have

$$\int_0^\infty [1 - F(x)] dx = \int_0^\infty xf(x) dx = E(X)$$

as required.

Work on the discrete case as an exercise.


Exercise 4.1.1

For a non-negative discrete rv X with cdf F and $E(X) < \infty$, prove that

$$E(X) = \sum_{x=0}^{\infty} [1 - F(x)]$$

□

Example 4.1.1 (Example 2.20)

Suppose $X \sim \text{Exp}(\theta)$. Use  Theorem 11 to calculate $E(X)$.

Solution

Note that X is a non-negative rv. The cdf of $X \sim \text{Exp}(\theta)$ is

$$F_X(x) = 1 - e^{-\frac{x}{\theta}}.$$

Then

$$\begin{aligned} E(X) &= \int_0^\infty 1 - F_X(x) dx = \int_0^\infty e^{-\frac{x}{\theta}} dx \\ &= -\theta e^{-\frac{x}{\theta}} \Big|_0^\infty = \theta \end{aligned}$$

□

Theorem 12 (Expected Value of a Function of X)

Suppose $h(x)$ is a real-valued function.

If X is a discrete rv with pmf f and support set A , then

$$E[h(x)] = \sum_{x \in A} h(x)f(x) \quad (4.3)$$

provided that the sum converges absolutely.

If X is a continuous rv with pdf f , then

$$E[h(x)] = \int_{-\infty}^{\infty} h(x)f(x) dx \quad (4.4)$$

provided that the integral converges absolutely.

The proof is, unfortunately, not trivial. One would have to look into Lebesgue integrals (or at the very least, Riemann-Stieltjes integrals) in order to prove this statement. This “theorem” is also called **The Law of the Unconscious Statistician** [Reference - Wikipedia]. An idea of the proof is given on Math SE.

Example 4.1.2

Suppose $X \sim \text{Unif}(0, \theta)$. Calculate $E(X^2)$.

Solution

$$E(X^2) = \int_0^{\theta} \frac{x^2}{\theta} dx = \frac{1}{\theta} \frac{x^3}{3} \Big|_{x=0}^{\theta} = \frac{\theta^2}{3}$$

Exercise 4.1.2

Find the pdf of $Y = X^2$ and find $E(Y)$ by evaluating $\int_{-\infty}^{\infty} y f_Y(y) dy$


Theorem 13 (Linearity of Expectation)

Suppose X is an rv with pf f . Let $a_i, b_i \in \mathbb{R}$, for $i = 1, \dots, n$, be constants, and $g_i(x)$, for $i = 1, \dots, n$, are real-valued functions. Then

$$E \left[\sum_{i=1}^n (a_i g_i(X) + b_i) \right] = \sum_{i=1}^n (a_i E[g_i(X)] + b_i) \quad (4.5)$$

provided that $E[g_i(X)] < \infty$ for $i = 1, \dots, n$.

This theorem essentially states that the expectation is a linear operator.

 **Proof**

Suppose X is a discrete rv with support set A . Then

$$\begin{aligned}
 E\left[\sum_{i=1}^n (a_i g_i(X) + b_i)\right] &= \sum_{x \in A} \left[\sum_{i=1}^n (a_i g_i(x) + b_i) \right] f(x) \quad \because \text{Theorem 12} \\
 &= \sum_{x \in A} \sum_{i=1}^n [a_i g_i(x) f(x) + b_i f(x)] \\
 &= \sum_{i=1}^n \sum_{x \in A} [a_i g_i(x) f(x) + b_i f(x)] \quad (*) \\
 &= \sum_{i=1}^n \left[a_i \sum_{x \in A} g_i(x) f(x) + b_i \sum_{x \in A} f(x) \right] \\
 &= \sum_{i=1}^n (a_i E[g_i(X)] + b_i)
 \end{aligned}$$

where note that $(*)$ is valid because a_i, b_i are constants, and $g_i(x), f(x)$ are finite real-valued functions.


“ Note

In general, $E(g(X)) \neq g(E(X))$ unless if g is a linear function. For example, for $a, b \in \mathbb{R}$, we have

$$E(aX + b) = aE(X) + b$$


4.1.2 Moments and Variance

Since these concepts were introduced in STAT230 and were given little treatment in the lecture, we shall only cover over them briefly.

 **Definition 24 (Variance)**

The expectation of the squared deviation of an rv from its mean is called the **variance**, i.e. for an rv X with mean $\mu = E(X)$,

$$\sigma^2 = \text{Var}(X) = E[(X - \mu)^2] = E(X^2) - E(X)^2$$

 **Definition 25 (Moments)**

Let X be an rv with mean μ .

The k^{th} **moment about the origin** is defined as:

$$E(X^k)$$

The k^{th} **moment about the mean** is defined as:

$$E[(X - \mu)^k]$$

The k^{th} **factorial moment** is defined as:

$$E[X^{(k)}] = E[X(X-1)\dots(X-k+1)] = E\left[\frac{X!}{(X-k)!}\right]$$

Theorem 14 (Variance of a Linear Function)

Suppose X is an rv with pf f and $a, b \in \mathbb{R}$. Then

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Proof

Observe that

$$\begin{aligned} \text{Var}(aX + b) &= E[(aX + b)^2] - E(aX + b)^2 \\ &= E[a^2X^2 + 2abX + b^2] - (aE(X) + b)^2 \\ &= a^2E(X^2) + 2abE(X) + b^2 - (a^2E(X)^2 + 2abE(X) + b^2) \\ &= a^2E(X^2) - a^2E(X)^2 = a^2 \text{Var}(X) \end{aligned}$$

□

Example 4.1.3 (Example 2.22 (course notes - 2.6.10 (1)))

If $X \sim \text{Poi}(\theta)$, then $E[X^{(k)}] = \theta^k$ for $k = 1, 2, \dots$

 **Solution**

Note

$$f_X(x) = \begin{cases} \frac{e^{-\theta}\theta^x}{x!} & x = 0, 1, 2, \dots \\ 0 & \text{otherwise} \end{cases}$$

So

$$\begin{aligned} E[X^{(k)}] &= E(X(X-1)(X-2)\dots(X-k+1)) \\ &= \sum_{x=0}^{\infty} x(x-1)(x-2)\dots(x-k+1) \frac{e^{-\theta}\theta^x}{x!} \\ &= 0 + \sum_{x=k}^{\infty} x(x-1)(x-2)\dots(x-k+1) \frac{e^{-\theta}\theta^x}{x!} \quad (*) \\ &= \sum_{x=k}^{\infty} \frac{x!}{(x-k)!} \frac{e^{-\theta}\theta^x}{x!} \quad \because x(x-1)\dots(x-k+1) = \frac{x!}{(x-k)!} \\ &= e^{-\theta}\theta^k \sum_{x=k}^{\infty} \frac{\theta^{x-k}}{(x-k)!} \\ &= e^{-\theta}\theta^k \sum_{y=0}^{\infty} \frac{\theta^y}{y!} \quad \text{let } y = x - k \\ &= e^{-\theta}\theta^k e^{\theta} = \theta^k \end{aligned}$$

where for (*) we have that $\sum_{x=0}^{k-1} x(x-1)\dots(x-k+1)A = 0$ for any $A \in \mathbb{R}$.

Note that it is not necessarily true that

$$x(x-1)\dots(x-k+1) = \frac{x!}{(x-k)!}$$

for $0 \leq x \leq k-1$. And so we can only say that the equality is true for $x \geq k$, and hence we have the approach that we use in (*).

4.2 Inequalities**4.2.1 Markov/Chebyshev Style Inequalities** **Theorem 15 (Markov's Inequality)**

If X is a non-negative rv and $a > 0$, then the probability that X is no less than a is no greater than the expectation of X divided by a , i.e.

$$P(X \geq a) \leq \frac{E(X)}{a} \quad (4.6)$$

 **Proof**

We shall prove for the discrete case. Suppose X is a non-negative discrete rv with pf f . Let $A \subset S$, where S is the sample space, such

that

$$A = \{w \in S : X(w) \geq a\}.$$

$$\begin{aligned} E(X) &= \sum_{x \in S} xf(x) \\ &= \sum_{x \in A} xf(x) + \sum_{x \notin A} xf(x) \\ &\geq \sum_{x \in A} xf(x) \quad \because \sum_{x \notin A} xf(x) \geq 0 \\ &\geq \sum_{x \in A} af(x) \\ &= a \sum_{x \in A} f(x) = a \cdot P(A) \\ &= a \cdot P(\{w \in S : X(w) \geq a\}) = aP(X \geq a). \end{aligned}$$

Exercise 4.2.1

Prove Markov's Inequality for a continuous rv.

□

📖 Theorem 16 (Markov's Inequality 2)

If X is a non-negative rv and $a, k > 0$, then the probability that X is no less than a is no greater than the expectation of X divided by a , i.e.

$$P(|X| \geq a) \leq \frac{E(|X|^k)}{a^k} \tag{4.7}$$

✎ Proof

We shall, again, prove for the discrete case. Suppose X is a non-negative discrete rv with pf f . $A := \{w \in S : |X(w)| \geq a\} \subseteq S$.

Then

$$\begin{aligned} E(|X|^k) &= \sum_{x \in S} |x|^k f(x) \\ &= \sum_{x \in A} |x|^k f(x) + \sum_{x \notin A} |x|^k f(x) \\ &\geq \sum_{x \in A} |x|^k f(x) \geq \sum_{x \in A} af(x) \\ &= a^k P(A) = a^k P(|X| \geq a). \end{aligned}$$


Question: Can we write

$$P(\{w \in S : |X(w)| \geq a\}) = P(|X| \geq a)?$$

Exercise 4.2.2

Prove for the continuous case.


□

 Theorem 17 (Chebyshev's Inequality)

Suppose X is an rv with finite mean μ and finite variance σ^2 . Then for any $k > 0$,

$$P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2} \quad (4.8)$$

 Proof

By  Theorem 16,


$$P(|X - \mu| \geq k\sigma) \leq \frac{E(|X - \mu|^2)}{(k\sigma)^2} = \frac{1}{k^2}$$

since $E(|X - \mu|^2) = \text{Var}(X) = \sigma^2$. □

Example 4.2.1 (Example 2.23)

A post office handles, on average, 10000 letters a day. What can be said about the probability that it will handle at least 15000 letters tomorrow?

 Solution

X := number of letters handled in a day. Note that by its definition, X is a non-negative discrete rv. Then, using  Theorem 15, since $E(X) = 10000$

$$P(X \geq 15000) \leq \frac{10000}{15000} = \frac{2}{3}.$$


Thus, we know that there is less than two-third of chance that the post office will handle more than 15000 tomorrow.

5 Lecture 5 May 15th 2018

5.1 Inequalities (Continued)

5.1.1 Markov/Chebyshev Style Inequalities (Continued)

Example 5.1.1 (Example 2.24)

A post office handles 10000 letters per day with a variance of 2000 letters. What can be said about the probability that this post office handles between 8000 and 12000 letters tomorrow? What about the probability that more than 15000 letters come in (use  Theorem 17)?

1. Probability that this post office handles between 8000 and 12000 letters tomorrow:


$$\begin{aligned} & P(8000 < X < 12000) \\ &= P(-2000 < X - 10000 < 2000) \\ &= P(|X - 10000| < 2000) = 1 - P(|X - 10000| \geq 2000) \\ &\geq 1 - \frac{1}{(\frac{2000}{\sigma})^2} \quad \because 1 \text{ Theorem 17} \wedge k = \frac{2000}{\sigma} = \sqrt{2000} \\ &= \frac{1999}{2000} \end{aligned}$$

2. Probability that more than 15000 letters come in:

$$\begin{aligned} P(X > 15000) &= P(X - 10000 > 15000 - 10000) \\ &= P(X - 10000 > 5000) \\ &\leq P(X - 10000 > 5000) + P(X - 10000 < 5000) \\ &\leq P(|X - 10000| > 5000) \\ &\leq \frac{1}{\left(\frac{5000}{\sqrt{2000}}\right)^2} = \frac{2000}{5000^2} \end{aligned}$$

5.2 Moment Generating Function

Moment generating functions are important because they uniquely define the distribution of an rv.

 Definition 26 (Moment Generating Function)

If X is an rv, then $M_X(t) = E(e^{tx})$ is called the **moment generating function** (mgf) of X provided this expectation exists for all $t \in (-h, h)$ for some $h > 0$.

“ Note

When determining the mgf of an rv, the values of t for which the expectation exists must always be stated. The range of t where the expectation is defined is “essentially” the **radius of convergence**.

Exercise 5.2.1 (Example 2.25 (2.9.2 (1) of the course notes))

Find the mgf of $X \sim \Gamma(\alpha, \beta)$. Make sure you specify the domain on which the mgf is defined.

 Solution

Note that the pdf of the Gamma distribution is:

$$f(x) = \begin{cases} \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-\frac{x}{\beta}} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$

Therefore

$$\begin{aligned}
 M_X(t) &= E(e^{tx}) = \int_0^\infty e^{tx} \frac{1}{\beta^\alpha} x^{\alpha-1} e^{-\frac{x}{\beta}} dx \\
 &= \frac{1}{\beta^\alpha} \int_0^\infty \frac{1}{\Gamma(\alpha)} x^\alpha e^{-x\left(\frac{1}{\beta}-t\right)} dx \\
 &= \frac{\left(\frac{\beta}{1-t\beta}\right)^\alpha}{\beta^\alpha} \int_0^\infty \underbrace{\frac{1}{\left(\frac{\beta}{1-t\beta}\right)^\alpha} x^{\alpha-1} e^{-\frac{x}{\frac{\beta}{1-t\beta}}}}_{\text{sum over all values for pdf of } \Gamma(\alpha, \frac{\beta}{1-t\beta}=1)} dx \quad \text{for } \frac{1}{\beta} - t > 0 \\
 &= (1-t\beta)^{-\alpha} \quad \text{for } t < \frac{1}{\beta}
 \end{aligned}$$

Definition 27 (Indicator Function)

The function $\mathbb{1}_A$ is called the **indicator function** of the set A , i.e.

$$\mathbb{1}_A = \begin{cases} 1 & \text{if } A \text{ occurs} \\ 0 & \text{if } A^C \text{ occurs} \end{cases} \quad (5.1)$$

Example 5.2.1

The pdf

$$f(x) = \begin{cases} \frac{1}{\theta} & 0 \leq x \leq \theta \\ 0 & \text{otherwise} \end{cases}$$

can be represented as

$$f(x) = \frac{1}{\theta} \mathbb{1}_{\{0 \leq x \leq \theta\}}$$

Example 5.2.2 (Example 2.26)

Find the mgf of $X \sim \text{Poi}(\lambda)$. Make sure you specify the domain on which the mgf is defined.

Solution

Note that the pmf of X is

$$f_X(x) = \frac{e^{-\lambda} \lambda^x}{x!} \mathbb{1}_{\{0,1,2,\dots\}}$$

The mgf is thus

$$\begin{aligned} M_X(t) &= E(e^{tX}) = \sum_{x=0}^{\infty} e^{tx} \frac{e^{-\lambda} \lambda^x}{x!} \\ &= e^{-\lambda} \sum_{x=0}^{\infty} \frac{(e^t \lambda)^x}{x!} = e^{-\lambda} e^{e^t \lambda} \\ &= e^{\lambda(e^t - 1)} \quad \forall t \in \mathbb{R} \end{aligned}$$

♦ Proposition 18 (Properties of the MGF)

Suppose X is an rv. Then

1. $M_X(0) = 1$
2. Suppose the derivatives $M_X^{(k)}(t)$, for $k = 1, 2, \dots$, exists for $t \in (-h, h)$ for some $h > 0$, then the **Maclaurin Series**¹ of $M_X(t)$ is

$$M_X(t) = \sum_{k=0}^{\infty} \frac{M_X^{(k)}(t) \Big|_{t=0}}{k!} t^k$$

¹ The Maclaurin series is the Taylor expansion around 0.

3. If the mgf exists, then the k^{th} moment of X is:

$$E(X^k) = \frac{d^k M_X(t)}{dt^k} \Big|_{t=0}$$

4. Putting 2 and 3 together, we have

$$M_X(t) = \sum_{k=0}^{\infty} \frac{E(X^k)}{k!} t^k$$

The final item shows why $M_X(t)$ is called the **moment generating function**.

Proof

1. $M_X(t) \Big|_{t=0} = E(e^{tX}) \Big|_{t=0} = E(e^0) = 1$
2. This is simply a result of using the Maclaurin series.

3. Note that

$$\begin{aligned} E(e^{tX}) &= E \left[1 + tX + \frac{1}{2}(tX)^2 + \frac{1}{3!}(tX)^3 + \dots \right] \\ &= 1 + tE(X) + \frac{t^2}{2}E(X^2) + \frac{t^3}{3!}E(X^3) + \dots \end{aligned}$$

So

$$\frac{d^k}{dt^k} E(e^{tX}) \Big|_{t=0} = \frac{k!}{k!} E(X^k) + \underbrace{\frac{k! \cdot t}{(k+1)!} E(X^{k+1}) + \dots}_{=0 \text{ when } t=0} \Big|_{t=0} = E(X^k)$$

□

Example 5.2.3 (Example 2.27)

A discrete random variable X has the pmf

$$f(x) = \left(\frac{1}{2}\right)^{x+1} \mathbb{1}_{\{0,1,2,\dots\}}$$

Derive the mgf of X and use it calculate its mean and variance.

$$\begin{aligned} M_X(t) &= \sum_{x=0}^{\infty} e^{tx} \left(\frac{1}{2}\right)^{x+1} \\ &= \frac{1}{2} \cdot \sum_{x=0}^{\infty} \left(\frac{e^t}{2}\right)^x \\ &= \frac{1}{2} \cdot \frac{1}{1 - \frac{e^t}{2}} \quad \text{for } \left|\frac{e^t}{2}\right| < 1 \text{ or } t < \ln 2 \\ &= \frac{1}{2 - e^t} \end{aligned}$$

To get the first two moments,

$$\begin{aligned} E(X) &= \frac{d}{dt} M_X(t) \Big|_{t=0} \\ &= \frac{e^t}{(2 - e^t)^2} \Big|_{t=0} = 1 \\ E(X^2) &= \frac{d^2}{dt^2} M_X(t) \Big|_{t=0} \\ &= \frac{e^t}{(2 - e^t)^2} + \frac{2e^t}{(2 - e^t)^3} \Big|_{t=0} \\ &= 1 + 2 = 3 \end{aligned}$$

Thus we have that the expected value and variance are

$$\begin{aligned} E(X) &= 1 \\ \text{Var}(X) &= E(X^2) - E(X)^2 = 3 - 1 = 2 \end{aligned}$$

respectively.

5.2.1 MGF of a Linear Transformation

Theorem 19 (MGF of a Linear Transformation)

Suppose the rv X has an mgf $M_X(t)$ defined for $t \in (-h, h)$ for some $h > 0$. Let $Y = aX + b$, where $a, b \in \mathbb{R}$ and $a \neq 0$. Then the mgf of Y is

$$M_Y(t) = e^{bt} M_X(at), \quad |t| \leq \frac{h}{|a|}. \quad (5.2)$$

Proof

Observe that

$$M_Y(t) = E(e^{tY}) = E(e^{t(aX+b)}) = E(e^{atX} e^{tb}) = e^{bt} M_X(at).$$

The range of t is

$$|at| < h \stackrel{a \neq 0}{\iff} |t| < \frac{h}{|a|}$$


Example 5.2.4 (Example 2.28)

Consider $X \sim \text{Unif}(\theta_1, \theta_2)$. Find the mgf of $Y = 5X + 3$.

 **Solution**

Note that

$$\begin{aligned}
 M_X(t) &= \int_{\theta_1}^{\theta_2} \frac{e^{tx}}{\theta_2 - \theta_1} dx \\
 &= \begin{cases} \frac{e^{tx}}{t(\theta_2 - \theta_1)} \Big|_{\theta_1}^{\theta_2} & t \neq 0 \\ 1 & t = 0 \end{cases} \\
 &= \begin{cases} \frac{e^{t\theta_2} - e^{t\theta_1}}{t(\theta_2 - \theta_1)} & t \neq 0 \\ 1 & t = 0 \end{cases}
 \end{aligned}$$

Thus by  Theorem 19,

$$M_Y(t) = e^{3t} M_X(5t) = \begin{cases} e^{3t} \frac{e^{5t\theta_2} - e^{5t\theta_1}}{5t(\theta_2 - \theta_1)} & t \neq 0 \\ 1 & t = 0 \end{cases}$$

5.2.2 Uniqueness of the MGF

Theorem 20 (Uniqueness of the MGF)

Suppose the rv X has mgf $M_X(t)$ and the rv Y has mgf $M_Y(t)$. Suppose also that $M_X(t) = M_Y(t)$ for all $t \in (-h, h)$ for some $h > 0$. Then X and Y have the same distribution, that is, $\forall s \in \mathbb{R}$,

$$P(X \leq s) = F_X(s) = F_Y(s) = P(Y \leq s)$$

Proof

The proof of this theorem is not trivial. See [this comment on Math SE](#) for information. It appears that the 2nd bullet point points to a material that I might be able to understand. If I can find that material, and understand it, I may change this proof section to become my own notes.

Example 5.2.5 (Example 2.29)

Suppose $X \sim \text{Unif}(0, 1)$. Define $Y = -2 \log X$, and use the mgf method to show that $Y \sim \chi_2^2$.

(Hint: Find mgf of χ_2 and show that Y has the same mgf)

 Solution

Let $Z = \chi_2^2$. The pdf of Z is therefore

$$f_Z(z) = \frac{1}{2} e^{-\frac{z}{2}} \mathbb{1}_{\{z > 0\}}.$$

Then

$$\begin{aligned} M_Z(t) &= E(e^{tZ}) = \int_0^\infty e^{tz} \frac{1}{2} e^{-\frac{z}{2}} dz \\ &= \frac{1}{2} \int_0^\infty e^{(t-\frac{1}{2})z} dz \\ &= \begin{cases} \frac{1}{2} \frac{1}{t-\frac{1}{2}} e^{(t-\frac{1}{2})z} \Big|_{z=0}^\infty & t \neq \frac{1}{2} \\ \infty & t = \frac{1}{2} \end{cases} \\ &= \frac{1}{2t-1} \quad t \neq \frac{1}{2} \end{aligned}$$

6 Lecture 6 May 17th 2018

6.1 Joint Distributions

6.1.1 Introduction to Joint Distributions

“ Note (Motivation)

Most studies collect information for multiple variables per subject rather than just one variable. Because these variables may interfere/interact with each other and hence give us results that may not be fully reliant on a single variable, it is in our interest to study the interaction of these variables.

To start off with the basics, we will first look at the bivariate case of a joint distribution.

6.1.2 Joint and Marginal CDFs


📖 Definition 28 (Joint CDF)

Suppose X and Y are rvs defined on a sample space S . The **joint cdf** of X and Y is given by

$$\forall (x, y) \in \mathbb{R}^2 \quad F(x, y) = P(X \leq x, Y \leq y).$$

“ Note

- Depending on whether X and Y are both discrete or both continuous, we can derive the joint pmf or joint pdf of (X, Y) , respectively.

-  Definition 28 only concerns two variables (a bivariate case), but we can certainly extend the idea to a k -dimensional joint cdf for the rvs X_1, X_2, \dots, X_k as $\forall (x_1, x_2, \dots, x_k) \in \mathbb{R}^k$,

$$F(x_1, x_2, \dots, x_k) = P(X_1 \leq x_1, X_2 \leq x_2, \dots, X_k \leq x_k).$$

♦ Proposition 21 (Properties of Joint CDF)

Suppose X, Y are rvs, either both continuous or discrete, and has a joint cdf F . Then

1. F is non-decreasing in x for fixed y .
2. F is non-decreasing in y for fixed x .
3. $\lim_{x \rightarrow -\infty} F(x, y) = 0$ and $\lim_{y \rightarrow -\infty} F(x, y) = 0$.
4. $\lim_{(x, y) \rightarrow (-\infty, -\infty)} F(x, y) = 0$ and $\lim_{(x, y) \rightarrow (\infty, \infty)} F(x, y) = 1$

Proof

1. Suppose not, i.e. that we have instead that F is decreasing for x .
Then for $x_1 < x_2 \in \mathbb{R}$, we would have

$$\begin{aligned} F(x_1, y) &> F(x_2, y) \\ \implies P(X \leq x_1, Y \leq y) &> P(X \leq x_2, Y \leq y) \end{aligned}$$

In other words,

$$\begin{aligned} &P(\{(w, v) : (w, v) \in S, X(w) \leq x_1, Y(v) \leq y\}) \\ &> P(\{(w, v) : (w, v) \in S, X(w) \leq x_2, Y(v) \leq y\}) \end{aligned}$$

However, note that for fixed y , since $x_1 < x_2$, we must have that

$$\begin{aligned} &\{(w, v) \in S : X(w) \leq x_1, Y(v) \leq y\} \\ &\subseteq \{(w, v) \in S : X(w) \leq x_2, Y(v) \leq y\}. \end{aligned}$$

By  Proposition 1, we have that

$$\begin{aligned} &P(\{(w, v) : (w, v) \in S, X(w) \leq x_1, Y(v) \leq y\}) \\ &\leq P(\{(w, v) : (w, v) \in S, X(w) \leq x_2, Y(v) \leq y\}). \end{aligned}$$

This is clearly a contradiction.

2. The proof for this statement is similar to the above.
3. Note that

$$\begin{aligned}\lim_{x \rightarrow -\infty} F(x, y) &= \lim_{x \rightarrow -\infty} P(X \leq x, Y \leq y) \\ &= P(X \leq -\infty, Y \leq y) \\ &= P([X \leq -\infty] \cap [Y \leq y]) \\ &= P(\emptyset \cap [Y \leq y]) = P(\emptyset) = 0\end{aligned}$$

The proof for the case where $y \rightarrow -\infty$ is similar.

4. This is simply a consequence of 3.

“ Note

We say that F is a joint cdf if it satisfies all the conditions in [Proposition 2.1](#).¹¹ Many literature actually claims this, and it does look like it will be assumed so for this class.

Example 6.1.1 (Example 3.1)

Consider the following joint cdf of two rvs (X_1, X_2) :


$$F(x_1, x_2) = \begin{cases} 0 & x_1 < 0 \vee x_2 < 0 \\ 0.49 & 0 \leq x_1 < 1 \wedge 0 \leq x_2 < 1 \\ 0.7 & 0 \leq x_1 < 1 \wedge x_2 > 1 \\ 0.7 & x_1 \geq 1 \wedge 0 \leq x_2 < 1 \\ 1 & x_1 \geq 1 \wedge x_2 \geq 1 \end{cases}$$

Flipping an unfair coin with $P(\{H\}) = 0.3$ twice independently, we define for $i = 1, 2$


$$X_i = \begin{cases} 1 & \text{if the } i^{\text{th}} \text{ flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

The joint cdf of (X_1, X_2) is the given F above. Verify that under this experiment, F is indeed a cdf.

 **Solution**

Note that conditions 3 and 4 of  Proposition 21 are automatically satisfied by the definition of F .

incomplete example

 **Definition 29 (Marginal CDF)**

For the rvs X, Y with joint cdf F , the **marginal cdf** of X is

$$F_X(x) = P(X \leq x) = \lim_{y \rightarrow \infty} F(x, y) = F(x, \infty) \quad \forall x \in \mathbb{R}$$

and the **marginal cdf** of Y is

$$F_Y(y) = P(Y \leq y) = \lim_{x \rightarrow \infty} F(x, y) = F(\infty, y) \quad \forall y \in \mathbb{R}$$

Note that the marginal cdf is defined for both discrete and continuous cases.

Example 6.1.2


Based on Example 6.1.1, derive $F_{X_i}(x_i)$ for $i = 1, 2$.

 **Solution**

$$F_{X_1}(x_1) = \lim_{x_2 \rightarrow \infty} F(x_1, x_2) = \begin{cases} 0 & x_1 < 0 \\ 0.7 & 0 \leq x_1 < 1 \\ 1 & x_1 \geq 1 \end{cases}$$

The solution for $F_{X_2}(x_2)$ is similar.

6.1.3 Joint Discrete RVs

 **Definition 30 (Joint Discrete RV)**

Suppose X and Y are rvs defined on a sample space S . If S is discrete

then X and Y are discrete rvs. The **joint pmf** of X and Y is given by

$$\forall (x, y) \in \mathbb{R}^2 \quad f(x, y) = P(X = x, Y = y).$$

The set $A = \{(x, y) : f(x, y) > 0\}$ is called the **support set** of (X, Y) .

💧 Proposition 22 (Properties of Joint PMF)

Suppose X, Y are discrete rvs with joint pmf f and support set A . Then

- $\forall (x, y) \in \mathbb{R}^2 \quad f(x, y) \geq 0$

- $\sum_{(x,y) \in A} f(x, y) = 1$

- $\forall R \subset \mathbb{R}^2,$

$$P[(X, Y) \in R] = \sum_{(x,y) \in R} f(x, y)$$

The proof is analogous to the univariate case as seen in 💧 Proposition 6

Example 6.1.3 (Example 3.2)

Consider the following joint pmf where the numbers inside the table show $P(X = x, Y = y)$. Find c . Then, calculate $P(X + Y \leq 2)$.

	$x = -2$	$x = 0$	$x = 2$
$y = 0$	0.05	0.1	0.15
$y = 1$	0.07	0.11	c
$y = 2$	0.02	0.25	0.05

✎ Solution

Since the sum of all the probabilities must be 1, thus

$$c = 1 - 0.05 - 0.07 - 0.02 - \dots - 0.15 - 0.05 = 0.2.$$

Notice that the only cases where $X + Y > 2$ is when

- $X = 2, Y = 1$; and
- $X = 2, Y = 2$.

Thus

$$\begin{aligned} P(X + Y \leq 2) &= 1 - P(X = 2, Y = 1) - P(X = 2, Y = 2) \\ &= 1 - 0.2 - 0.05 = 0.75 \end{aligned}$$

Example 6.1.4 (Example 3.3)

A small college has 90 male and 30 female professors. An ad hoc committee of 5 is selected at random to write the vision and mission of the college. Let X and Y be the number of men and women in this committee, respectively. Derive the joint distribution of (X, Y) .

Solution

Observe that the support set of this distribution is

$$A = \{(x, y) : x + y = 5, x, y = 0, 1, 2, 3, 4, 5\}.$$

We have that the distribution is

$$P(X = x, Y = y) = \begin{cases} \frac{\binom{90}{x} \binom{30}{y}}{\binom{120}{5}} & x, y = 0, 1, 2, 3, 4, 5 \\ & x + y = 5 \\ 0 & \text{otherwise} \end{cases}$$

Definition 31 (Marginal Distribution - Discrete Case)

Suppose X and Y are discrete rvs with joint pf f . Then the *marginal pf* of X is

$$\forall x \in \mathbb{R}^2 \quad f_X(x) = P(X = x) = \sum_{y \in \mathbb{R}} f(x, y),$$

and the *marginal pf* of Y is

$$\forall y \in \mathbb{R}^2 \quad f_Y(y) = P(Y = Y) = \sum_{x \in \mathbb{R}} f(x, y).$$

Example 6.1.5 (Example 3.4)

Consider the joint pmf from Example 6.1.3. Find the marginal distri-

butions, i.e. marginal pmfs of X and Y .

	$x = -2$	$x = 0$	$x = 2$
$y = 0$	0.05	0.1	0.15
$y = 1$	0.07	0.11	0.2
$y = 2$	0.02	0.25	0.05

 **Solution**

Using the definition, we have that

$$f_X(x) = \sum_{y \in \mathbb{R}} f(x, y) = \begin{cases} 0.14 & x = -2 \\ 0.46 & x = 0 \\ 0.40 & x = 2 \end{cases}$$

and

$$f_Y(y) = \sum_{x \in \mathbb{R}} f(x, y) = \begin{cases} 0.3 & y = 0 \\ 0.38 & y = 1 \\ 0.32 & y = 2 \end{cases}$$

Example 6.1.6 (Example 3.5)

Suppose that a penny and a nickel are each tossed 10 times so that every pair of sequences of tosses (n tosses in each sequence) is equally likely to occur. Let X be the number of heads obtained with the penny, and Y be the number of heads obtained with the nickel. It can be shown that (show it!) the joint pmf of X and Y is as follows.

$$P(X = x, Y = y) = \begin{cases} \binom{10}{x} \binom{10}{y} \left(\frac{1}{2}\right)^{20} & x, y = 0, \dots, 10 \\ 0 & \text{otherwise} \end{cases}$$

 **Solution**

Note that the support set of X and Y are the same, i.e.

$$A_X = A_Y = \{0, 1, \dots, 10\}.$$


We may assume that the penny and the nickel are fair coins, i.e. if we let p_x and p_y be the probability of getting a head for a penny and nickel, respectively, then $p_x = p_y = \frac{1}{2}$. Since there are 10 ways to get x heads with the penny, and similarly so for the nickel, we have that

$$\begin{aligned} P(X = x, Y = y) &= \begin{cases} \binom{10}{x} \binom{10}{y} \left(\frac{1}{2}\right)^{10} \left(\frac{1}{2}\right)^{10} & x, y = 0, 1, \dots, 10 \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} \binom{10}{x} \binom{10}{y} \left(\frac{1}{2}\right)^{20} & x, y = 0, 1, \dots, 10 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

as required.

“ Note

It is interesting to observe that the two rvs in the last example have seemingly no relationship with one another in terms of the experiment conducted, since they do not affect each other. This leads us to introducing the next concept.

6.1.4 Independence of Discrete RVs **Definition 32 (Independence of Discrete RVs)**

Two rvs X and Y with joint cdf F are said to be **independent** if and only if

$$\forall x, y \in \mathbb{R} \quad F(x, y) = F_X(x)F_Y(y)$$

 **Theorem 23 (Independence by PF)**

Suppose X and Y are rvs with joint cdf F , joint pf f , marginal cdf F_X and F_Y respectively, and marginal pf f_X and f_Y respectively. Also, suppose that $A_X = \{x : f_X(x) > 0\}$ is the support set of X and $A_Y = \{y : f_Y(y) > 0\}$ is the support set of Y . Then X and Y are independent rvs if and only if either

$$\forall (x, y) \in A_X \times A_Y \quad f(x, y) = f_X(x)f_Y(y)$$

holds, or

$$\forall x, y \in \mathbb{R} \quad F(x, y) = F_X(x)F_Y(y)$$

I am not certain as to why this is presented as a theorem that repeats the definition. As so, the prove for the 2nd equation will not be shown.

 **Proof**

The (\implies) direction is simply a result of **Clairaut's Theorem**². While the (\impliedby) direction is a direct result of applying double integrals. □

² Work needs to be done to show that our statement actually satisfies the condition for Clairaut's Theorem to apply. Clairaut's Theorem states that:

 **Theorem 24 (Clairaut's Theorem)**

If (x_0, y_0) is a point in the domain of a function f with

- f is defined on all points in an open disk centered at (x_0, y_0) ;
- the first partial derivatives, f_{xy} and f_{yx} are all continuous for all points in the open disk.


Then $f_{xy}(x_0, y_0) = f_{yx}(x_0, y_0)$.

Example 6.1.7 (Example 3.6)

Suppose X and Y are discrete rvs with joint pf

$$f(x, y) = \frac{\theta^{x+y} e^{-2\theta}}{x!y!} \mathbb{1}_{\{x, y=0, 1, \dots\}}.$$

Are X and Y independent of each other?

 **Solution**

Note that we may write f as

$$f(x, y) = \left(\frac{\theta^x e^{-\theta}}{x!} \cdot \frac{\theta^y e^{-\theta}}{y!} \right) \mathbb{1}_{\{x, y=0, 1, \dots\}}$$

and so this suggests that we can indeed break down f into two parts, each only affected by x and y respectively, "indenpdent" of each

other. Indeed, since

$$\begin{aligned}
 f_X(x) &= \sum_{y=0}^{\infty} \frac{\theta^{x+y} e^{-\theta}}{x!y!} \mathbb{1}_{\{x,y=0,1,\dots\}} \\
 &= \sum_{y=0}^{\infty} \left(\frac{\theta^x e^{-\theta}}{x!} \cdot \frac{\theta^y e^{-\theta}}{y!} \right) \mathbb{1}_{\{x=0,1,\dots\}} \\
 &= \frac{\theta^x e^{-\theta}}{x!} \underbrace{\sum_{y=0}^{\infty} \frac{\theta^y e^{-\theta}}{y!}}_{\text{sum of pmf of Poi}(\theta)=1} \\
 &= \frac{\theta^x e^{-\theta}}{x!}
 \end{aligned}$$

Similarly, we can obtain

$$f_Y(y) = \frac{\theta^y e^{-\theta}}{y!}$$

Multiplying $f_X(x)$ and $f_Y(y)$ together, we indeed get back to the original joint pmf.

7 Lecture 7 May 24th 2018

7.1 Joint Distributions (Continued)

7.1.1 Independence of Discrete RVs (Continued)

Example 7.1.1 (Example 3.7)

Consider the joint pmf below from Example 6.1.3. Are X and Y independent? Prove or disprove.

	$x = -2$	$x = 0$	$x = 2$	$P(Y = y)$
$y = 0$	0.05	0.1	0.15	0.3
$y = 1$	0.07	0.11	0.2	0.38
$y = 2$	0.02	0.25	0.05	0.32
$P(X = x)$	0.14	0.46	0.4	

Solution

Note that

$$P(X = -2, Y = 0) = 0.05 \text{ but}$$
$$P(X = -2)P(Y = 0) = 0.14 \cdot 0.3 = 0.042 \neq 0.05.$$

Thus X and Y are not independent.

7.1.2 Joint Continuous RVs

Definition 33 (Joint Continuous RVs)

Two random variables X and Y are said to be **jointly continuous** if there

exists a function $f(x, y)$ such that the joint cdf of X and Y can be written as

$$\forall (x, y) \in \mathbb{R}^2 \quad F(x, y) = \int_{-\infty}^x \int_{-\infty}^y f(t_1, t_2) dt_2 dt_1.$$

The function f is called the **joint density function** of X and Y . It follows from the above definition that when the second partial derivative exists, we have

$$f(x, y) = \frac{\partial^2}{\partial x \partial y} F(x, y)$$

The set $\{(x, y) : f(x, y) > 0\}$ is called the **support set** of (X, Y) .

“ Note (Convention)

Define $f(x, y) = 0$ when $\frac{\partial^2}{\partial x \partial y} F(x, y)$ does not exist.

Example 7.1.2 (Example 3.8)

Suppose X and Y have joint pdf $f(x, y) = \mathbb{1}_{\{0 < x, y < 1\}} = \mathbb{1}_{\{0 < x < 1, 0 < y < 1\}}$. Calculate the joint cdf of X and Y .

Solution

$$F(x, y) = \begin{cases} 0 & x \leq 0, \forall y \leq 0 \\ \int_0^x \int_0^y 1 ds dt = xy & 0 < x < 1 \wedge 0 < y < 1 \\ \int_0^1 \int_0^y 1 ds dt = y & x \geq 1 \wedge 0 < y < 1 \\ \int_0^x \int_0^1 1 ds dt = x & 0 < x < 1 \wedge y \geq 1 \\ \int_0^1 \int_0^1 1 ds dt = 1 & x \geq 1 \wedge y \geq 1 \end{cases}$$

♦ Proposition 25 (Properties of Joint PDF)

1. $\forall (x, y) \in \mathbb{R}^2 \quad f(x, y) \geq 0$
2. $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) dx dy = 1$

3. $\forall B \subset \mathbb{R}^2$,

$$P[(X, Y) \in B] = \int \int_{(x,y) \in B} f(x, y) dx dy$$

 **Proof**

Example 7.1.3 (Example 3.9)

Suppose that $f(x, y) = Kxy \cdot \mathbb{1}_{\{0 < x, y < 1\}}$ for some constant $K > 0$. Find K so that f is a valid joint pdf. If X and Y have the joint density f , calculate $P(X > Y)$.


 **Solution**

Note that

$$1 = \int_0^1 \int_0^1 Kxy dx dy = \frac{K}{4}.$$

Thus $K = 4$. To solve the next part, observe that for $X > Y$, we have the diagram to the right to show the support set of the joint distribution. The shaded region is the support set. We then have

$$\begin{aligned} P(X > Y) &= \int_0^1 \int_0^x 4xy dy dx = \int_0^1 2xy^2 \Big|_0^x dx \\ &= \int_0^1 2x^3 dx = \frac{1}{2} x^3 \Big|_0^1 = \frac{1}{2} \end{aligned}$$



Example 7.1.4 (Example 3.10)


Suppose that

$$f(x, y) = \begin{cases} Cxy & 0 < x, y < 1, x + y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Find C so that $f(x, y)$ is a valid joint probability density function, and calculate $P(Y^2 < X)$.


 **Solution**

Note that the diagram on the right shows the support set of (X, Y) .



To calculate $P(Y^2 < X)$, note the diagram to the right. Then

$$\begin{aligned} P(Y^2 < X) &= \int_0^{\frac{3+\sqrt{5}}{2}} \int_0^{\sqrt{x}} 24xy \, dy \, dx + \int_{\frac{3+\sqrt{5}}{2}}^1 \int_0^{1-x} 24xy \, dy \, dx \\ &= \int_0^{\frac{3+\sqrt{5}}{2}} 12xy^2 \Big|_0^{\sqrt{x}} \, dx + \int_{\frac{3+\sqrt{5}}{2}}^1 12xy^2 \Big|_0^{1-x} \, dx \\ &= 4x^3 \Big|_0^{\frac{3+\sqrt{5}}{2}} + \int_{\frac{3+\sqrt{5}}{2}}^1 24(x^3 - 2x^2 + x) \, dx \\ &= 4 \left(\frac{3 + \sqrt{5}}{2} \right)^3 + 24 \left[\frac{1}{4}x^4 - \frac{2}{3}x^3 + \frac{1}{2}x^2 \right] \Big|_{\frac{3+\sqrt{5}}{2}}^1 = \dots \end{aligned}$$



Solve for $y = 1 - x$ and $y^2 = x$ to get the intersection.

We shall not proceed to get the final solution since it is a messy process and the result is not important.

7.1.3 Marginal Distribution (Continuous)

Definition 34 (Marginal PDF)

Suppose X and Y are continuous rvs with joint pdf f . Then the marginal pdf of X is given by

$$\forall x \in \mathbb{R} \quad f_X(x) = \int_{-\infty}^{\infty} f \, dy,$$

and the marginal pdf of Y is

$$\forall y \in \mathbb{R} \quad f_Y(y) = \int_{-\infty}^{\infty} f \, dx.$$

Example 7.1.5 (Example 3.11)

Suppose X and Y have joint pdf $f(x, y) = K(x + y)\mathbb{1}_{0 \leq x < y \leq 1}$ for some constant K . Find K . Then, calculate the marginal density of X .

Solution

A diagram showing the region of the support set is on the right.

To get K ,

$$\begin{aligned} 1 &= \int_0^1 \int_x^1 K(x+y) dy dx = \int_0^1 \left(Kxy + \frac{1}{2}Ky^2 \right) \Big|_x^1 dx \\ &= \int_0^1 Kx + \frac{K}{2} - Kx^2 - \frac{1}{2}Kx^2 dx \\ &= \frac{K}{2} \left(x^2 + x - x^3 \right) \Big|_0^1 = \frac{K}{2} \end{aligned}$$

Thus $K = 2$.

To get the marginal density of X , note that our joint pdf is now the following:


$$f(x, y) = 2(x+y)\mathbb{1}_{\{0 \leq x < y \leq 1\}}$$

Thus

$$\int_x^1 2(x+y) dy = 2xy + y^2 \Big|_x^1 = 2x + 1 - 3x^2$$

And hence

$$f_X(x) = \begin{cases} -3x^2 + 2x + 1 & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$



7.1.4 Independence of Continuous RVs

Definition 35 (Independence of Continuous RVs)

Two random variables X and Y with joint cdf F and joint pdf f are independent iff

$$\forall x, y \in \mathbb{R} \quad F(x, y) = F_X(x)F_Y(y)$$

or¹

$$\forall x, y \in \mathbb{R} \quad f(x, y) = f_X(x)f_Y(y).$$

¹ It's really an "AND"

Note

A necessary, but insufficient, condition for X and Y to be independent is that

$$\text{supp}(X, Y) = \text{supp}(X) \times \text{supp}(Y)$$

Example 7.1.6 (Example 3.12)

Are random variables X and Y introduced in Example 7.1.5 independent? Explain.

 Solution

Recall that the pdf was given as

$$f(x, y) = 2(x + y)\mathbb{1}_{\{1 \leq x < y \leq 1\}}.$$

We derived the marginal pdf of X in the earlier example:

$$f_X(x) = (-3x^2 + 2x + 1)\mathbb{1}_{\{0 \leq x \leq 1\}}.$$

To get the marginal pdf of Y , note

$$\int_0^y 2(x + y) dx = x^2 + 2xy \Big|_0^y = 3y^2.$$

Thus

$$f_Y(y) = \begin{cases} 3y^2 & 0 \leq y \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

Note that

$$f_X(x)f_Y(y) = -9x^2y^2 + 6xy^2 + 3y^2 \quad 0 \leq x < y \leq 1$$

which is not equal to f . Thus, X and Y are not independent.

8 Lecture 8 May 29th 2018

8.1 Joint Distributions (Continued 2)

8.1.1 Independence of Continuous RVs (Continued)

Example 8.1.1 (Example 3.12 (3.4.8 course note))

Suppose X and Y are continuous with joint pdf

$$f(x, y) = \frac{3}{2}y(1 - x^2)\mathbb{1}_{\{-1 \leq x \leq 1\}}\mathbb{1}_{\{0 \leq y \leq 1\}}$$

Are X and Y independent?

 **Solution**

The marginal pdf of X is

$$\begin{aligned} f_X(x) &= \int_0^1 \frac{3}{2}y(1 - x^2) dy = \frac{3}{4}y^2(1 - x^2) \Big|_0^1 \\ &= \begin{cases} \frac{3}{4}(1 - x^2) & -1 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

The marginal pdf of Y is

$$\begin{aligned} f_Y(y) &= \int_{-1}^1 \frac{3}{2}y(1 - x^2) dx = \frac{3}{2}y \left(x - \frac{1}{3}x^3 \right) \Big|_{-1}^1 \\ &= \begin{cases} 2y & 0 \leq y \leq 1 \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Clearly, we have

$$f_X(x)f_Y(y) = \frac{3}{2}y(1 - x^2) = f(x, y) \quad -1 \leq x \leq 1, 0 \leq y \leq 1.$$

Thus X and Y are independent.

▣ Theorem 26 (Factorization Theorem for Independence)

Suppose X and Y are rvs with joint pf f , and marginal pf f_X and f_Y , respectively. Suppose also that

$A = \{(x, y) : f(x, y) > 0\}$ is the support set of (X, Y)

$A_X = \{x : f_X(x) > 0\}$ is the support set of X , and

$A_Y = \{y : f_Y(y) > 0\}$ is the support set of Y

Then X and Y are independent rvs iff $A = A_X \times A_Y$ and there exist non-negative functions g and h such that

$$f(x, y) = g(x)h(y)$$

for all $(x, y) \in A_X \times A_Y$.

✎ Proof

The \implies direction is straightforward: Since X and Y are independent, we have that $f = f_X f_Y$, and so clearly, $A = A_X \times A_Y$ and so $\forall (x, y) \in A = A_X \times A_Y$, we have that f_X and f_Y are non-negative.

For the \impliedby direction, note that

$$\begin{aligned} f_Y(y) &= \int_{x \in A_X} g(x)h(y) dx = h(y) \int_{x \in A_X} g(x) dx \\ f_X(x) &= \int_{y \in A_Y} g(x)h(y) dy = g(x) \int_{y \in A_Y} h(y) dy. \end{aligned}$$


Thus,

$$\begin{aligned} f_X(x)f_Y(y) &= g(x)h(y) \int_{x \in A_X} g(x) dx \int_{y \in A_Y} h(y) dy \\ &= g(x)h(y) \int_{x \in A_X} \int_{y \in A_Y} g(x)h(y) dy dx = g(x)h(y) \end{aligned}$$

where line 2 is by **linearity of integration**. Thus $f(x, y) = f_X(x)f_Y(y)$.

Thus X and Y are independent. \square

“ Note

1. If  Theorem 26 holds, then f_X will be proportional to g and f_Y will be proportional to h . Clearly so, since

$$g(x) \cdot h(y) = f_X(x)f_Y(y)$$

$$g(x) \propto f_X(x) \wedge h(y) \propto f_Y(y)$$

2. The definitions and theorems can be easily extended to the random vector (X_1, X_2, \dots, X_n) . Indeed, if we apply mathematical induction on the proof above, we will be able to get our desired result.¹

¹ I wonder if this statement is equivalent to the Fisher-Neyman Factorization Theorem.

8.1.2 Conditional Distributions

Definition 36 (Conditional Distributions)

Suppose X and Y are rvs with joint pf f , and marginal pfs f_X and f_Y , respectively. Suppose also that $A = \{(x, y) : f(x, y) > 0\}$. The **conditional pf** of X given $Y = y$ is given by

$$f_X(x|y) = \frac{f(x, y)}{f_Y(y)}$$

for $(x, y) \in A$ provided that $f_Y(y) \neq 0$. The **conditional pf** of Y given $X = x$ is given by

$$f_Y(y|x) = \frac{f(x, y)}{f_X(x)}$$

for $(x, y) \in A$ provided that $f_X(x) \neq 0$.

Remark

If X and Y are discrete rvs then

$$f_X(x|y) = P(X = x|Y = y) = \frac{P(X = x, Y = y)}{P(Y = y)} = \frac{f(x, y)}{f_Y(y)}$$

and

$$\sum_x f_X(x|y) = \sum_x \frac{f(x, y)}{f_Y(y)} = \frac{1}{f_Y(y)} \sum_x f(x, y) = \frac{f_Y(y)}{f_Y(y)} = 1,$$

and similarly so for $f_Y(y|x)$. Similarly, if X and Y are both continuous rvs,

then

$$\int_{-\infty}^{\infty} f_X(x|y) dx = \int_{-\infty}^{\infty} \frac{f(x,y)}{f_Y(y)} dx = \frac{1}{f_Y(y)} \int_{-\infty}^{\infty} f(x,y) dx = \frac{f_Y(y)}{f_Y(y)} = 1$$

Now consider if X is a continuous rv such that $f_X(x) \neq P(X = x)$ and $P(X = x) = 0$ for all x . Then to justify the definition of the conditional pdf of Y given $X = x$, when X and Y are both continuous rvs, we consider $P(Y \leq y|X = x)$ using the limit approach:

$$\begin{aligned} P(Y \leq y|X = x) &= \lim_{h \rightarrow 0} P(Y \leq y|x \leq X \leq x+h) \\ &= \lim_{h \rightarrow 0} \frac{\int_x^{x+h} \int_{-\infty}^y f(u,v) dv du}{\int_x^{x+h} f_X(u) du} \\ &= \lim_{h \rightarrow 0} \frac{\frac{d}{dh} \int_x^{x+h} \int_{-\infty}^y f(u,v) dv du}{\frac{d}{dh} \int_x^{x+h} f_X(u) du} \quad \text{by L'Hôpital's Rule} \\ &= \lim_{h \rightarrow 0} \frac{\int_{-\infty}^y \frac{d}{dh} \int_x^{x+h} f(u,v) du dv}{\frac{d}{dh} \int_x^{x+h} f_X(u) du} \quad (1) \\ &= \lim_{h \rightarrow 0} \frac{\int_{-\infty}^y f(x+h,v) dv}{f_X(x+h)} \quad (2) \\ &= \frac{\int_{-\infty}^y f(x,v) dv}{f_X(x)} \end{aligned}$$

where (1) is by assuming that the integrands are all convergent so that we may interchange the integral signs and the differential operator, and (2) by the **Fundamental Theorem of Calculus**. If we differentiate the last line with respect to y , by the Fundamental Theorem of Calculus, we have

$$\frac{d}{dy} P(Y \leq y|X = x) = \frac{f(x,y)}{f_X(x)}$$

which justifies the using of our definition

$$f_Y(y|x) = \frac{f(x,y)}{f_X(x)}.$$

Example 8.1.2

A fair coin is flipped 10 times independently.

1. What is the distribution of Y , the number of heads in 10 flips?
2. Suppose the first 4 flips have all landed on tails. What is the distribution of Y given this information?

 **Solution**

- Clearly, we know that $Y \sim \text{Bin}\left(10, \frac{1}{2}\right)$.
- Since each flip is independent of each other and the first four flips have already been determined, the range of values for Y changes from $\{0, \dots, 10\}$ to $\{0, \dots, 6\}$. Since the experiment is still essentially the same, we have that

$$Y \mid \text{first 4 flips are tails} \sim \text{Bin}\left(6, \frac{1}{2}\right).$$

Example 8.1.3 (Example 3.13)

Consider the experiment carried out in [Example 8.1.2](#). Let

$X :=$ number of heads in the first 4 flips

$Y :=$ number of heads in 10 flips

Derive the conditional distribution of Y given that the first 4 flips landed on heads, i.e. derive the distribution for $Y \mid X = 4$.

 **Solution**

Let W be the number of heads in the last 6 flips. Then W has the same distribution as in part 2 of our earlier example. Also, $X \sim \text{Bin}\left(4, \frac{1}{2}\right)$. Clearly, $Y = X + W$. We proceed to derive the joint pf of X and Y :

$$\begin{aligned} P(X = x, Y = y) &= P(X = x, X + W = y) = P(X = x, W = y - x) \\ &= P(X = x)P(W = y - x) \quad \text{by Independence} \\ &= \binom{4}{x} \left(\frac{1}{2}\right)^x \left(\frac{1}{2}\right)^{4-x} \cdot \binom{6}{y-x} \left(\frac{1}{2}\right)^{y-x} \left(\frac{1}{2}\right)^{6-y+x} \\ &= \binom{4}{x} \binom{6}{y-x} \left(\frac{1}{2}\right)^{10} \end{aligned}$$

Then

$$\begin{aligned} P(Y \mid X = 4) &= \frac{P(X = 4, Y = y)}{P(X = 4)} \\ &= \frac{\binom{4}{4} \binom{6}{y-4} \left(\frac{1}{2}\right)^{10}}{\binom{4}{4} \left(\frac{1}{2}\right)^4 \left(\frac{1}{2}\right)^0} \\ &= \binom{6}{y-4} \left(\frac{1}{2}\right)^6 \quad y \in \{4, 5, \dots, 10\}. \end{aligned}$$

We may also re-label the conditional distribution to have

$$\binom{6}{y^*} \left(\frac{1}{2}\right)^6 \quad y^* \in \{0, \dots, 6\}$$

Example 8.1.4 (Example 3.14)

From Example 7.1.5, we had that

$$f(x, y) = 2(x + y)\mathbb{1}_{\{0 \leq x < y \leq 1\}}$$

and the marginal density of X is

$$f_X(x) = (2x - 3x^2 + 1)\mathbb{1}_{\{0 \leq x < 1\}}.$$

Derive the conditional distribution of $Y|X = \frac{1}{2}$.

Solution

Observe that

$$f(y|X = \frac{1}{2}) = \frac{f(\frac{1}{2}, y)}{f_X(\frac{1}{2})} = \frac{2(\frac{1}{2} + y)}{2(\frac{1}{2}) - 3(\frac{1}{2})^2 + 1} = \frac{8}{13}(1 + 2y)$$

for $\frac{1}{2} < y \leq 1$.

Proposition 27 (Properties of Conditional Distributions)


Let X and Y be rvs. If both X and Y are discrete, then

- $\sum_x f(x|y) = 1$;
- $F(x|y) = \sum_{\{w:w \leq x\}} f(w|y)$; and
- $f(x|y) = F(x|y) - F(x^-|y)$.

If X and Y are both continuous, then

- $\int_x f(x|y) dx = 1$;
- $F(x|y) = \int_{-\infty}^x f(t|y) dt$; and
- $f(x|y) = \frac{\partial}{\partial x} F(x|y)$

Exercise 8.1.1

Prove  Proposition 27.

Theorem 28 (Product Rule)

Suppose X and Y are rvs with joint pf f , marginal pfs $f_X(x)$ and $f_Y(y)$ respectively, and conditional pfs $f_X(x|y)$ and $f_Y(y|x)$ respectively. Then

$$f(x, y) = f_X(x|y)f_Y(y) = f_Y(y|x)f_X(x).$$

 **Proof**

Notice once and for all that by rearranging the definition of conditional distribution

$$f_X(x|y)f_Y(y) = f(x, y) = f_Y(y|x)f_X(x)$$

□

 **Proposition 29 (Independence from Conditionality)**

Suppose X and Y are rvs with marginal pfs $f_X(x)$ and $f_Y(y)$ respectively, and conditional pfs $f_X(x|y)$ and $f_Y(y|x)$ respectively. Let $A_X = \{x : f_X(x) > 0\}$ and $A_Y = \{y : f_Y(y) > 0\}$. X and Y are independent rvs iff either of the following holds:

$$\forall x \in A_X \quad f_X(x|y) = f_X(x)$$

or

$$\forall y \in A_Y \quad f_Y(y|x) = f_Y(y).$$

 **Proof**

Suppose that X and Y are independent rvs. Then


$$f(x, y) = f_X(x)f_Y(y).$$

Then

$$f_X(x|y) = \frac{f(x, y)}{f_Y(y)} = \frac{f_X(x)f_Y(y)}{f_Y(y)} = f_X(x)$$

$$f_Y(y|x) = \frac{f(x, y)}{f_X(x)} = \frac{f_X(x)f_Y(y)}{f_X(x)} = f_Y(y).$$

for $x \in A_X$ and $y \in A_Y$.

Suppose WLOG that $f_X(x|y) = f_X(x)$. Thus by  Theorem 28

$$f(x, y) = f_X(x|y)f_Y(y) = f_X(x)f_Y(y) \text{ for } x \in A_X, y \in A_Y.$$

□

Example 8.1.5 (Example 3.15)

In a game of chance, a random number is generated from $P \sim \text{Beta}(\alpha, \beta)$. Given $P = p$, a coin with $P(\{H\}) = p$ is flipped independently n times, where the player is rewarded the same amount of dollars as the number of heads in n . Calculate the probability that a random player earns at least \$1 in this game.

Solution

Let X be the number of heads that appear in n flips, which equates to the total amount of \$1 earned. Then

$$X | P = p \sim \text{Bin}(n, p)$$

However, note that

$$P(X \geq 1) = P(\text{earn at least } \$1) = 1 - P(\text{earn nothing}) = 1 - P(X = 0)$$

To get $P(X = x)$, we need to do the following: note that the support set of P is from 0 to 1, then

$$\begin{aligned} \Pr(X = x) &= \int_0^1 \Pr(X = x, P = p) dp \\ &= \int_0^1 \Pr(X = x | P = p) \Pr(P = p) dp \quad \text{by 1 Theorem 28} \\ &= \int_0^1 \binom{n}{x} p^x (1-p)^{n-x} \frac{p^{\alpha-1} (1-p)^{\beta-1}}{B(\alpha, \beta)} dp \\ &= \binom{n}{x} \frac{1}{B(\alpha, \beta)} \int_0^1 p^{\alpha+x-1} (1-p)^{\beta+n-x-1} dp \\ &= \binom{n}{x} \frac{B(\alpha+x, \beta+n-x)}{B(\alpha, \beta)} \underbrace{\int_0^1 \frac{p^{\alpha+x-1} (1-p)^{\beta+n-x-1}}{B(\alpha+x, \beta+n-x)} dp}_{\text{pdf of Beta}(\alpha+x, \beta+n-x)} \\ &= \binom{n}{x} \frac{B(\alpha+x, \beta+n-x)}{B(\alpha, \beta)} \end{aligned}$$

Definition 37 (Beta Distribution)

If $X \sim \text{Beta}(\alpha, \beta)$, then

$$f(x) = \frac{x^{\alpha-1} (1-x)^{\beta-1}}{B(\alpha, \beta)}$$

where

$$B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$$

Therefore,

$$\begin{aligned} P(X \geq 1) &= 1 - P(X = 0) = 1 - \binom{n}{0} \frac{\Gamma(\alpha)\Gamma(\beta+n)}{\Gamma(\alpha+\beta+n)} \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \\ &= 1 - \frac{\Gamma(\alpha+\beta)\Gamma(\beta+n)}{\Gamma(\alpha+\beta+n)\Gamma(\beta)} \end{aligned}$$

8.1.3 Joint Expectations

Definition 38 (Joint Expectation)

Suppose $h(x, y)$ is a real-valued function. If X and Y are discrete rvs with joint pf f and support A , then

$$E[h(x, y)] = \sum_{(x, y) \in A} h(x, y) f(x, y).$$

provided that the joint sum converges absolutely.

If X and Y are continuous rvs with joint pf f , then

$$E[h(x, y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} h(x, y) f(x, y) dx dy$$

provided that the joint integral converges absolutely.

This is also known as the **Law of the Unconscious Statistician** for two rvs.

Example 8.1.6 (Example 3.16)

Consider X and Y with the following joint probability distribution. Calculate $E(XY)$.

	$x = -2$	$x = 0$	$x = 2$
$y = 0$	0.05	0.1	0.15
$y = 1$	0.07	0.1	0.2
$y = 2$	0.02	0.25	0.05

9 Lecture 9 May 31st 2018

9.1 Joint Distributions (Continued 3)

9.1.1 Joint Expectations (Continued)


Theorem 30 (Linearity of Expectation in Bivariate Case)

Suppose X and Y are two rvs with joint pf f , a_i, b_i , for $i = 1, \dots, n$, are constants, and $g_i(x, y)$, for $i = 1, \dots, n$, are real-valued functions. Then

$$E \left[\sum_{i=1}^n (a_i g_i(X, Y) + b_i) \right] = \sum_{i=1}^n (a_i E[g_i(X, Y)]) + \sum_{i=1}^n b_i$$

provided that $E[g_i(X, Y)]$ is finite for $i = 1, \dots, n$.

Proof

This is simply an extension of  Theorem 13.

Theorem 31 (Implication of Independence on Joint Expectation)

If X and Y are independent rvs with joint pf f , and $g(x)$ and $h(y)$ are real valued functions, then

$$E[g(X)h(Y)] = E[g(X)]E[h(Y)]$$

✎ Proof

We shall prove for the discrete case and leave the continuous case for future exercises.

Observe that

$$\begin{aligned} E[g(X)h(Y)] &= \sum_x \sum_y g(x)h(y)f(x,y) \quad \because \Phi \text{ Definition 38} \\ &= \sum_x \sum_y g(x)h(y)f_X(x)f_Y(y) \\ &= \sum_x g(x)f_X(x) \sum_y h(y)f_Y(y) \\ &= E[g(X)]E[h(Y)] \end{aligned}$$

where $f_X(x)$ and $f_Y(y)$ are the marginal pfs of X and Y respectively. □

We may repeatedly apply the above proof for n rvs through induction and get the following result.

📖 Theorem 32 (Generalized Implication of Independence on Joint Expectation)

If X_1, X_2, \dots, X_n , for some $n \in \mathbb{N}$, are independent rvs and h_1, h_2, \dots, h_n are real valued functions, then

$$E \left[\prod_{i=1}^n h_i(X_i) \right] = \prod_{i=1}^n E[h_i(X_i)].$$

9.1.2 Covariance

INDEPENDENCE of two rvs X and Y implies that knowledge of the value of X does not provide any information whatsoever about the distribution of Y . Essentially, we can say that there is no “relationship” between X and Y . In statistics, **linear relationships** are often the subject of interest. The strength of a linear relationship is related to **covariance** and measured by the **correlation coefficient**, usually denoted by ρ .

Exercise 9.1.1

Prove 📖 Theorem 31 for the continuous case.

It can be shown that when X and Y have no linear relationship iff their covariance is 0.

On a related thought, does covariance relate to independence? If so, how?

Definition 39 (Covariance)

The **covariance** of rvs X and Y is given by

$$\text{Cov}(X, Y) = E[(X - \mu_X)(Y - \mu_Y)] = E(XY) - \mu_X\mu_Y. \quad (9.1)$$


where $\mu_X = E[X]$ and $\mu_Y = E[Y]$.

If $\text{Cov}(X, Y) = 0$, then X and Y are called **uncorrelated** rvs.

“ Note

Note that the 2nd and 3rd term are equivalent in Equation (9.1) since

$$\begin{aligned} E[(X - \mu_X)(Y - \mu_Y)] &= E[XY - \mu_X Y - \mu_Y X + \mu_X \mu_Y] \\ &= E[XY] - \mu_X E[Y] - \mu_Y E[X] + \mu_X \mu_Y \quad \because \text{Theorem 30} \\ &= E[XY] - \mu_X \mu_Y - \mu_X \mu_Y + \mu_X \mu_Y = E[XY] - \mu_X \mu_Y \end{aligned}$$

From here, it is easy to see from  Theorem 31, that since $E[XY] = E[X]E[Y] = \mu_X \mu_Y$, we have that the independence of X from Y will imply that $\text{Cov}(X, Y) = 0$.

However, the converse of the above is **not true**.

Example 9.1.1

Source: Stats SE

Let X be an rv that it is -1 or 1 with probability 0.5 . Then let Y be an rv such that $Y = 0$ if $X = -1$, and Y is randomly -1 or 1 with probability 0.5 if $X = 1$.

Clearly X and Y are highly dependent (since knowing Y allows me

to perfectly know X). They both have zero mean:

$$E[X] = -1 \left(\frac{1}{2} \right) + 1 \left(\frac{1}{2} \right) = 0$$

$$E[Y] = -1 \left(\frac{1}{2} \right) + 1 \left(\frac{1}{2} \right) = 0$$

and

$$E[XY] = (-1) \cdot 0P(X = -1, Y = 0) + 1(-1) \cdot P(X = 1, Y = -1) \\ + 1(1)P(X = 1, Y = 1) \\ = -\frac{1}{4} + \frac{1}{4} = 0$$

Thus $\text{Cov}(X, Y) = 0$

Or more generally, take any distribution $P(X)$ and any $P(Y|X)$ such that $P(Y = a|X) = P(Y = -a|X)$ for all X (i.e., a joint distribution that is symmetric around the x axis), and you will always have zero covariance. But you will have non-independence whenever $P(Y|X) \neq P(Y)$, i.e., the conditionals are not all equal to the marginal, and vice versa for symmetry around the y axis.

“ Note

- If $\text{Cov}(X, Y) = 0$, then X and Y are called **uncorrelated** rvs.
- By definition, $\text{Cov}(X, X) = \text{Var}(X)$, since

$$\text{Cov}(X, X) = E[(X - \mu_X)^2] = \text{Var}(X)$$

Example 9.1.2 (Example 3.17)


Consider the rvs X and Y with the joint pdf

$$f(x, y) = \begin{cases} 2 & 0 \leq x \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Calculate $\text{Cov}(X, Y)$.

Solution

Observe the diagram of the support set of X and Y to our right.



Then we can calculate

$$\begin{aligned} E[XY] &= \int_0^1 \int_x^1 2xy \, dy \, dx = \int_0^1 xy^2 \Big|_x^1 \, dx \\ &= \int_0^1 x - x^3 \, dx = \frac{1}{2} - \frac{1}{4} = \frac{1}{4} \end{aligned}$$

$$f_X(x) = \int_x^1 2 \, dy = \begin{cases} 2 - 2x & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$E[X] = \int_0^1 x(2 - 2x) \, dx = 1 - \frac{2}{3} = \frac{1}{3}$$

$$f_Y(y) = \int_0^y 2 \, dx = \begin{cases} 2y & 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$E[Y] = \int_0^1 2y^2 \, dy = \frac{2}{3}$$

Thus

$$\text{Cov}(X, Y) = \frac{1}{4} - \frac{1}{3} \left(\frac{2}{3} \right) = \frac{1}{36}$$

We observe that the covariance is positive. This implies a **positive linear relationship**. However, we cannot tell from this value the strength of the relationship between X and Y .

Example 9.1.3 (Example 3.18)

Consider rvs X and Y with the joint pf

$$f(x, y) = \frac{xy}{7} \mathbb{1}_{\{y=1,2\}} \mathbb{1}_{\{x=0,\dots,y\}}.$$

Calculate $\text{Cov}(X, Y)$.

Solution

The following table captures all the probabilities that can be found from the given pf

	$x = 0$	$x = 1$	$x = 2$
$y = 1$	0	$\frac{1}{7}$	0
$y = 2$	0	$\frac{2}{7}$	$\frac{4}{7}$

Observe that we thus have

$$f_X(x) = \begin{cases} \frac{3}{7} & x = 1 \\ \frac{4}{7} & x = 2 \end{cases}$$

$$E[X] = \frac{3}{7} + 2\left(\frac{4}{7}\right) = \frac{11}{7}$$

$$f_Y(y) = \begin{cases} \frac{1}{7} & y = 1 \\ \frac{6}{7} & y = 2 \end{cases}$$

$$E[Y] = \frac{1}{7} + 2\left(\frac{6}{7}\right) = \frac{13}{7}$$

Also,

$$E[XY] = \frac{1}{7} + 2\left(\frac{2}{7}\right) + 4\left(\frac{4}{7}\right) = 3$$

Therefore,

$$\text{Cov}(X, Y) = E[XY] - E[X]E[Y] = 3 - \frac{11}{7} \frac{13}{7} = \frac{4}{49}$$

Theorem 33 (Variance of Linear Combinations)

Suppose X and Y are rvs and a, b, c are real constants. Then

$$\text{Var}(aX + bY + c) = a^2 \text{Var}(X) + b^2 \text{Var}(Y) + 2ab \text{Cov}(X, Y)$$

Proof

Let $E[aX + bY + c] = \mu$. Observe that

$$\begin{aligned} & \text{Var}[aX + bY + c] \\ &= E \left[[(aX + bY + c) - \mu]^2 \right] \\ &= E \left[(aX + bY + c)^2 - 2\mu(aX + bY + c) + \mu^2 \right] \\ &= E \left[a^2X^2 + abXY + acX + abXY + b^2Y^2 + bcY + c^2 \right. \\ & \quad \left. - 2\mu(aX + bY + c) + \mu^2 \right] \\ &= a^2E[X^2] + 2abE[XY] + acE[X] + b^2E[Y^2] + bcE[Y] + c^2 - \mu^2 \end{aligned}$$

Note that

$$\begin{aligned}\mu^2 &= E[aX + bY + c]^2 \\ &= (aE[X] + bE[Y] + c)^2 \\ &= a^2E[X]^2 + b^2E[Y]^2 + 2abE[X]E[Y] + acE[X] + bcE[Y] + c^2\end{aligned}$$

Therefore, we have that

$$\begin{aligned}\text{Var}[aX + bY + c] &= a^2E[X^2] + a^2E[X]^2 + b^2E[Y^2] - b^2E[Y]^2 + 2abE[XY] - 2abE[X]E[Y] \\ &= a^2(E[X^2] - E[X]^2) + b^2(E[Y^2] - E[Y]^2) + 2ab(E[XY] - E[X]E[Y]) \\ &= a^2\text{Var}(X) + b^2\text{Var} Y + 2ab\text{Cov}(X, Y)\end{aligned}$$

as required. \square

By applying [Theorem 33](#) repeatedly, we have the following generalized theorem.

[Theorem 34 \(Generalized Variance of Linear Combinations\)](#)

Suppose X_1, X_2, \dots, X_n are rvs with $\text{Var}(X_i) = \sigma_i^2$, and a_1, a_2, \dots, a_n are real constants. Then

$$\text{Var}\left(\sum_{i=1}^n a_i X_i\right) = \sum_{i=1}^n a_i^2 \sigma_i^2 + 2 \sum_{i=1}^{n-1} \sum_{j=i+1}^n a_i a_j \text{Cov}(X_i, X_j)$$

Note that to prove the above, we have to also use the fact that

$$\text{Cov}(X_i, X_j) = \text{Cov}(X_j, X_i)$$

“ Note

Note that in [Theorem 34](#), if the rvs are independent rvs, then $\text{Cov}(X_i, X_j) = 0$ for $i \neq j$, thus wiping off the 2nd term in the equation, leaving us with

$$\text{Var}\left(\sum_{i=1}^n a_i X_i\right) = \sum_{i=1}^n a_i^2 \sigma_i^2$$

Example 9.1.4 (Example 3.19)

To build a ship engine piece, suppose two pole-shaped components A and B are attached at one end to each other to make one long pole-shaped component C . Suppose the length of part A is an rv with a mean of 3 inches and a variance of 0.25 inch^2 . Similarly, the length of component B is an rv with a mean of 25 inches and a variance of 0.5 inch^2 .

Find the mean and the variance of the length of part C if

1. the lengths of A and B are independent;
2. the covariance between lengths of A and B is -0.3 inch^2 .

 **Solution**

Note that we are given that $C = A + B$

1. We have that

$$E(C) = E(A + B) = E(A) + E(B) = 3 + 25 = 28.$$

For variance, since A and B are independent, $\text{Cov}(A, B) = 0$, thus

$$\begin{aligned} \text{Var}(C) &= \text{Var}(A + B) = \text{Var}(A) + \text{Var}(B) + 2\text{Cov}(A, B) \\ &= 0.25 + 0.5 + 0 = 0.75 \end{aligned}$$

2. Since C is a linear equation, the covariance does not affect the expectation and thus we still have

$$E(C) = 28.$$

Now, given that $\text{Cov}(A, B) = -0.3$, we have

$$\text{Var}(C) = \text{Var}(A) + \text{Var}(B) + 2\text{Cov}(A, B) = 0.75 - 0.6 = 0.15.$$

9.1.3 Correlation

The **covariance** is a real number which depends on the units of measurement of X and Y . The information part of a covariance is its **sign**, unless if it is used as the context.

To use the covariance as the context, and to quantitatively measure the strength of a linear relationship, which we have discussed and

desired before, we use the **correlation coefficient**.


 **Definition 40 (Correlation Coefficient)**

The **correlation coefficient** of rvs X and Y is given by

$$\rho(X, Y) = \frac{\text{Cov}(X, Y)}{\sigma_X \sigma_Y}$$

where $\sigma_X = \sqrt{\text{Var}(X)}$ and $\sigma_Y = \sqrt{\text{Var}(Y)}$.

Note that this is the definition of the **Pearson Correlation Coefficient**. There are other correlation coefficients but we will be using only Pearson, at it seems.

 **Proposition 35 (Properties of the Correlation Coefficient)**

Let X and Y be rvs, and $\rho(X, Y)$ the correlation coefficient of X and Y .

Then

1. $|\rho(X, Y)| \leq 1$;
2. (**perfect positive linear relationship**)
 $\rho(X, Y) = 1 \iff Y = aX + b$ for some $a > 0$;
3. (**perfect inverse linear relationship**)
 $\rho(X, Y) = -1 \iff Y = aX + b$ for some $a < 0$.

 **Proof**

1. This is somewhat beyond the scope of what we can cover now but we shall use this result presented on [Wikipedia](#):

$$|\text{Cov}(X, Y)| \leq \sqrt{\text{Var}(X) \text{Var}(Y)}.$$

Then given the formula of $\rho(X, Y)$, the proof is complete:

$$|\rho(X, Y)| = \left| \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}} \right| \leq \frac{\sqrt{\text{Var}(X) \text{Var}(Y)}}{\sqrt{\text{Var}(X) \text{Var}(Y)}} = 1$$

Proving 2 and 3 is currently outside of my abilities. Refer to this [Math SE Q&A](#) for a hint on how to prove this statement.

Example 9.1.5 (Example 3.20)


Consider rvs X and Y with the joint pdf

$$f(x, y) = \begin{cases} 2 & 0 \leq x \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Calculate $\rho(X, Y)$.

 **Solution**

The diagram to the right is an illustration of the region of support for X and Y . We now calculate the following values:



$$E(XY) = \int_0^1 \int_x^1 2xy \, dy \, dx = \int_0^1 x - x^3 \, dx = \frac{1}{4}$$

$$f_X(x) = \int_x^1 2 \, dy = 2 - 2x \quad 0 \leq x \leq 1$$

$$f_Y(y) = \int_0^y 2 \, dx = 2y \quad 0 \leq y \leq 1$$

$$E(X) = \int_0^1 2x - 2x^2 \, dx = \left(x^2 - \frac{2}{3}x^3 \right) \Big|_0^1 = \frac{1}{3}$$

$$E(X^2) = \int_0^1 2x^2 - 2x^3 \, dx = \left(\frac{2}{3}x^3 - \frac{1}{2}x^4 \right) \Big|_0^1 = \frac{1}{6}$$

$$E(Y) = \int_0^1 2y^2 \, dy = \frac{2}{3}$$

$$E(Y^2) = \int_0^1 2y^3 \, dy = \frac{1}{2}$$

$$\text{Var}(X) = E(X^2) - E(X)^2 = \frac{1}{6} - \left(\frac{1}{3} \right)^2 = \frac{1}{18}$$

$$\text{Var}(Y) = E(Y^2) - E(Y)^2 = \frac{1}{2} - \left(\frac{2}{3} \right)^2 = \frac{1}{18}$$

Therefore we have that

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y) = \frac{1}{4} - \frac{2}{9} = \frac{1}{36}$$

$$\sqrt{\text{Var}(X) \text{Var}(Y)} = \frac{1}{18}$$

and so

$$\rho(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}} = \frac{\frac{1}{36}}{\frac{1}{18}} = \frac{1}{2}$$

9.1.4 Conditional Expectation

Definition 41 (Conditional Expectation)

Let G be a real-valued function. The **conditional expectation** of $g(Y)$ given $X = x$, denoted as $g(Y) | (X = x)$ is given by

$$E[g(Y) | x] = \sum_y g(y) f_Y(y | x)$$

if $Y | (X = x)$ is a discrete rv and

$$E[g(Y) | x] = \int_y g(y) f_Y(y | x)$$

if $Y | (X = x)$ is a continuous rv. This definition only holds provided that the sum and the integral converges absolutely. The conditional expectation of $h(X)$ given $Y = y$, where h is a real-valued function, is defined in a similar manner.

We also call $E[Y | X = x]$ the **conditional mean**, which may be denoted as $E(Y | x)$, and $\text{Var}(Y | X = x)$ the **conditional variance**, which may be denoted as $\text{Var}(Y | x)$.

“ Note

Note that there is also the notation $E(Y | X)$, which is an rv, and hence different from $E(Y | x)$.

Example 9.1.6 (Example 3.21)

Consider $f(x, y) = 8xy \mathbb{1}_{\{0 < x < y < 1\}}$. Calculate the conditional mean and the conditional variance of $X | (Y = \frac{1}{2})$.


Solution

The diagram to the right illustrates the region of support for X and Y . To derive the conditional distribution, we first need $f_Y(y)$.

$$f_Y(y) = \int_0^y 8xy \, dx = 4x^2 y \Big|_0^y = 4y^3$$

Thus

$$f_X \left(X | Y = \frac{1}{2} \right) = \frac{f \left(x, \frac{1}{2} \right)}{f_Y \left(\frac{1}{2} \right)} = \frac{4x}{\frac{1}{2}} = 8x \quad 0 < x < \frac{1}{2}$$



10 Lecture 10 Jun 05th 2018

10.1 Joint Distribution (Continued 4)

10.1.1 Conditional Expectation (Continued)

Example 10.1.1 (Example 3.22)

Given the joint distribution below, calculate $\text{Var}(X | Y = 1)$ and compare it to $\text{Var}(X)$.

	x = -2	x = 0	x = 2	P(Y = y)
y = 0	0.05	0.1	0.15	0.3
y = 1	0.07	0.11	0.2	0.38
y = 2	0.02	0.25	0.05	0.32
P(X = x)	0.14	0.46	0.4	

Solution

Note that

$$\begin{aligned}\text{Var}(X) &= E(X^2) - E(X)^2 \\ &= 4 \cdot 0.14 + 4 \cdot 0.4 - (-2 \cdot 0.14 + 2 \cdot 0.4)^2 = 1.8896\end{aligned}$$

To get $\text{Var}(X | Y = 1)$, we first need

$$f(X | Y = 1) = \frac{P(X = x, Y = 1)}{P(Y = 1)} = \begin{cases} \frac{0.07}{0.38} = 0.1842 & x = -2 \\ \frac{0.11}{0.38} = 0.2895 & x = 0 \\ \frac{0.2}{0.38} = 0.5263 & x = 2 \end{cases}$$

Thus

$$\begin{aligned}\text{Var}(X | Y = 1) &= E[X^2 | Y = 1] - E[X | Y = 1]^2 \\ &= \frac{14}{19} + \frac{40}{19} - \left(\frac{13}{19}\right)^2 = \frac{857}{361} = 2.3740\end{aligned}$$

💡 Proposition 36 (Independence on Conditional Expectation)

If X and Y are independent rvs then $E[g(Y) | x] = E[g(Y)]$ and $E[h(X) | y] = E[h(X)]$.

Proof


We shall prove one of the above for the other will follow a similar argument. Also, we shall prove the continuous case and leave the discrete case as an exercise.

Observe that

$$\begin{aligned}E[g(Y) | X = x] &= \int_y g(y) \frac{f(x, y)}{f_X(x)} dy \\ &= \int_y g(y) \frac{f_X(x) f_Y(y)}{f_X(x)} dy \quad \because \text{independence} \\ &= \int_y g(y) f_Y(y) dy = E[g(Y)]\end{aligned}$$

□

Exercise 10.1.1

Prove the discrete case for  Proposition 36.

Theorem 37 (Law of Total Expectation)

Suppose X and Y are rvs, then

$$E(E[g(Y) | X]) = E[g(Y)]$$


If g is the identity function, we have $E(E[Y | X]) = E(Y)$.

Proof

We shall prove for the discrete case and leave the continuous case as an exercise. Observe that

Exercise 10.1.2

Prove the continuous case for

 Theorem 37.

$$\begin{aligned}
E[g(Y) | X] &= \sum_y [g(y) \cdot P(Y = y | X)] \\
E[E[g(Y) | X]] &= \sum_x \left[\sum_y [g(y) \cdot P(Y = y | X)] \right] P(X = x) \\
&= \sum_x \sum_y g(y) \cdot P(X = x, Y = y) \\
&= \sum_y g(y) \sum_x P(X = x, Y = y) \\
&= \sum_y [g(y) \cdot P(Y = y)] = E[g(Y)]
\end{aligned}$$

□

Example 10.1.2 (Example 3.23 - A Classical Example)

This is a very classical example to illustrate the power of the **Law of Total Expectation**.

A man is lost in a mine, and 3 paths are in front of him. If he takes path 1, after 3 hours, he will be back at his current place. If he takes path 2, the time to get out of the mine (in hours) follows an $\text{Exp}(1)$ distribution. If he takes the 3rd path, he will be back to his current place after 2 hours. Suppose that the man cannot recognize which path he took every time he comes back to the original spot (after going through either path 1 or 3), and so he randomly chooses a path every time he comes back to this original spot. What is the expected time that he will take to get out of the mine?

 **Solution**

Let X an rv that represents the path number, i.e. $X = 1, 2$ or 3 , and let Y represent the total time that the man takes to exit the mine. We are given that

$$P(X = 1) = P(X = 2) = P(X = 3) = \frac{1}{3}.$$

We are also given that

$$\begin{aligned}
E[Y | X = 1] &= 3 + E[Y] \\
E[Y | X = 2] &= 1 \quad \because Y | (X = 2) \sim \text{Exp}(1) \\
E[Y | X = 3] &= 2 + E[Y].
\end{aligned}$$

Therefore, to get the expected time, by  Theorem 37,

$$\begin{aligned}
 E[Y] &= E[E[Y | X]] \\
 &= \frac{1}{3} \cdot E[Y | X = 1] + \frac{1}{3} \cdot E[Y | X = 2] + \frac{1}{3} \cdot E[Y | X = 3] \\
 &= \frac{1}{3} \cdot (3 + E[Y]) + \frac{1}{3} \cdot 1 + \frac{1}{3} \cdot (2 + E[Y]) \\
 &= 2 + \frac{2}{3}E[Y]
 \end{aligned}$$

and hence

$$E[Y] = 6$$

Theorem 38 (Law of Total Variance)

Suppose X and Y are rvs. Then

$$\text{Var}(Y) = E[\text{Var}(Y | X)] + \text{Var}[E(Y | X)]$$

Proof

Note that

$$\begin{aligned}
 \text{Var}(Y|X) &= E(Y^2|X) - E(Y|X)^2 \\
 E[\text{Var}(Y|X)] &= E[E(Y^2|X) - E(Y|X)^2] \\
 &= E[Y^2] - E[E(Y|X)^2] \\
 &= E[Y^2] - \left[\text{Var}(E(Y|X)) + E(E(Y|X))^2 \right] \\
 &= \text{Var}(Y) - \text{Var}(E(Y|X))
 \end{aligned}$$

By rearranging the above, we get

$$\text{Var}(Y) = E[\text{Var}(Y|X)] + \text{Var}[E(Y|X)]$$

□

Example 10.1.3 (Example 3.24 (Course Note 3.7.11))

Suppose $P \sim \text{Unif}(0, 0.1)$ and $Y | P = p \sim \text{Bin}(10, p)$. Find $E(Y)$ and $\text{Var}(Y)$.

 Solution

$$\begin{aligned}
 E[Y] &= E[E[Y|P]] = E[10P] = 10E[P] \\
 &= 10 \cdot \int_0^{0.1} \frac{p}{0.1} dp = \frac{10}{0.2} p^2 \Big|_0^{0.1} = 0.05 \\
 \text{Note: } E[P^2] &= \int_0^{0.1} \frac{p^2}{0.1} dp = \frac{p^3}{0.3} \Big|_0^{0.1} = \frac{0.001}{0.3} = \frac{1}{300} \\
 \text{Var}(Y) &= E[\text{Var}(Y|P)] + \text{Var}[E[Y|P]] \\
 &= E[10P(1-P)] + \text{Var}[10P] \\
 &= 10E[P] - 10E[P^2] + 100 \text{Var}(P) \\
 &= 0.05 - \frac{1}{30} + 100 \left[E[P^2] - E[P]^2 \right] \\
 &= \frac{1}{60} + 100 \left[\frac{1}{300} - 0.05^2 \right] = \frac{1}{60} + 100 \left[\frac{1}{300} - \frac{1}{400} \right] \\
 &= \frac{1}{60} + \frac{1}{12} = \frac{1}{10}
 \end{aligned}$$

10.1.2 Joint Moment Generating Functions

Definition 42 (Joint Moment Generating Functions)

The **joint moment generating function** of two rvs X and Y is defined as

$$M(t_1, t_2) = E \left(e^{t_1 X + t_2 Y} \right)$$

if the expectation exists for all $t_1 \in (-h_1, h_1)$ and $t_2 \in (-h_2, h_2)$ for some $h_1, h_2 > 0$.

More generally, if X_1, X_2, \dots, X_n are rvs, then

$$M(t_1, t_2, \dots, t_n) = E \left[\exp \left(\sum_{i=1}^n t_i X_i \right) \right]$$

is called the **joint mgf** of X_1, X_2, \dots, X_n if the expectation exists for all $t_i \in (-h_i, h_i)$ for some $h_i > 0$, where $i = 1, \dots, n$.

Definition 43 (Joint Moments and Marginal MGF)

Given the joint mgf $M(t_1, t_2)$, we can calculate the joint moments. In

particular,

$$E(X^j Y^k) = \frac{\partial^{j+k}}{\partial t_1^j \partial t_2^k} M(t_1, t_2) \Big|_{(t_1, t_2) = (0, 0)}$$

If $M(t_1, t_2)$ exists for all $t_1 \in (-h_1, h_1)$ and $t_2 \in (-h_2, h_2)$ for some $h_1, h_2 > 0$, then the mgf of X is given by

$$M_X(t) = E(e^{tX}) = M(t, 0) \quad t \in (-h_1, h_1)$$

and the mgf of Y is given by

$$M_Y(t) = E(e^{tY}) = M(0, t) \quad t \in (-h_2, h_2).$$

Example 10.1.4 (Example 3.25)

Given the joint distribution below, calculate the joint mgf $M(t_1, t_2)$, the first joint moment, $E[XY]$, from the joint mgf, and the marginal mgf of X and that of Y .

	$x = -1$	$x = 1$
$y = 1$	0.5	0.3
$y = 2$	0.1	0.1

Solution

Since all probabilities are provided,

$$M(t_1, t_2) = E(e^{t_1 X + t_2 Y}) = \sum_x \sum_y e^{t_1 x + t_2 y} P(X = x, Y = y)$$

$$= 0.5e^{-t_1 + t_2} + 0.3e^{t_1 + t_2} + 0.1e^{-t_1 + 2t_2} + 0.1e^{t_1 + 2t_2}$$

$$\begin{aligned} E(XY) &= \frac{\partial^2}{\partial t_1 \partial t_2} M(t_1, t_2) \Big|_{t_1=0, t_2=0} \\ &= \frac{\partial}{\partial t_1} \left[0.5e^{-t_1 + t_2} + 0.3e^{t_1 + t_2} + 0.2e^{-t_1 + 2t_2} + 0.2e^{t_1 + 2t_2} \right] \Big|_{t_1=0, t_2=0} \\ &= -0.5e^{-t_1 + t_2} + 0.3e^{t_1 + t_2} - 0.2e^{-t_1 + 2t_2} + 0.2e^{t_1 + 2t_2} \Big|_{t_1=0, t_2=0} \\ &= -0.2 \end{aligned}$$

$$\begin{aligned} M_X(t_1) &= M(t_1, 0) = 0.5e^{-t_1} + 0.3e^{t_1} + 0.1e^{-t_1} + 0.1e^{t_1} \\ &= 0.6e^{-t_1} + 0.4e^{t_1} \end{aligned}$$

$$\begin{aligned} M_Y(t_2) &= M(0, t_2) = 0.5e^{t_2} + 0.3e^{t_2} + 0.1e^{2t_2} + 0.1e^{2t_2} \\ &= 0.8e^{t_2} + 0.2e^{2t_2} \end{aligned}$$

◆ **Proposition 39 (Independence on Joint MGF)**

Suppose X and Y are rvs with joint mgf $M(t_1, t_2)$ which exists $\forall t_1 \in (-h_1, h_1), t_2 \in (-h_2, h_2)$, for some $h_1, h_2 > 0$. Then X and Y are independent rvs iff

$$\forall t_1 \in (-h_1, h_1), t_2 \in (-h_2, h_2) \quad M(t_1, t_2) = M_X(t_1)M_Y(t_2)$$

where $M_X(t_1) = M(t_1, 0)$ and $M_Y(t_2) = M(0, t_2)$.

✎ **Proof**
to be proven later

Example 10.1.5 (Example 3.26 (Course Note 3.8.5))

Suppose X and Y are continuous rvs with joint pdf

$$f(x, y) = e^{-y} \quad 0 < x < y < \infty$$


Find the joint mdf of X and Y . Are X and Y independent rvs? What is the marginal mgf of X and Y ?

✎ **Solution**

$$\begin{aligned} M(t_1, t_2) &= E[e^{t_1 X + t_2 Y}] = \int_0^\infty \int_0^y e^{t_1 x + t_2 y} e^{-y} dx dy \\ &= \int_0^\infty \frac{1}{t_1} e^{t_1 x + t_2 y - y} \Big|_0^y dy \\ &= \int_0^\infty \frac{1}{t_1} [e^{y(t_2 - 1)} - e^{y(t_1 + t_2 - 1)}] dy \\ &= \frac{1}{t_1} \left[\frac{1}{t_2 - 1} e^{y(t_2 - 1)} - \frac{1}{t_1 + t_2 - 1} e^{y(t_1 + t_2 - 1)} \right] \Big|_0^\infty \\ &= \frac{1}{t_1} \left[\frac{t_1}{(t_2 - 1)(t_1 + t_2 - 1)} \right] \\ &= \frac{1}{(t_2 - 1)(t_1 + t_2 - 1)} \quad t_2 < 1 \wedge t_1 + t_2 < 1 \\ M_X(t_1) &= M(t_1, 0) = \frac{1}{t_1 - 1} \quad t_1 < 1 \\ M_Y(t_2) &= M(0, t_2) = \frac{1}{(t_2 - 1)^2} \quad t_2 < 1 \end{aligned}$$

Observe that

$$M_X(t_1)M_Y(t_2) = \frac{1}{(t_1 - 1)(t_2 - 1)^2} \neq M(t_1, t_2)$$

and so by  Proposition 39, X and Y are not independent.

Example 10.1.6 (Example 3.27)

Investigate the independence of X and Y in Example 10.1.4 using the mgf method.

 **Solution**

We had that

$$M_X(t_1) = 0.6^{-t_1} + 0.4e^{t_1} \quad t_1 \in \mathbb{R}$$

$$M_Y(t_2) = 0.8e^{t_2} + 0.2e^{2t_2} \quad t_2 \in \mathbb{R}.$$

Since

$$M_X\left(\frac{1}{2}\right)M_Y\left(\frac{1}{2}\right) \neq M\left(\frac{1}{2}, \frac{1}{2}\right)$$

we have that X and Y are not independent.

11 Lecture 11 Jun 07th 2018

11.0.1 Working with Multivariate Cases

Almost everything that has been introduced above can be extended to cases where we have more than just 2 rvs. For example:

Definition 44 (k-variate CDF)

The *k-variate CDF*, $k > 2$, rvs X_1, \dots, X_k is defined as

$$F(x_1, \dots, x_k) = P(X_1 \leq x_1, X_2 \leq x_2, \dots, X_k \leq x_k).$$

In the continuous case, we may write

$$f(x_1, \dots, x_k) = \frac{\partial^k}{\partial x_1 \dots \partial x_k} F(x_1, \dots, x_k).$$

Definition 45 (k-variate Support Set)

The support set of the distribution for X_1, X_2, \dots, X_k is

$$\{(x_1, \dots, x_k) : f(x_1, \dots, x_k) > 0\}$$

We also have the following:

Proposition 40 (Law of Total Probability - Multivariate)

If X_1, \dots, X_k are continuous rvs, then

$$\int_{x_1} \dots \int_{x_k} f(x_1, \dots, x_k) dx_1 \dots dx_k = 1.$$

Should they be discrete, then

$$\sum_{x_1} \dots \sum_{x_k} f(x_1, \dots, x_k) = 1$$

Definition 46 (k-Variate Marginal Distribution)

To get the marginal distribution of a subset of m variables from X_1, \dots, X_k ($1 \leq m \leq k$), we will sum or integrate over the other ones if they are discrete or continuous, respectively. For example,

$$f(x_1, x_2, x_3) = \int_{x_4} \dots \int_{x_k} f(x_1, \dots, x_k) dx_4 \dots dx_k$$

Definition 47 (k-Variate Joint MGF)

The joint mgf of X_1, \dots, X_k is defined as

$$M(t_1, t_2, \dots, t_k) = E\left(e^{t_1 X_1 + \dots + t_k X_k}\right)$$

Proposition 41 (Independence for Multivariate Cases)

If X_1, \dots, X_k are independent, then

$$f(x_1, \dots, x_k) = \prod_{i=1}^k f_{X_i}(x_i) \quad F(x_1, \dots, x_k) = \prod_{i=1}^k F_{X_i}(x_i)$$

$$M(t_1, \dots, t_k) = \prod_{i=1}^k M_{X_i}(t_i)$$

THERE ARE many different examples of multivariate distributions.

We shall discuss two:

- Multinomial Distribution
- Multivariate Normal Distribution

The **multinomial distribution** is an extension of the binomial distribution to cases where there are more categories than two results.

For a multinomial distribution, we have that

- the experiment involves n trials, each with k categories
- the outcome of trials are independent of each other
- the probability of each category, p_i , remains the same across n trials
- $X = (X_1, \dots, X_k) \sim \text{Mult}(n, p_1, \dots, p_k)$ counts the number of elements in each category among the n trials.

Definition 48 (Multinomial Distribution)

Suppose X_1, \dots, X_k are discrete rvs with joint pf

$$f(x_1, \dots, x_k) = \frac{n!}{x_1! x_2! \dots x_k! x_{k+1}!} p_1^{x_1} p_2^{x_2} \dots p_k^{x_k} p_{k+1}^{x_{k+1}}$$

where $x_i = 0, \dots, n$, $x_{k+1} = n - \sum_{i=1}^k x_i$, $0 < p_i < 1$, $p_{k+1} = 1 - \sum_{i=1}^k p_i$, for $i = 1, \dots, k+1$.

Under these conditions, (X_1, \dots, X_k) is said to have a multinomial distribution, and we write $(X_1, \dots, X_k) \sim \text{Mult}(n, p_1, \dots, p_k)$.

Note

Observe that $\text{Bin}(n, p) = \text{Mult}(n, p, p)$.

Proposition 42 (Properties of Multinomial Distribution)

Suppose $(X_1, \dots, X_k) \sim \text{Mult}(n, p_1, \dots, p_k)$, then

1. $\forall (t_1, \dots, t_k) \in \mathbb{R}^k$, the random vector (X_1, \dots, X_k) has joint mgf

$$M(t_1, \dots, t_k) = E\left(e^{t_1 X_1 + \dots + t_k X_k}\right) = (p_1 e^{t_1} + \dots + p_k e^{t_k} + p_{k+1})^n$$

2. Any subset of X_1, \dots, X_{k+1} also has a multinomial distribution. In particular, $X_i \sim \text{Bin}(n, p_i)$, $i = 1, \dots, k+1$.
3. If $T = X_i + X_j$, for $i \neq j$, then $T \sim \text{Bin}(n, p_i + p_j)$
4. $\text{Cov}(X_i, X_j) = -np_i p_j$, for $i \neq j$

5. The conditional distribution of any subset of (X_1, \dots, X_{k+1}) given the rest of the coordinates is a multinomial distribution. In particular, the conditional pf of X_i given $X_j = x_j$, $i \neq j$, is

$$X_i | X_j = x_j \sim \text{Bin} \left(n - x_j, \frac{p_i}{1 - p_j} \right)$$

6. The conditional distribution of X_i given $T = X_i + X_j = t$, for $i \neq j$, is

$$X_i | X_i + X_j = t \sim \text{Bin} \left(t, \frac{p_i}{p_i + p_j} \right)$$

WE SHALL look at the bivariate normal distribution so that it is easier to be explained. The same idea can be extended to a multivariate Normal distribution.

Definition 49 (Bivariate Normal Distribution)

Let X_1 and X_2 be rvs with joint pdf

$$\begin{aligned} f(x_1, x_2) &= \frac{1}{2\pi |\Sigma|^{\frac{1}{2}}} \exp \left\{ -\frac{1}{2} (x - \mu)^T \Sigma^{-1} (x - \mu) \right\} \\ &= \frac{1}{2\pi \sigma_1 \sigma_2 \sqrt{1 - \rho^2}} \exp \left\{ -\frac{1}{2(1 - \rho^2)} \left[\frac{(x_1 - \mu_1)^2}{\sigma_1^2} \right. \right. \\ &\quad \left. \left. + \frac{(x_2 - \mu_2)^2}{\sigma_2^2} - \frac{2\rho(x_1 - \mu_1)(x_2 - \mu_2)}{\sigma_1 \sigma_2} \right] \right\} \end{aligned}$$

where $(x_1, x_2) \in \mathbb{R}^2$ and

$$x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \mu = \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}, \Sigma = \begin{bmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 \\ \rho\sigma_1\sigma_2 & \sigma_2^2 \end{bmatrix}$$

where Σ is a nonsingular matrix. Then $X = (X_1, X_2)^T$ is said to have a bivariate normal distribution, and we write $X \sim \text{BVN}(\mu, \Sigma)$.

Proposition 43 (Properties of Bivariate Normal Distribution)

Suppose $X \sim \text{BVN}(\mu, \Sigma)$, where

$$\mu = \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}, \Sigma = \begin{bmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 \\ \rho\sigma_1\sigma_2 & \sigma_2^2 \end{bmatrix}$$

1. X has a joint mgf of

$$M(t_1, t_2) = E[\exp(t^T X)] = E(e^{t_1 X_1 + t_2 X_2}) = \exp\left(\mu^T t + \frac{1}{2} t^T \Sigma t\right)$$

$$\forall (t_1, t_2) \in \mathbb{R}^2.$$

2. $X_1 \sim N(\mu_1, \sigma_1^2)$ and $X_2 \sim N(\mu_2, \sigma_2^2)$

3. $\text{Cov}(X_1, X_2) = \rho\sigma_1\sigma_2$ and $\text{Corr}(X_1, X_2) = \rho$ where $|\rho| \leq 1$

4. X_1 and X_2 are independent rvs iff $\rho = 0$

5. $c = (c_1, c_2)^T$ is a nonzero vector of constants \implies

$$c^T X = \sum_{i=1}^2 c_i X_i \sim N(c^T \mu, c^T \Sigma c).$$

6. If A is a 2×2 nonsingular matrix and b is a 2×1 vector, then $Y = AX + b \sim \text{BVN}(A\mu + b, A\Sigma A^T)$.

7.

$$X_2 | X_1 = x_1 \sim N\left(\mu_2 + \rho \frac{\sigma_2}{\sigma_1} (x_1 - \mu_1), \sigma_2^2 (1 - \rho^2)\right)$$

$$X_1 | X_2 = x_2 \sim N\left(\mu_1 + \rho \frac{\sigma_1}{\sigma_2} (x_2 - \mu_2), \sigma_1^2 (1 - \rho^2)\right)$$

8. $(X - \mu)^T \Sigma^{-1} (X - \mu) \sim \chi^2(2)$

Definition 50 (Random Sample (IID))

Rvs X_1, \dots, X_n are said to form a **simple random sample** or are said to be **independent and identically distributed (IID)** if X_1, \dots, X_n are independent, and $f_{X_i} = f_{X_j}, \forall i \neq j$.

Example 11.0.1 (Example 3.28)

Let X_1, \dots, X_{10} be a random sample of standard normal distribution.

Let Y_1 denote the number of these variables that are between -1 and 1 , let Y_2 denote the number that have absolute value between 1 and 2 , and let Y_3 denote the number that have absolute value larger than 2 .

Calculate:

1. $P(Y_1 \leq 2)$
2. $E[Y_2 \mid Y_1 = 5]$

12 Lecture 12 Jun 12th 2018

12.1 Functions of Random Variables

12.1.1 Transformation of Two or More Random Variables

In earlier lectures we discussed about basic transformations from one random variable to another, for example, from a continuous rv X to $Y = g(X)$. In particular, we methods were presented:

- THE DIRECT METHOD, i.e. $P(Y \leq y) = P(g(X) \leq y)$, and taking the derivative of $P(Y \leq y)$ with respect to y .
- USING THE MGF OF Y , and then translate it as the mgf of X .

“ Note (Recall)

In Section 2.4.4, we used the following idea to obtain the result that we desire: for rvs X and $Y = g(X)$ where g is some continuous and injective function

$$P(Y \leq y) = P(g(X) \leq y) = P(X \leq g^{-1}(y)) = F_X(g^{-1}(y))$$
$$f_Y(y) = \frac{d}{dy} F_X(g^{-1}(y)) = (g^{-1}(y))' f_X(g^{-1}(y))$$

In this chapter, we will now study the case where we have more than one rv involved. In particular, let X and Y be two continuous rvs with joint pdf $f(x, y)$. Our questions are:

1. What is the distribution of $U = h_1(X, Y)$?
2. What is the joint distribution of $U = h_1(X, Y)$ and $V = h_2(X, Y)$?

To answer the first question, we can actually still employ the direct method:

Example 12.1.1 (Example 4.1 (Course Notes 4.1.1))

Suppose X and Y are continuous rvs with joint pdf

$$f(x, y) = 3y\mathbb{1}_{0 \leq x \leq y \leq 1}$$

Find the pdf of $T = XY$.

 Solution

First, note that¹


$$P(T \leq t) = P(XY \leq t) = P\left(Y \leq \frac{t}{X}\right)$$

The diagram to the right shows us the support of the joint probability. We observe that if $t \leq 0$, then $P(T \leq t) = 0$, and if $t \geq 1$, then $P(T \leq t) = 1$. Now if $0 < t < 1$, the region that we are looking for is the shaded region with the label A, and so we consider

$$\begin{aligned} P(T \leq t) &= 1 - P(B) = 1 - \int \int_B f(x, y) dx dy \\ &\stackrel{(1)}{=} 1 - \int_{\sqrt{t}}^1 \int_{\frac{t}{y}}^{\sqrt{t}} f(x, y) dx dy - \int_{\sqrt{t}}^1 \int_{\sqrt{t}}^y f(x, y) dx dy \\ &\stackrel{(2)}{=} 1 - \int_{\sqrt{t}}^1 \int_{\frac{t}{y}}^y f(x, y) dx dy \\ &= 1 - \int_{\sqrt{t}}^1 \int_{\frac{t}{y}}^y 3y dx dy \\ &= 1 - \int_{\sqrt{t}}^1 3y \left(y - \frac{t}{y}\right) dy \quad \because \text{FTC} \\ &= 1 - \frac{\sqrt{t}}{1} \left(3y^2 - 3t\right) dy \\ &= 1 - \left[y^3 - 3ty\right]_{\sqrt{t}}^1 = 1 - (1 - 3t - \sqrt{t}^3 + 3t\sqrt{t}) \\ &= 3t - 2t\sqrt{t} \text{ for } 0 < t < 1 \end{aligned}$$

where for step (1), we broke B into two parts, in particular at $x = \sqrt{t}$ where $y = x$ and $y = \frac{t}{x}$ coincide, and step (2) is true by linearity of

¹ I wonder if the last step is actually valid. The support of X definitely includes 0, so the division would not make sense with $\frac{t}{X}$. We can, however, still make sense of the event in the 2nd term, which we would have $P(0 \leq t)$. Should we be concerned about $X = 0$, or can we neglect that single point given that X is a continuous rv?



integration. With that, i.e. with the CDF of T , we can then obtain


$$f_T(t) = \begin{cases} 3 - 3\sqrt{t} & 0 < t < 1 \\ 0 & \text{otherwise} \end{cases}$$

Example 12.1.2 (Example 4.2 (Course Note 4.1.2))

Using the info in Example 12.1.1, find the pdf of $T = \frac{X}{Y}$.

Solution

The diagram to the right shows the support of X, Y and the function $t = \frac{x}{y}$. We observe that if $t = 0$, then $x = 0$, and we would have the line on the axis, and so $P(T \leq t) = 0$. If $t < 0$, we would have $y = mx$ where $m = \frac{1}{t} < 0$, which, regardless of what $t < 0$ is, will not interact with the support of X and Y . So for $t < 0$, $P(T \leq t) = 0$. Now if $t > 0$, we have



$$P(T \leq t) = P\left(\frac{X}{Y} \leq t\right) = P\left(Y \geq \frac{1}{t}X\right).$$

Consider the case where $t \geq 1$, we have that the event would still cover the entire support set of X and Y , and so $P(T \leq t) = 1$ for $t \geq 1$. With that, the only remaining case is when $0 < t < 1$. In this case,

$$P(T \leq t) = \int_0^1 \int_0^{ty} 3y \, dx \, dy = \int_0^1 3ty^2 \, dy = t$$

and so the pdf of T is

$$f_T(t) = \begin{cases} 1 & 0 < t < 1 \\ 0 & \text{otherwise} \end{cases}$$

Example 12.1.3 (Example 4.3 (Course Note 4.1.3) - Order Statistics)

Suppose X_1, \dots, X_n are IID samples, each from a continuous distribution, and with pdf f and cdf F . Find the pdf of

1. $T = \min(X_1, \dots, X_n) = X_{(1)}$
2. $Y = \max(X_1, \dots, X_n) = X_{(n)}$

Solution

1. For T , we have that its cdf is²

² The use the **Law of Total Probability** here so that we can use the following argument: "the *smallest* rv is larger than t , and so rest of the rvs must be the same."

$$\begin{aligned}
P(T \leq t) &= 1 - P(T > t) = 1 - P(\min(X_1, \dots, X_n) > t) \\
&= 1 - P(X_1 > t, X_2 > t, \dots, X_n > t) \\
&= 1 - \prod_{i=1}^n P(X_i > t) \quad \because \text{independence} \\
&= 1 - \prod_{i=1}^n P(X_1 > t) \quad \because \text{identical distribution} \\
&= 1 - P(X_1 > t)^n = 1 - [1 - F_{X_1}(t)]^n
\end{aligned}$$

and so its pdf is

$$\begin{aligned}
f_T(t) &= -\frac{d}{dt}[1 - F_{X_1}(t)]^n = -n(-F_{X_1}(t))'[1 - F_{X_1}(t)]^{n-1} \\
&= nf_{X_1}(t)[1 - F_{X_1}(t)]^{n-1}.
\end{aligned}$$

Since T relies entirely on X_1 (due to IID), and since we did not have to condition on the values of t , we have that

$$\text{supp}(T) = \text{supp}(X_1) = \text{supp}(X_i) \quad \text{for } i = 1, \dots, n.$$

2. For Y , we have that its cdf is³

$$\begin{aligned}
P(Y \leq t) &= P(\max(X_1, \dots, X_n) \leq y) = P(X_1 \leq y, \dots, X_n \leq y) \\
&= \prod_{i=1}^n P(X_i \leq y) \quad \because \text{independence} \\
&= \prod_{i=1}^n P(X_1 \leq y) \quad \because \text{identical distribution} \\
&= P(X_1 \leq t)^n = F_{X_1}(t)^n
\end{aligned}$$

³ This time, we do not have to employ the **Law of Total Probability**, because we simply have that “the *largest* rv is smaller than t , and so must the rest of the rvs.”

and therefore its pdf is

$$f_Y(y) = \frac{d}{dy} F_{X_1}(y)^n = n f_{X_1}(y) F_{X_1}(y)^{n-1}.$$

Exercise 12.1.1

From Example 12.1.3, find the joint distribution of $X_{(1)}$ and $X_{(n)}$.

12.1.2 One-to-One Bivariate Transformations

Definition 51 (One-to-One Bivariate Transformation)

Let X and Y be rvs, and $R_{XY} = \text{supp}[(X, Y)] \in \mathbb{R}^2$. We define

$$U = h_1(X, Y) \quad V = h_2(X, Y)$$

$$S : R_{XY} \rightarrow \mathbb{R}^2 \text{ by } (x, y) \mapsto (h_1(x, y), h_2(x, y))$$

The mapping S is called a **one-to-one mapping** if and only if ⁴ $\forall (u, v) \in R_{UV}, \exists!(x, y) \in R_{XY}, \exists w_1, w_2$ that are functions such that

$$x = w_1(u, v) \quad y = w_2(u, v)$$

i.e. $\exists S^{-1} : R_{UV} \rightarrow R_{XY}$ such that $(u, v) \mapsto (x, y)$. The **Jacobian** of the transformation S^{-1} is

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \left| \frac{\partial(u, v)}{\partial(x, y)} \right|^{-1}$$

where $\frac{\partial(u, v)}{\partial(x, y)}$ is the Jacobian of the transformation S .

⁴ There is nothing magnificent about this definition, since this is simply the definition of a one-to-one function.

📌 Theorem 44 (One-to-One Bivariate Transformations)

Let X and Y be continuous rvs with joint pdf f_{XY} and let $R_{XY} = \{(x, y) : f(x, y) > 0\}$ be the support set of (X, Y) , and R_{UV} be the support set of (U, V) . Suppose the transformation $S : R_{XY} \rightarrow R_{UV}$ defined by

$$U = h_1(X, Y) \quad V = h_2(X, Y)$$

is a one-to-one transformation, with inverse transformation

$$X = w_1(U, V) \quad Y = w_2(U, V).$$

Then $g(u, v)$, the joint pdf of U and V , is given by

$$\forall (u, v) \in R_{UV} \quad g(u, v) = f(w_1(u, v), w_2(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right|.$$

This is a generalization of **📌 Theorem 8**.

✏ Proof

Let the inverse transformation be labelled as $S^{-1} : R_{UV} \supset B \rightarrow A \subset$

R_{XY} . Then

$$\begin{aligned} \int_B g(u, v) \, dv \, du &= P[(U, V) \in B] = P[(X, Y) \in A] \\ &= \int_A f(x, y) \, dx \, dy \\ &= \int_B f(w_1(u, v), w_2(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \, du \, dv \end{aligned}$$

where the last step is by the **Change of Variables Theorem**. And so by comparing integrands, we have

$$\forall (u, v) \in R_{UV} \quad g(u, v) = f(w_1(u, v), w_2(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right|$$

as required. \square

Proposition 45 (Properties of the Jacobian)

Given the setup in [Definition 51](#), we have that

1. if S is a linear transformation, i.e. $\exists a_1, b_1, c_1, a_2, b_2, c_2 \in \mathbb{R}$ such that $u(x, y) = a_1x + b_1y + c_1$ and $v(x, y) = a_2x + b_2y + c_2$, then the Jacobian is a constant;
2. if S is a one-to-one transformation, then $\left| \frac{\partial(x, y)}{\partial(u, v)} \right| \neq 0$

Proof

1. We have

$$\begin{aligned} \frac{\partial u}{\partial x} &= a_1 & \frac{\partial u}{\partial y} &= b_1 \\ \frac{\partial v}{\partial x} &= a_2 & \frac{\partial v}{\partial y} &= b_2 \end{aligned}$$

and so

$$|J| = \frac{\partial(u, v)}{\partial(x, y)} = \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} = a_1b_2 - a_2b_1$$

which is a constant, as required.

2. **I have no idea how to prove this.**

Example 12.1.4 (Example 4.4 (Course Notes 4.2.4))

Suppose $X \sim \text{Gam}(a, 1)$ and $Y \sim \text{Gam}(b, 1)$ independently. Find the joint pdf of $U = X + Y$ and $V = \frac{X}{X+Y}$. Show that $U \sim \text{Gam}(a + b, 1)$ and $V \sim \text{Beta}(a, b)$, independently. Find $E(V)$.

 Solution

Given $U = X + Y$ and $V = \frac{X}{X+Y}$, rearranging variables, we have

$$X = UV \text{ and } Y = U(1 - V)$$

In order for U to have a Gamma distribution, we need U to be non-negative, which we do since both X and Y have Gamma distributions. Note that the transformation is indeed one to one, since $\forall (u_1, v_1), (u_2, v_2) \in R_{UV}$ with


$$\begin{aligned} u_1 = x_1 + y_1 \quad v_1 &= \frac{x_1}{x_1 + y_1} \\ u_2 = x_2 + y_2 \quad v_2 &= \frac{x_2}{x_2 + y_2}, \end{aligned}$$

we have that, if we let ϕ denote the transformation,


$$\begin{aligned} \phi(u_1, v_1) &= \phi(u_2, v_2) \\ \implies \left(x_1 + y_1, \frac{x_1}{x_1 + y_1} \right) &= \left(x_2 + y_2, \frac{x_2}{x_2 + y_2} \right) \end{aligned}$$

which then

$$\begin{aligned} x_1 + y_1 &= x_2 + y_2 & (12.1) \\ \frac{x_1}{x_2 + y_2} &= \frac{x_2}{x_2 + y_2} \\ \xRightarrow{\text{Equation (12.1)}} x_1 &= x_2 \\ \xRightarrow{\text{Equation (12.1)}} y_1 &= y_2. \end{aligned}$$

We shall now get the Jacobian so that we may use  Theorem 44, so that we may consequently get the distributions for U and V .

$$\begin{aligned} J &= \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} v & u \\ 1-v & -u \end{vmatrix} = -vu - u(1-v) = -u \\ &|J| = u \end{aligned}$$

By  Theorem 44, and since X and Y are independent, we have

$$\begin{aligned}
 f_{U,V}(u,v) &= f_{X,Y}(x,y) \cdot |J| = f_X(x)f_Y(y) \cdot |J| \\
 &= \frac{x^{a-1}e^{-x}}{\Gamma(a)} \frac{y^{b-1}e^{-y}}{\Gamma(b)} \cdot |J| \\
 &= \frac{e^{-a}e^{-b}}{\Gamma(a)\Gamma(b)} (uv)^{a-1}u^{b-1}(1-v)^{b-1}u \\
 &= \underbrace{\frac{u^{a+b-1}e^{-(a+b)}}{\Gamma(a+b)}}_{\text{pdf of Gam}(a+b,1)} \cdot \underbrace{\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} v^{a-1}(1-v)^{b-1}}_{\text{pdf of Beta}(a,b)}
 \end{aligned}$$

We have already shown that U is a non-negative rv, and so $U \sim \text{Gam}(a+b,1)$ as required. Note that for V , we have $X+Y > X > 0 \implies 1 > \frac{X}{X+Y} > 0 \quad \because X+Y \neq 0$ and so $0 < V < 1$. Therefore $V \sim \text{Beta}(a,b)$.

With that, we can look for $E[V]$.

$$\begin{aligned}
 E[V] &= \int_0^1 v \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} v^{a-1}(1-v)^{b-1} dv \\
 &= \frac{\Gamma(a+b)\Gamma(a+1)}{\Gamma(a)\Gamma(a+b+1)} \int_0^1 \underbrace{\frac{\Gamma(a+b+1)}{\Gamma(a+1)\Gamma(b)} v^a(1-v)^{b-1} dv}_{\text{pdf of Beta}(a+1,b)} \\
 &= \frac{a\Gamma(a)\Gamma(a+b)}{(a+b)\Gamma(a)\Gamma(a+b)} = \frac{a}{a+b}
 \end{aligned}$$

13 Lecture 13 Jun 14th 2018

13.1 Functions of Random Variables (Continued)

13.1.1 One to One Bivariate Transformations (Continued)

Example 13.1.1 (Example 4.5 (Course Notes 4.2.8))

Suppose X and Y are continuous rvs with joint pdf

$$f(x, y) = e^{-x-y} \mathbb{1}_{\{0 < x, y < \infty\}}.$$

Let $U = X + Y$ and $V = X - Y$. Find the joint pdf of U and V . (Note: Be sure to specify the support of (U, V) .) Then, find the marginal pdf of U and V respectively.

Solution

Note that because $0 < x, y < \infty$,

$$\begin{aligned} u = x + y &\implies 0 < u < \infty \\ v = x - y &\implies -\infty < v < \infty \end{aligned}$$

The Jacobian is

$$\left| \frac{\partial(u, v)}{\partial(x, y)} \right| = \left| \begin{vmatrix} 1 & 1 \\ 1 & -1 \end{vmatrix} \right| = |-2| = 2$$

and so

$$\left| \frac{\partial(x, y)}{\partial(u, v)} \right| = \left| \frac{\partial(u, v)}{\partial(x, y)} \right|^{-1} = \frac{1}{2}.$$

Note that

$$X = \frac{U - V}{2} \quad Y = \frac{U + V}{2}.$$

and so

$$x = \frac{u+v}{2} > 0 \implies u > -v$$

$$y = \frac{u-v}{2} > 0 \implies u > v.$$

The diagram to the right shows the support of U, V . With that,

$$g(u, v) = f\left(\frac{u+v}{2}, \frac{u-v}{2}\right) \cdot \frac{1}{2}$$

$$= \frac{1}{2} e^{-\left(\frac{u+v}{2}\right) - \left(\frac{u-v}{2}\right)} = \frac{1}{2} e^{-u}.$$


With that, the marginal pdf of V is¹

$$v < 0 \implies f_V(v) = \int_{-v}^{\infty} \frac{1}{2} e^{-u} du = \frac{1}{2} e^v$$

$$v \geq 0 \implies f_V(v) = \int_v^{\infty} \frac{1}{2} e^{-u} du = \frac{1}{2} e^{-v}.$$

For the marginal pdf of U , we have

$$f_U(u) = \int_{-u}^u \frac{1}{2} e^{-u} dv = u e^{-u} \quad \text{for } 0 < u \leq \infty.$$



¹ V is also called the Double Exponential Distribution.

13.1.2 Moment Generating Function Method

This method is particularly useful in finding distributions of sums of independent rvs.

Theorem 46 (Sums of MGF)

Suppose X_1, \dots, X_n are independent rvs and X_i has mgf $M_i(t)$ which exists for $t \in (-h, h)$ for some $H > 0$. The mgf of $Y = \sum_{i=1}^n X_i$ is given by

$$M_Y(t) = \prod_{i=1}^n M_i(t)$$

for $t \in (-h, h)$.

 **Proof**

Observe that

$$\begin{aligned} M_Y(t) &= E \left[e^{tY} \right] = E \left[e^{t \sum_{i=1}^n X_i} \right] = E \left[\prod_{i=1}^n e^{tX_i} \right] \\ &= \prod_{i=1}^n E \left[e^{tX_i} \right] = \prod_{i=1}^n M_i(t) \end{aligned}$$

and $t \in (-h, h)$ is preserved. □

“ Note

1. If X_i 's are IID rvs each with mgf $M(t)$ then Y has mgf

$$M_Y(t) = [M(t)]^n \quad \text{for } t \in (-h, h).$$

2. Used in conjunction with the Uniqueness Theorem for mgfs, this theorem can be used to find the distribution of Y .

Exercise 13.1.1

Show the following results:

1. If $X \sim \text{Gam}(\alpha, \beta)$, where $\alpha \in \mathbb{N}$, then $\frac{2X}{\beta} \sim \chi^2(2\alpha)$.

2. If $X_i \sim \text{Gam}(\alpha_i, \beta)$, $i = 1, \dots, n$ independently, then

$$\sum_{i=1}^n X_i \sim \text{Gam} \left(\sum_{i=1}^n \alpha_i, \beta \right).$$

3. If $X_i \sim \text{Gam}(1, \beta) = \text{Exp}(\beta)$, $i = 1, \dots, n$ independently, then

$$\sum_{i=1}^n X_i \sim \text{Gam}(n, \beta).$$

4. If $X_i \sim \text{Gam} \left(\frac{k_i}{2}, 2 \right) = \chi^2(k_i)$, $i = 1, \dots, n$ independently, then

$$\sum_{i=1}^n X_i \sim \chi^2 \left(\sum_{i=1}^n k_i \right).$$

5. If $X_i \sim N(\mu, \sigma^2)$, $i = 1, \dots, n$ independently, then

$$\sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma} \right)^2 \sim \chi^2(n).$$

6. If $X_i \sim \text{Poi}(\mu_i)$, $i = 1, \dots, n$ independently, then

$$\sum_{i=1}^n X_i \sim \text{Poi} \left(\sum_{i=1}^n \mu_i \right).$$

7. If $X_i \sim \text{Bin}(n_i, p)$, $i = 1, \dots, n$ independently, then

$$\sum_{i=1}^n X_i \sim \text{Bin} \left(\sum_{i=1}^n n_i, p \right).$$


8. If $X_i \sim \text{NB}(k_i, p)$, $i = 1, \dots, n$ independently, then

$$\sum_{i=1}^n X_i \sim \text{NB} \left(\sum_{i=1}^n k_i, p \right).$$

Solution

1. Using the mgf method,

$$\begin{aligned} M_Y(t) &= E[e^{tY}] = E \left[e^{t \cdot \frac{2X}{\beta}} \right] = \int_0^\infty e^{\frac{2tx}{\beta}} \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-\frac{x}{\beta}} dx \\ &= \frac{1}{\beta^\alpha} \int_0^\infty \frac{1}{\Gamma(\alpha)} x^{\alpha-1} e^{-\frac{x(1-2t)}{\beta}} dx \\ &\stackrel{(*)}{=} \frac{1}{\beta^\alpha} \left(\frac{\beta}{1-2t} \right)^\alpha \underbrace{\int_0^\infty \frac{1}{\left(\frac{\beta}{1-2t} \right) \Gamma(\alpha)} x^{\alpha-1} e^{-\frac{x}{\frac{\beta}{1-2t}}} dx}_{\text{pdf of } \text{Gam} \left(\alpha, \frac{\beta}{1-2t} \right)} \\ &= \left(\frac{1}{1-2t} \right)^\alpha = (1-2t)^{-\alpha} \end{aligned}$$

where in (*) we note that $1-2t > 0$ and so $t < \frac{1}{2}$. Observe that the mgf of Y is the mgf of $\chi^2(2\alpha)$ and so by the  Theorem 20, $Y = \frac{2X}{\beta} \sim \chi^2(2\alpha)$.

2. Using the mgf method, let $Y = \sum_{i=1}^n X_i$

$$\begin{aligned} M_Y(t) &= E[e^{tY}] = E \left[\prod_{i=1}^n e^{tX_i} \right] = \prod_{i=1}^n E[e^{tX_i}] \\ &= \prod_{i=1}^n M_{X_i}(t) = \prod_{i=1}^n (1-\beta t)^{-\alpha_i} = (1-\beta t)^{-\sum_{i=1}^n \alpha_i} \end{aligned}$$

and we observe that the last term is the mgf of $\text{Gam}(\sum_{i=1}^n \alpha_i, \beta)$ and so the result follows.

3. From (2) is $\alpha_i = 1$ for $i = 1, \dots, n$, then $\sum_{i=1}^n \alpha_i = \sum_{i=1}^n 1 = n$. The result follows.

4. By (2), we have $\sum_{i=1}^n X_i \sim \text{Gam}\left(\sum_{i=1}^n \frac{k_i}{2}, 2\right)$. Then by (1), we have that

$$\sum_{i=1}^n X_i \sim \chi^2\left(\sum_{i=1}^n k_i\right)$$

as required.

5. We know that $Y_i = \frac{X_i - \mu}{\sigma} \sim Z(0, 1)$. Let $W_i = Y_i^2$. Then

$$\begin{aligned} M_{W_i}(t) &= E\left[e^{tY_i}\right] = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{y^2}{2}} e^{ty^2} dy \\ &= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left\{-\frac{y^2}{2(1-2t)^{-1}}\right\} dy \\ &= (1-2t)^{-\frac{1}{2}} \int_{-\infty}^{\infty} \underbrace{\frac{1}{\sqrt{2\pi}(1-2t)^{-\frac{1}{2}}} \exp\left\{\frac{y^2}{2(1-2t)^{-1}}\right\}}_{\text{pdf of } N\left(0, (1-2t)^{-\frac{1}{2}}\right)} dy \\ &= (1-2t)^{-\frac{1}{2}}, \end{aligned}$$

which is the mgf of $\chi^2(1)$. Let $W = \sum_{i=1}^n W_i$. Then

$$M_W(t) = \prod_{i=1}^n (1-2t)^{-\frac{1}{2}} = (1-2t)^{-\frac{n}{2}}$$

which is the mgf of $\chi^2(n)$.

6. Let $Y = \sum_{i=1}^n X_i$. Then

$$M_Y(t) = \prod_{i=1}^n e^{\mu_i(e^t - 1)} = \exp\left[(e^t - 1) \sum_{i=1}^n \mu_i\right]$$

is the mgf of $\text{Poi}(\sum_{i=1}^n \mu_i)$.

7. Let $Y = \sum_{i=1}^n X_i$. Then

$$M_Y(t) = \prod_{i=1}^n (pe^t + q)^{n_i} = (pe^t + q)^{\sum_{i=1}^n n_i}$$

is the mgf of $\text{Bin}(\sum_{i=1}^n n_i, p)$.

14 Lecture 14 Jun 19th 2018

14.1 Functions of Random Variables (Continued 2)

14.1.1 Moment Generating Function Method (Continued)

Theorem 47 (Gaussian Distribution)


If $X_i \sim N(\mu_i, \sigma_i^2)$, with $i = 1, \dots, n$ independently, then

$$\sum_{i=1}^n a_i X_i \sim N\left(\sum_{i=1}^n a_i \mu_i, \sum_{i=1}^n a_i^2 \sigma_i^2\right).$$

Proof

Let $Y_i = a_i X_i$. Then by  Theorem 19, we have

$$M_{Y_i}(t) = M_{X_i}(a_i t) = \exp\left[\mu_i a_i t + \frac{\sigma_i^2 a_i^2 t^2}{2}\right],$$

which implies that $Y_i \sim N(a_i \mu_i, a_i^2 \sigma_i^2)$ by  Theorem 20. Then let $Y = \sum_{i=1}^n Y_i$. Then

$$M_Y(t) = \prod_{i=1}^n \exp\left[\mu_i a_i t + \frac{\sigma_i^2 a_i^2 t^2}{2}\right] = \exp\left[t \sum_{i=1}^n a_i \mu_i + \frac{t^2}{2} \sum_{i=1}^n a_i^2 \sigma_i^2\right],$$

which implies that $Y \sim N\left(\sum_{i=1}^n a_i \mu_i, \sum_{i=1}^n a_i^2 \sigma_i^2\right)$. □



Theorem 48 (Properties of the Gaussian Distribution)

Assume $X_1, \dots, X_n \sim N(\mu, \sigma^2)$, independently, where $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$,

and $S^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}$. Then

1. $\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$.
2. (**Cochran's Theorem**) $\frac{(n-1)S^2}{\sigma^2} = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{\sigma^2} \sim \chi^2(n-1)$.
3. (**t-test**) $\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$.

Proof

1. Let $Y_i = \frac{X_i}{n}$.  Theorem 47 implies that $Y_i \sim N\left(\frac{\mu}{n}, \frac{\sigma^2}{n^2}\right)$.
Let $Y = \sum_{i=1}^n Y_i$. Then  Theorem 47 implies that $Y \sim N\left(\sum_{i=1}^n \frac{\mu}{n}, \sum_{i=1}^n \frac{\sigma^2}{n^2}\right) = N\left(\mu, \frac{\sigma^2}{n}\right)$. □
2. The proof of Cochran's Theorem is beyond the scope of this course, for it uses knowledge from linear algebra and involving Fourier Transforms. [Reference - Wikipedia]
3. The proof of this statement is non-trivial: [Reference - Wikipedia]
[Reference - Math SE]

Theorem 49 (F Distribution)

Suppose X_1, \dots, X_n is a random sample from the $N(\mu_1, \sigma_1^2)$ distribution and independently Y_1, \dots, Y_m is a random sample from the $N(\mu_2, \sigma_2^2)$ distribution. Let

$$S_1^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 \text{ and } S_2^2 = \frac{1}{m-1} \sum_{i=1}^m (Y_i - \bar{Y})^2.$$

Then

$$\frac{S_1^2/\sigma_1^2}{S_2^2/\sigma_2^2} \sim F(n-1, m-1).$$

Proof

This is, once again, an incomplete proof. Rigor shall be appended where it is due or they shall be stated with reference, or if it is at a level that I cannot understand and that I cannot find a reference, I shall state that as well.

Note that the definition of the F distribution is as follows:

$$F_{\gamma_1, \gamma_2} = \frac{\chi^2(\gamma_1)/\gamma_1}{\chi^2(\gamma_2)/\gamma_2}$$

where $\chi^2(\gamma_1)$ and $\chi^2(\gamma_2)$ are independent.

Now we are given that X_1, \dots, X_n and Y_1, \dots, Y_m are independent of one another. It can likely be shown, using induction, that the following two are independent of one another:

$$\sum_{i=1}^n \left(\frac{X_i - \bar{X}}{\sigma_1} \right)^2 \sim \chi^2(n-1)$$

$$\sum_{j=1}^m \left(\frac{Y_j - \bar{Y}}{\sigma_2} \right)^2 \sim \chi^2(m-1).$$

Then

$$\frac{\sum_{i=1}^n \left(\frac{X_i - \bar{X}}{\sigma_1} \right)^2}{\sum_{j=1}^m \left(\frac{Y_j - \bar{Y}}{\sigma_2} \right)^2} = \frac{(n-1)S_1^2/\sigma_1^2}{(m-1)S_2^2/\sigma_2^2}.$$

With that, we have that

$$\frac{S_1^2/\sigma_1^2}{S_2^2/\sigma_2^2} = \frac{\frac{(n-1)S_1^2}{\sigma_1^2}/(n-1)}{\frac{(m-1)S_2^2}{\sigma_2^2}/(m-1)}.$$

The result follows.

15 Lecture 15 Jun 21st 2018

15.1 Limiting or Asymptotic Distributions

15.1.1 Convergence in Distribution

Definition 52 (Convergence in Distribution)

Let $X_1, X_2, \dots, X_n, \dots$ be a sequence of rvs such that X_i has cdf F_i , for $i \in \mathbb{N}$. Let X be an rv with cdf F . We say that X_i **converges in distribution** to X and we write

$$X_n \xrightarrow{D} X$$

if

$$\lim_{i \rightarrow \infty} F_i(x) = F(x)$$

at all points x at which F is continuous. We call F the **limiting** or **asymptotic distribution** of X_i .

The following theorem, that we shall not prove, will be useful for the remainder of the notes.

Theorem 50 (Taylor Series with Lagrange's Remainder)

Suppose that $f : [a, b] \rightarrow \mathbb{R}$ is infinitely differentiable, and $x \in [a, b]$. Then $\forall x \in [a, b]$ and $\forall k \in \mathbb{N}$,

$$f(x) = \sum_{i=0}^k \frac{f^{(i)}(c)(x-c)^i}{i!} + \frac{f^{(k+1)}(\zeta_x)(x-c)^{k+1}}{(k+1)!}$$

for some $\zeta_x \in [c, x]$.

The proof for the above involves the use of the **Mean Value Theorem for Integrals**.

 **Theorem 51 (Generalized Limit Definition of e)**

If $b, c \in \mathbb{R}$ are constants and $\lim_{n \rightarrow \infty} \psi(n) = 0$, then

$$\lim_{n \rightarrow \infty} \left[1 + \frac{b}{n} + \frac{\psi(n)}{n} \right]^{cn} = e^{bc}.$$

 **Proof**


The above equation can be rewritten as

$$\lim_{n \rightarrow \infty} e^{cn \log \left(1 + \frac{b}{n} + \frac{\psi(n)}{n} \right)} = e^{bc}$$

and so it suffices to prove that

$$\lim_{n \rightarrow \infty} cn \log \left(1 + \frac{b}{n} + \frac{\psi(n)}{n} \right) = bc$$

Note that the Taylor Expansion with Lagrange's Remainders

( **Theorem 50**) of $\log(1+x)$, where we pick $c = 0$ and $k = 1$ for convenience,

$$\log(1+x) = \frac{\log(1)x^0}{0!} + \frac{\left(\frac{1}{1-0}\right)x^1}{1!} + \frac{-\frac{1}{(1+\zeta_x)^2}x^2}{2!} = x - \frac{x^2}{2(1+\zeta_x)}$$

where $\zeta \in [0, x]$. Then

$$cn \log \left[1 + \frac{b}{n} + \frac{\psi(n)}{n} \right] = cb + c\psi(n) - \frac{c(b + \psi(n))^2}{2n(1 + \zeta)^2}$$

where $\zeta \in \left[0, \frac{b + \psi(n)}{n} \right]$. Note that by **L'Hôpital's Rule**, we have that

$$\lim_{n \rightarrow \infty} \frac{b + \psi(n)}{n} = \lim_{n \rightarrow \infty} \frac{\psi'(n)}{1} = 0$$

and so the possible highest value for ζ goes to 0 as $n \rightarrow \infty$. Then

$$\lim_{n \rightarrow \infty} \frac{c(b + \psi(n))^2}{2n(1 + \zeta)^2} = 0.$$

As a result, we have

$$\lim_{n \rightarrow \infty} cn \log \left(1 + \frac{b}{n} + \frac{\psi(n)}{n} \right) = bc$$

as required. \square

✦ **Corollary 52 (Limit definition of e)**

If $b, c \in \mathbb{R}$ are constants, then

$$\lim_{n \rightarrow \infty} \left(1 + \frac{b}{n} \right)^{cn} = e^{bc}$$

Example 15.1.1 (Example 5.1 (Course Notes 5.1.4))

Let $X_i \sim \text{Exp}(1)$, where $i \in \mathbb{N}$, independently so. Consider the sequence of rvs $Y_1, Y_2, \dots, Y_n, \dots$, where $Y_n = \max(X_1, \dots, X_n) - \log n$. Find the limiting distribution of Y_n .

 **Solution**

Firstly, observe that to find the support set of Y_n , note

$$\begin{aligned} 0 &< x_1, \dots, x_n < \infty \\ 0 &< \max(x_1, \dots, x_n) < \infty \\ -\log n &< \max(x_1, \dots, x_n) - \log n < \infty \end{aligned}$$

so

$$\text{supp}(Y) = (-\log n, \infty)$$

Now the pf of Y_n is

$$\begin{aligned} F_n(y) &= P(Y_n \leq y) = P(\max(X_1, \dots, X_n) - \log n \leq y) \\ &= P(\max(X_1, \dots, X_n) \leq y + \log n) \\ &= \prod_{i=1}^n P(X_i \leq y + \log n) \quad \because X_1, \dots, X_n \text{ are IID} \\ &= \prod_{i=1}^n \left[1 - e^{-y - \log n} \right] = \prod_{i=1}^n \left[1 - \frac{e^{-y}}{n} \right] = \left[1 - \frac{e^{-y}}{n} \right]^n. \end{aligned}$$

Thus the limiting distribution of Y_i is

$$\lim_{n \rightarrow \infty} F_n(y) = \lim_{n \rightarrow \infty} \left[1 - \frac{e^{-y}}{n} \right]^n = e^{-e^{-y}} \text{ for } y \in (-\log n, \infty).$$

Example 15.1.2 (Example 5.2 (Course Notes 5.1.5))

Let $X_i \sim \text{Unif}(0, \theta)$, $i \in \mathbb{N}$, independently so. Consider the sequence of random variables $Y_1, Y_2, \dots, Y_n, \dots$, where $Y_n = \max(X_1, \dots, X_n)$. Find the limiting distribution of Y_n .

 Solution

Clearly, support of $Y_n = (0, \theta)$. Note that

$$f_{X_i}(x) = \frac{1}{\theta} \mathbb{1}_{0 < x < \theta}.$$

Now since X_i are IID,

$$F_n(y) = P(Y_n \leq y) = \prod_{i=1}^n P(X_i \leq y) = \prod_{i=1}^n \frac{y}{\theta} = \left(\frac{y}{\theta}\right)^n \quad y \in (0, \theta)$$

Then the limiting distribution is

$$\lim_{n \rightarrow \infty} F_n(y) = \lim_{n \rightarrow \infty} \left(\frac{y}{\theta}\right)^n = \begin{cases} 0 & y < \theta \\ 1 & y \geq \theta \end{cases}$$

Define Y to have a distribution such that $P(Y = \theta) = 1$. Then

$$Y_n \xrightarrow{D} Y.$$

 Definition 53 (Degenerate Distribution)

A function F is the cdf of a degenerate distribution at value $y = c$ if

$$F(y) = \begin{cases} 0 & y < c \\ 1 & y \geq c \end{cases}.$$

In other words, F is the CDF of a discrete distribution where

$$P(Y = y) = \begin{cases} 1 & y = c \\ 0 & \text{otherwise} \end{cases}$$

We have that the earlier example gives us that the limiting distribution is a degenerate distribution.

15.1.2 Convergence in Probability

Definition 54 (Convergence in Probability)

A sequence of rvs $X_1, X_2, \dots, X_n, \dots$ **converges in probability** to an rv X if, for every $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} P(|X_n - X| \geq \varepsilon) = 0$$

or equivalently

$$\lim_{n \rightarrow \infty} P(|X_n - X| < \varepsilon) = 1$$

We write

$$X_n \xrightarrow{P} X.$$

Example 15.1.3 (Example 5.3)

Consider $X \sim \text{Bernoulli}(0.3)$. Define the sequence $X_n = \left(1 + \frac{1}{n}\right) X$, $n \in \mathbb{N}$. Show that $X_n \xrightarrow{P} X$.

Solution

Note that

$$P(X = x) = \begin{cases} 0.3 & x = 1 \\ 0.7 & x = 0 \end{cases}$$

Since $X_n = \left(1 + \frac{1}{n}\right) X$, we have

$$X_1 = 2X, X_2 = \frac{3}{2}X, X_3 = \frac{4}{3}X, \dots$$

Note that $|X_n - X| = \left| \left(1 + \frac{1}{n}\right) X - X \right| = \left| \frac{1}{n} X \right| = \frac{1}{n} X$, so

$$\begin{aligned} P(|X_n - X| < \varepsilon) &= P\left(\frac{1}{n} X < \varepsilon\right) = P(X < n\varepsilon) \\ &= \begin{cases} P(X = 0) = 0.7 & n < \frac{1}{\varepsilon} \\ P(X = 1) + P(X = 0) = 1 & n \geq \frac{1}{\varepsilon} \end{cases} \end{aligned}$$

Therefore

$$\lim_{n \rightarrow \infty} P(|X_n - X| < \varepsilon) = 1$$

and so

$$X_n \xrightarrow{P} X.$$

We shall look into the following proposition in the next lecture.

♦ **Proposition**

Suppose $\forall n \in \mathbb{N}$, $\{X_n\}$ is a sequence of rvs. Then

$$X_n \xrightarrow{P} X \implies X_n \xrightarrow{D} X$$

where X is an rv.

16 Lecture 16 Jun 26th 2018

16.1 Limiting or Asymptotic Distributions (Continued)

16.1.1 Convergence in Probability (Continued)

Before proving the proposition introduced in last class, we require the following lemma.

🌲 Lemma 53

Let X, Y be rvs, $a \in \mathbb{R}$, and $\varepsilon > 0$. Then

$$P(Y \leq a) \leq P(X \leq a + \varepsilon) + P(|Y - X| > \varepsilon)$$

✎ Proof

Note that

$$P(Y \leq a) = P(Y \leq a, X \leq a + \varepsilon) + P(Y \leq a, X > a + \varepsilon) \quad (16.1)$$

$$\leq P(X \leq a + \varepsilon) + P(Y - X \leq a - X, a - X < -\varepsilon) \quad (16.2)$$

$$\leq P(X \leq a + \varepsilon) + P(Y - X < -\varepsilon) \quad (16.3)$$

$$\leq P(X \leq a + \varepsilon) + P(Y - X < -\varepsilon) + P(Y - X > \varepsilon) \quad (16.4)$$

$$= P(X \leq a + \varepsilon) + P(|Y - X| > \varepsilon)$$

where Equation (16.1) is by **Law of Total Probability**, Equation (16.2) and Equation (16.3) are by the fact that for non-empty sets A and B , $P(A \cap B) \leq P(A)$, and Equation (16.4) is because $P(Y - X > \varepsilon) \geq 0$. □

♦ **Proposition 54 (Convergence in Probability Implies Convergence in Distribution)**

Suppose $\forall n \in \mathbb{N}$, $\{X_n\}$ is a sequence of rvs. Then

$$X_n \xrightarrow{P} X \implies X_n \xrightarrow{D} X$$

where X is an rv.

This only states for the case of scalar random variables.

✎ **Proof**

By Lemma 53, we have that¹

$$\begin{aligned} P(X_n \leq a) &\leq P(X \leq a + \varepsilon) + P(|X_n - X| > \varepsilon) \\ P(X \leq a - \varepsilon) &\leq P(X_n \leq a) + P(|X_n - X| > \varepsilon) \end{aligned}$$

Then

$$\begin{aligned} P(X \leq a - \varepsilon) - P(|X_n - X| > \varepsilon) &\leq P(X_n \leq a) \\ &\leq P(X \leq a + \varepsilon) + P(|X_n - X| > \varepsilon) \end{aligned}$$

As $n \rightarrow \infty$, by assumption that $X_n \xrightarrow{P} X$, we have that $P(|X_n - X| > \varepsilon) \rightarrow 0$, and so

$$P(X \leq a - \varepsilon) \leq \lim_{n \rightarrow \infty} P(X_n \leq a) \leq P(X \leq a + \varepsilon)$$

Now as $\varepsilon \rightarrow 0^+$, note that we must have the cdf of X be continuous on a by assumption, and so

$$P(X \leq a) \leq \lim_{n \rightarrow \infty} P(X_n \leq a) \leq P(X \leq a)$$

and so by the **Squeeze Theorem**,

$$\lim_{n \rightarrow \infty} P(X_n \leq a) = P(X \leq a)$$

which then

$$X_n \xrightarrow{D} X$$

as required. □

¹ We choose to use Lemma 53 to have $P(X_n \leq a)$ in two inequalities. Note that our goal is to show that

$$\lim_{n \rightarrow \infty} P(X_n \leq a) = P(X \leq a).$$

We already know that $\varepsilon \rightarrow 0$, so we should use that to our advantage in the case of X . Also, in hindsight, it is clear why we choose this method as the **Squeeze Theorem** is suitable to help us reach our conclusion.

Definition 55 (Convergence in Probability to a Constant)

A sequence of rvs $\{X_i\}_{i \in \mathbb{N}}$ converges in probability to a constant $b \in \mathbb{R}$ if $\forall \varepsilon > 0$,

$$\lim_{n \rightarrow \infty} P(|X_n - b| \geq \varepsilon) = 0$$

or equivalently

$$\lim_{n \rightarrow \infty} P(|X_n - b| < \varepsilon) = 1.$$

We write

$$X_n \xrightarrow{P} b.$$

Example 16.1.1 (Example 5.4 - Weak Law of Large Numbers)

Let X_1, \dots, X_n, \dots be a sequence of IID rvs, each having mean μ and variance σ^2 . Let

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i.$$

Then $\bar{X}_n \xrightarrow{P} \mu$.

Solution

Observe that

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n E[X_i] = \frac{n\mu}{n} = \mu,$$

Also, we have that

$$\text{Var}(\bar{X}_n) = \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n^2} \sum_{i=1}^n \text{Var}(X_i) = \frac{1}{n^2} \cdot n\sigma^2 = \frac{\sigma^2}{n}.$$

Then by [Theorem 16](#), we have that $\forall \varepsilon > 0$,

$$P(|\bar{X}_n - \mu| \geq \varepsilon) \leq \frac{\text{Var}(\bar{X}_n)}{\varepsilon^2} = \frac{\sigma^2}{n\varepsilon^2}.$$

Therefore, we have

$$0 \leq P(|\bar{X}_n - \mu| \geq \varepsilon) \leq \min\left(1, \frac{\sigma^2}{n\varepsilon^2}\right)$$

As $n \rightarrow \infty$, by [Squeeze Theorem](#), we have that

$$\lim_{n \rightarrow \infty} P(|\bar{X}_n - \mu| \geq \varepsilon) = 0$$

The Weak Law of Large Numbers is also known as [Bernoulli's Theorem](#).

i.e. $\bar{X}_n \xrightarrow{P} \mu$.

Note that the converse of  Proposition 54 is not generally true.

(Example required) However, with an additional condition, the converse becomes true.

 **Proposition 55 (Partial Converse of  Proposition 54)**

Suppose $\forall n \in \mathbb{N}$, $\{X_n\}$ is a sequence of rvs, each with cdf F_n . If

$$\lim_{n \rightarrow \infty} F_n(x) = \lim_{n \rightarrow \infty} P(X_n \leq x) = \begin{cases} 0 & x < c \\ 1 & x > c \end{cases},$$

then

$$X_n \xrightarrow{P} c.$$

 **Proof**

Note that for $\varepsilon > 0$, we have

$$\begin{aligned} P(|X_n - c| > \varepsilon) &= P(X_n - c < -\varepsilon) + P(X_n - c > \varepsilon) \\ &= P(X_n < c - \varepsilon) + P(X_n > c + \varepsilon) \\ &= 1 - P(X_n < c + \varepsilon) + P(X_n < c - \varepsilon) \end{aligned}$$


By assumption, we have that

$$\begin{aligned} \lim_{n \rightarrow \infty} P(X_n \leq c + \varepsilon) &= 1 \\ \lim_{n \rightarrow \infty} P(X_n \leq c - \varepsilon) &= 0 \end{aligned}$$

So

$$\lim_{n \rightarrow \infty} P(|X_n - c| > \varepsilon) = 1 - 1 + 0 = 0$$

and so by definition, we have $X_n \xrightarrow{P} c$. □

 **Definition 56 (Double Parameter Exponential Distribution)**

We say that an rv $X \sim \text{Exp}(\lambda, \theta)$ when X has pdf

$$f(x) = e^{-\frac{x-\theta}{\lambda}} \text{ for } x \in (\theta, \infty)$$

where λ is the scale parameter, and θ the location parameter.

Example 16.1.2 (Example 5.5 (Course Notes 5.2.5))

Let $X_i \sim \text{Exp}(1, \theta)$, $i = 1, 2, \dots$, independently. Consider the sequence of rvs Y_1, Y_2, \dots where $Y_n = \min(X_1, X_2, \dots, X_n)$, $n = 1, 2, \dots$. Show that $Y_n \xrightarrow{P} \theta$.

✎ Solution

Since we want to show that $Y_n \xrightarrow{P} \theta$, and θ is a constant, we can use

♠ Proposition 55. Thus we need to show that

$$\lim_{n \rightarrow \infty} F_{Y_n}(y) = \lim_{n \rightarrow \infty} P(Y_n \leq y) = \begin{cases} 0 & y < \theta \\ 1 & y > \theta \end{cases}$$

Since Y_n is defined as the minimum of n of the first X_i rvs, we need to use the **Law of Total Probability** in order to be able to make sense of the order statistics, i.e.

$$P(Y_n \leq y) = 1 - P(Y_n > y)$$

Now

$$\begin{aligned} P(Y_n > y) &= \prod_{i=1}^n P(X_i > y) = \prod_{i=1}^n \left[\int_y^{\infty} e^{-(x-\theta)} dx \right] \\ &= \prod_{i=1}^n \left[e^{-(y-\theta)} \right] = e^{-n(y-\theta)} \end{aligned}$$

Thus

$$\begin{aligned} \lim_{n \rightarrow \infty} P(Y_n \leq y) &= \begin{cases} 1 - \lim_{n \rightarrow \infty} P(Y_n > y) & y > \theta \\ 0 & y \leq \theta \end{cases} \\ &= \begin{cases} 1 - \lim_{n \rightarrow \infty} e^{-n(y-\theta)} = 1 & y > \theta \\ 0 & y \leq \theta \end{cases} \end{aligned}$$

since if $y < \theta$

$$P(Y_n \leq y) = P(\min(X_1, X_2, \dots, X_n) \leq y) = 0.$$

Note that if $y = \theta$, then

$$e^{-n(y-\theta)} = 1.$$

The proof is complete with \spadesuit Proposition 55.

16.1.2 Limit Theorems

\spadesuit Proposition 56 (Convergence in Distribution and MGF)

Let $X_1, X_2, \dots, X_n, \dots$ be a sequence of rvs such that X_n has mgf $M_n(t)$.

Let X be an rv with mgf $M(t)$.

$$X_n \xrightarrow{D} X \iff \left[\exists h > 0 \forall t \in (-h, h) \quad \lim_{n \rightarrow \infty} M_n(t) = M(t) \right]$$

\pencil Proof

Note that

$$\lim_{n \rightarrow \infty} M_n(t) = \lim_{n \rightarrow \infty} \int_{\text{supp}(X_n)} e^{tx} \frac{d}{dx} F_{X_n}(x) dx$$

and

$$M(t) = \int_{\text{supp}(X)} e^{tx} \frac{d}{dx} F_X(x) dx$$

The result follows assuming that the integral converges².

\square ² This allows us to “swap” the limit, the integral sign, and the differential operator.

Example 16.1.3

Consider the sequence Y_1, Y_2, \dots , where $Y_i \sim \text{Bin}\left(n, \frac{\mu}{n}\right)$, for $i = 1, 2, \dots$. Find the limiting distribution of Y_n .

 Solution (Example 5.6)

Note that $Y_n \sim \text{Bin}\left(n, \frac{\mu}{n}\right) \implies$

$$M_{Y_n}(t) = \left(\frac{\mu}{n}e^t + 1 - \frac{\mu}{n}\right)^n = \left(1 + \frac{\mu(e^t - 1)}{n}\right)^n.$$

So

$$\lim_{n \rightarrow \infty} M_{Y_n}(t) = \lim_{n \rightarrow \infty} \left(1 + \frac{\mu(e^t - 1)}{n}\right)^n = e^{\mu(e^t - 1)}$$

which is the mgf of a Poisson Distribution. Thus by  Proposition 56, we have that

$$Y_n \xrightarrow{D} Y \sim \text{Poi}(\mu).$$

Example 16.1.4 (Example 5.7)

Let $Y_1, Y_2, \dots \sim G(\mu, \sigma)$. Then

$$\frac{\bar{Y}_n - \mu}{\sigma/\sqrt{n}} \sim G(0, 1)$$


where $\bar{Y}_n = \frac{1}{n} \sum_{i=1}^n Y_i$.

 Solution

Note that by  Theorem 19, we have that

$$M_{\bar{Y}_n}(t) = \prod_{i=1}^n M_{Y_i}\left(\frac{t}{n}\right) = \prod_{i=1}^n e^{\frac{\mu t}{n} + \frac{(\sigma t)^2}{2n^2}} = e^{\mu t + \frac{\sigma^2 t^2}{n}}.$$

which is the mgf of $G\left(\mu, \frac{\sigma}{\sqrt{n}}\right)$. Let $X_n = \frac{\bar{Y}_n - \mu}{\sigma/\sqrt{n}}$.

Then again, by  Theorem 19, we have

$$\begin{aligned} M_{X_n}(t) &= e^{-\frac{\mu t}{\sigma/\sqrt{n}}} M_{\bar{Y}_n}\left(\frac{t}{\sigma/\sqrt{n}}\right) \\ &= e^{-\frac{\mu t}{\sigma/\sqrt{n}}} e^{\frac{\mu t}{\sigma/\sqrt{n}} + \frac{\sigma^2}{2} \left(\frac{t}{\sigma/\sqrt{n}}\right)^2} = e^{\frac{t^2}{2}} \end{aligned}$$


which is the mgf of $G(0, 1)$.

Example 16.1.5 (Example 5.8)

Let $Y_n \sim \text{Gam}(n, 2)$. Find the limiting distribution of

$$W_N = \frac{1}{2\sqrt{n}}(Y_n - 2n).$$

 **Solution**

By  Theorem 19, we have

$$\begin{aligned} M_{W_N}(t) &= e^{-t\sqrt{n}} M_{Y_n}(t) \left(\frac{t}{2\sqrt{n}} \right) = e^{-t\sqrt{n}} \left(1 - 2 \left(\frac{t}{2\sqrt{n}} \right) \right)^{-n} \\ &= \left[e^{\frac{t}{\sqrt{n}}} \left(1 - \frac{t}{\sqrt{n}} \right) \right]^{-n} = \left[\left(1 - \frac{t}{\sqrt{n}} \right) \sum_{k=0}^{\infty} \frac{(t/\sqrt{n})^k}{k!} \right]^{-n} \\ &= \left[\left(1 - \frac{t}{\sqrt{n}} \right) \left(1 + \frac{t}{\sqrt{n}} + \frac{t^2}{2n} + \frac{t^3}{3\sqrt{n}^3} + \dots \right) \right]^{-n} \\ &= \left(1 - \frac{t}{\sqrt{n}} + \frac{t}{\sqrt{n}} - \frac{t^2}{n} + \frac{t^2}{2n} - \frac{t^3}{2\sqrt{n}^3} + \frac{t^3}{3\sqrt{n}^3} - \frac{t^4}{3n^2} + \dots \right)^{-n} \\ &= \left(1 - \frac{t^2}{2n} - \underbrace{\frac{t^3}{6\sqrt{n}^3} - \frac{t^4}{3n^2} + \dots}_{\rightarrow 0 \text{ as } n \rightarrow \infty} \right)^{-n} \end{aligned}$$


So we have, by  Theorem 51,

$$\lim_{n \rightarrow \infty} M_{W_N}(t) = \lim_{n \rightarrow \infty} \left(1 - \frac{t^2/2}{n} - \frac{t^3}{6\sqrt{n}^3} - \frac{t^4}{3n^2} + \dots \right)^{-n} = e^{-(-\frac{t^2}{2})} = e^{\frac{t^2}{2}},$$

which is the mgf of $G(0, 1)$. So we know that

$$W_n \xrightarrow{D} W \sim G(0, 1)$$

for some limiting distribution W that has a $G(0, 1)$ distribution.

 **Theorem 57 (Central Limit Theorem)**

Suppose X_1, X_2, \dots is a sequence of IID rvs with $E(X_i) = \mu$ and

$\text{Var}(X_i) = \sigma^2 < \infty$, for $i = 1, 2, \dots$. Then

$$\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} = \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \xrightarrow{D} Z \sim N(0, 1)$$

where $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$.

⚡ We actually also need the condition that all moments of $\frac{X_i - \mu}{\sigma}$ exists and is bounded.

 **Proof**

Since the first and second moment exist, the mgf must be well-

defined, and is at least of class C^2 ³. We will use [Proposition 56](#) and

³You should remember this from your calculus courses. Otherwise, [Reference - Wiki].

[Theorem 50](#) in this proof. Firstly note that

$$E \left[\sum_{i=1}^n X_i \right] = n\mu \text{ and } \text{Var} \left(\sum_{i=1}^n X_i \right) = n\sigma^2$$

Let $\Sigma X_n = \sum_{i=1}^n X_i$ and $Z = \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma}$. Note that

$$Z = \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} = \frac{n\bar{X}_n - \mu n}{\sigma\sqrt{n}} = \frac{\Sigma X_n - n\mu}{\sigma\sqrt{n}} = \sum_{i=1}^n \frac{X_i - \mu}{\sigma\sqrt{n}}.$$

Let $Y_i = \frac{X_i - \mu}{\sigma}$ so that

$$Z = \sum_{i=1}^n \frac{1}{\sqrt{n}} Y_i.$$

Now the mgf of Z , by [Theorem 19](#) and [Proposition 39](#), is

$$M_Z(t) = \prod_{i=1}^n M_{Y_i} \left(\frac{t}{\sqrt{n}} \right) = M_{Y_i} \left(\frac{t}{\sqrt{n}} \right)^n. \quad (16.5)$$

Note that

$$E(Y_i) = E \left(\frac{X_i - \mu}{\sigma} \right) = \frac{1}{\sigma} E(X) - \frac{\mu}{\sigma} = 0$$

$$\text{Var}(Y_i) = \text{Var} \left(\frac{X_i - \mu}{\sigma} \right) = \frac{1}{\sigma^2} \text{Var}(X_i) = 1.$$

So by [Theorem 50](#) up to the second derivative, we have


$$M_{Y_i} \left(\frac{t}{\sqrt{n}} \right) = 1 + \frac{t^2}{2n} + \frac{M_{Y_i}^{(3)}(\zeta) \left(\frac{t}{\sqrt{n}} \right)^3}{3!}$$

where $\zeta \in \left[0, \frac{t}{\sqrt{n}} \right]$. Note that the last term in the above equation goes to 0 faster than the second term as $n \rightarrow \infty$, since we would have n^3 in the denominator, which is larger than n in the second term. With that, continuing with [Equation \(16.5\)](#) and now taking the limit as $n \rightarrow \infty$, we have

$$\lim_{n \rightarrow \infty} M_Z(t) = \lim_{n \rightarrow \infty} \left[1 + \frac{t^2}{2n} + \phi(n) \right]^n$$

where

$$\phi(n) = \frac{M_{Y_i}^{(3)}(\zeta) \left(\frac{t}{\sqrt{n}} \right)^3}{3!}.$$

Then by  Theorem 5.1, we have

$$\lim_{n \rightarrow \infty} M_Z(t) = e^{\frac{t^2}{2}}$$

which is the mgf of $N(0,1)$. This completes the proof. \square

Example 16.1.6 (Example 5.9)

Revisit Example 16.1.5 using the  Central Limit Theorem. Given

$$W_n = \frac{Y - 2n}{2\sqrt{n}}$$

where $Y \sim \text{Gam}(n,2)$, find sequences a_n and b_n such that

$$a_n(Y - b_n) \xrightarrow{D} Z \sim G(0,1).$$

Solution

Let X_1, X_2, \dots be a sequence of IID rvs with distribution $\text{Exp}(2)$. Note that we then have $E(X_i) = 2$ and $\text{Var}(X_i) = 4$. Let $Y_n = \sum_{i=1}^n X_i$. Note that

$$M_{Y_n}(t) = [M_{X_i}(t)]^n = (1 - 2t)^{-n}$$

which is the mgf of $\text{Gam}(n,2)$. By CLT, we have that

$$\frac{\frac{1}{n}Y_n - 2}{2/\sqrt{n}} \xrightarrow{D} G(0,1).$$

Note that

$$\frac{\frac{1}{n}Y_n - 2}{2/\sqrt{n}} = \frac{\sqrt{n}}{2n}(Y_n - 2n)$$

and so

$$a_n = \frac{1}{2}\sqrt{n} \text{ and } b_n = 2n$$

Example 16.1.7 (Example 5.10 (Course Notes 5.3.6))

Suppose $Y_n \sim \chi^2(n)$, $n = 1, 2, \dots$. Consider the sequence of rvs Z_1, Z_2, \dots where $Z_n = (Y_n - n)/\sqrt{2n}$. Show that

$$Z_n = \frac{Y_n - n}{\sqrt{2n}} \xrightarrow{D} Z \sim G(0,1).$$

Solution

Let $X_1, X_2, \dots \sim \chi^2(1)$ be IID rvs, and so $E[X_i] = 1$ and $\text{Var}(X_i) = 2$

This involves the knowledge that the sum of n exponential distributions with mean μ results in a gamma distribution with parameter n and μ . Since this has never been proven in this set of notes, it shall be proven here in this solution.

Once again our class pulls the card of "not explaining or showing stuff that they are supposed to". We will be using the fact that a sum of n Chi-Squared Distributions, each with degree of freedom k , will result in a gamma distribution with parameter $\frac{nk}{2}$ and 2. This will be proven in this exercise.

for $1 \leq i \leq n$, and let $Y_n = \sum_{i=1}^n X_i$. Note that

$$M_{Y_n}(t) = [M_{X_i}(t)]^n = (1 - 2t)^{-\frac{n}{2}}$$

is the mgf of $\chi^2(n)$. By CLT, we have that

$$\frac{\frac{1}{n} \sum_{i=1}^n X_i - 1}{\sqrt{2}/\sqrt{2}} \xrightarrow{D} Z \sim G(0, 1).$$

Note that

$$\frac{\frac{1}{n} \sum_{i=1}^n X_i - 1}{\sqrt{2}/\sqrt{n}} = \frac{Y_n - n}{\sqrt{2n}/\sqrt{n}} = \frac{Y_n - n}{\sqrt{2n}}.$$

This completes our example.

Example 16.1.8 (Example 5.11 (Course Notes 5.3.7))

Suppose $Y_n \sim \text{Bin}(n, p)$, $n = 1, 2, \dots$. Consider the sequence of rvs Z_1, Z_2, \dots , where $Z_n = \frac{Y_n - np}{\sqrt{np(1-p)}}$. Show that

$$Z_n = \frac{Y_n - np}{\sqrt{np(1-p)}} \xrightarrow{D} Z \sim N(0, 1).$$

Solution

Let X_1, X_2, \dots be a sequence of IID Bernoulli trials with success probability $0 \leq p \leq 1$. Note that for $i \in \mathbb{N}$, we have that $E[X_i] = p$ and $\text{Var}(X_i) = p(1-p)$. Now let $Y_n = \sum_{i=1}^n X_i$. We note that this satisfies our assumption, i.e. $Y_n \sim \text{Bin}(n, p)$ since

$$M_{Y_n}(t) = [M_{X_1}]^n = (pe^t + 1 - p)^n$$

is the mgf of $\text{Bin}(n, p)$. By CLT, we have that

$$\frac{\frac{1}{n} \sum_{i=1}^n X_i - p}{\sqrt{p(1-p)}/\sqrt{n}} \xrightarrow{D} Z \sim N(0, 1).$$

Note that

$$\frac{\frac{1}{n} \sum_{i=1}^n X_i - p}{\sqrt{p(1-p)}/\sqrt{n}} = \frac{Y_n - np}{\sqrt{np(1-p)}}$$

and so this completes our proof.

Proposition 58 (Other Limit Theorems)

1. $X_n \xrightarrow{P} a \wedge g$ continuous at $x = a \implies g(X_n) \xrightarrow{P} g(a)$.
2. $X_n \xrightarrow{P} a \wedge Y_n \xrightarrow{P} b \wedge g$ continuous at $(x, y) = (a, b) \implies$

$$g(X_n, Y_n) \xrightarrow{P} g(a, b).$$

3. (**Slutsky's Theorem**) $X_n \xrightarrow{D} X \wedge Y_n \xrightarrow{P} b \wedge g$ continuous at $(x, y) = (x, b)$ for all $x \in \text{supp}(X) \implies g(X_n, Y_n) \xrightarrow{D} g(X, b)$.
4. (**Continuous Mapping Theorem**) $X_n \xrightarrow{D} X \wedge g$ continuous at all $x \in \text{supp}(X) \implies g(X_n) \xrightarrow{D} g(X)$.

Proof

1. Since g is continuous at a , we have that

$$\forall \epsilon > 0 \exists \delta > 0 \forall x \in \text{supp}(X) (|x - a| < \delta \implies |g(x) - g(a)| < \epsilon)$$

It follows that

$$P[|g(X_n) - g(a)| < \epsilon] \geq P[|X_n - a| < \delta]$$

since $P(B) \geq P(A)$ whenever $A \subset B$ ⁴. Since $X_n \xrightarrow{P} a$, we have that for any $\delta > 0$, we have

$$1 \geq \lim_{n \rightarrow \infty} P[|g(X_n) - g(a)| < \epsilon] \geq \lim_{n \rightarrow \infty} P[|X_n - a| < \delta] = 1.$$

Thus

$$\lim_{n \rightarrow \infty} P[|g(X_n) - g(a)| < \epsilon] = 1$$

and so

$$g(X_n) \xrightarrow{P} g(a)$$

as required.

2. This is simply a more general case than (1). □
3. See this [Wikipedia article](#) for a proof. Requires knowledge of measure theory (in particular, convergence of measures).
4. See this [Wikipedia article](#) for a proof. Requires knowledge of measure theory (in particular, convergence of measures).

Example 16.1.9

If $X_n \xrightarrow{P} 10$ and $Y_n \xrightarrow{P} 2$, then $X_n Y_n \xrightarrow{P} 20$ since $g(x, y) = xy$ is continuous at $(10, 2)$.

If $Z_n \xrightarrow{D} Z \sim G(0, 1)$ and $a_n \xrightarrow{P} a$ where a is a constant, then


⁴ While I do not have a rigorous proof of this for our case here, it is a sensible result seeing that δ depends on ϵ . My hunch was right: thinking about the two probability measures using the definition of a random variable, we see that the $w \in S$ that works for $|X_n - a| < \delta$ will definitely work for $|g(X_n) - g(a)| < \epsilon$, but the converse is not necessarily true. Therefore, the events covered in the left term is a larger set that contains the event on the right.


$a_n Z_n \xrightarrow{D} aZ \sim (0, a)$ since $g(a, z) = az$ is continuous at (a, z) for all $z \in \text{supp}(Z)$.

Example 16.1.10 (Example 5.12 (Course Notes 5.3.10))

If $X_n \xrightarrow{P} a > 0$, $Y_n \xrightarrow{P} b \neq 0$ and $Z_n \xrightarrow{D} Z \sim G(0, 1)$, find the limiting distributions of each of the following:

1. $\sqrt{X_n}$
2. $X_n + Y_n$
3. $X_n Z_n$
4. $\frac{1}{Z_n}$

 **Solution**

1. Since $a > 0$, we have the function $g(x) = \sqrt{x}$ is continuous on a , and so by  Proposition 58, we have that $\sqrt{X_n} \xrightarrow{P} \sqrt{a}$.
2. Since the function $g(x, y) = x + y$ is continuous on the real number 2-tuple (a, b) , we have that $X_n + Y_n \xrightarrow{P} a + b$.
3. Since the function $g(x, z) = xz$ is continuous on the real number 2-tuple (x, z) for all $x \in \text{supp}(X_n)$ and $z \in \text{supp}(Z_n)$, by **Slutsky's Theorem**, we have $X_n Z_n \xrightarrow{D} aZ \sim G(0, a)$.
4. Note that the function $g(z) = \frac{1}{z}$ is continuous for all $z \in \text{supp}(X)$ except at $z = 0$. But since $Z \sim G(0, 1)$ is a continuous distribution, at a single point $z = 0$, $P(Z = 0) = 0$, and in a distribution it is of negligible value⁵. Therefore, we have that

$$\frac{1}{Z_n} \xrightarrow{D} \frac{1}{Z}$$

⁵ This is a painful thing to write down for me...

Example 16.1.11 (Example 5.13 (Course Notes 5.3.11))

Suppose $X_1, X_2, \dots \sim \text{Poi}(\mu)$ are IID rvs. Define $Z_n = \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sqrt{\bar{X}_n}}$. Find the limiting distribution of Z_n .

 **Solution**

Firstly, note that the parameter for a Poisson distribution is positive. Note that by CLT, we have

$$\frac{\bar{X}_n - \mu}{\sqrt{\mu}/\sqrt{n}} \sim Z \sim G(0, 1)$$

Note that since \sqrt{m} is a constant, we have that $\sqrt{m} \xrightarrow{P} \sqrt{m}$ is trivially

true. Thus by our limit theorems, we have

$$\sqrt{n}(\bar{X}_n - \mu) \sim \sqrt{m}Z \sim G(0, \sqrt{m}).$$

Now by the **Weak Law of Large Numbers**, we have

$$\bar{X}_n \xrightarrow{P} \mu.$$

Since a function $g(x) = \sqrt{x}$ is continuous for $x = \mu > 0$, we have that

$$\sqrt{\bar{X}_n} \xrightarrow{P} \sqrt{\mu}.$$

Then by the **Continuous Mapping Theorem**, we have that

$$Z_n = \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sqrt{\bar{X}_n}} \sim \frac{\sqrt{\mu}Z}{\sqrt{\mu}} \sim G(0, 1).$$

Theorem 59 (Generalized δ -Method)

Let X_1, X_2, \dots be a sequence of rvs such that

$$n^b(X_n - a) \xrightarrow{D} X$$

for some $b > 0$. Suppose the function $g(x)$ is differentiable at a and $g'(a) \neq 0$. Then

$$n^b[g(X_n) - g(a)] \xrightarrow{D} g'(a)X.$$

Proof

Using the **Mean Value Theorem**, we have that

$$g(X_n) = g(a) + g'(c)(X_n - a) \tag{16.6}$$

for some c in between X_n and a . Since $X_n \xrightarrow{P} a$ ⁶, and, WLOG, $X_n < c < a$, $c \xrightarrow{P} a$. Then by the **Continuous Mapping Theorem**, we have $g(c) \xrightarrow{P} g(a)$.

⁶ I have no idea why...

Now by rearranging Equation (16.6) and multiplying both sides by n^b , we get

$$n^b[g(X_n) - g(a)] = g'(c)n^b[X_n - a] \xrightarrow{D} g'(a)X$$

by **Slutsky's Theorem**. □

✦ **Corollary 60 (δ -Method)**


Let X_1, X_2, \dots be a sequence of IID rvs with mean μ and variance σ^2 . Suppose the function $g(x)$ is differentiable at μ and $g'(\mu) \neq 0$. Then

$$\sqrt{n}[g(\bar{X}_n) - g(\mu)] \xrightarrow{D} Z \sim N\left(0, [g'(\mu)]^2 \sigma^2\right)$$

✎ **Proof**

Note that by the same working as in Example 16.1.11 for $\sqrt{n}(\bar{X}_n - \mu)$, we have that⁷

$$\sqrt{n}(\bar{X}_n - \mu) \xrightarrow{D} Y \sim N(0, \sigma^2).$$

Then by  Theorem 59, we have that

$$\sqrt{n}(g(\bar{X}_n) - g(\mu)) \xrightarrow{D} g'(\mu)Y \sim N\left(0, [g'(\mu)]^2 \sigma^2\right)$$

as required. □

⁷ This is actually stated as an assumption on the δ -Method Wikipedia article - [Reference].

Example 16.1.12 (Example 5.14)


Let $X_1, X_2, \dots \sim \text{Geo}(p)$ is a sequence of IID rvs, each with support $\text{supp}(X_i) = \{0, 1, 2, \dots\}$. Derive the limiting distribution of

$$W_n = \sqrt{n} \left(\frac{1}{\bar{X}_n} - \frac{p}{1-p} \right)$$

✎ **Solution**

Note that by CLT, we have that

$$\frac{\bar{X}_n - \frac{1-p}{p}}{\sqrt{\frac{1-p}{p^2}} / \sqrt{n}} \xrightarrow{D} Z \sim N(0, 1)$$

Then using  Proposition 58, we have

$$\sqrt{n} \left(\bar{X}_n - \frac{1-p}{p} \right) \xrightarrow{D} \sqrt{\frac{1-p}{p^2}} Z \sim N\left(0, \frac{1-p}{p^2}\right)$$

We took $g(x) = \frac{1}{x}$ and the given solution in class somehow circumvents the fact that $x = 0$ is a case.

Example 16.1.13 (Example 5.15)

Suppose that $X_1, X_2, \dots \sim \text{Exp}(\theta)$ is a sequence of IID rvs. Find constants a_n and b_n such that

$$W_n = b_n(\bar{X}_n^2 - a_n)$$

has a non-degenerate limiting distribution.

 Solution

By CLT, we have

$$\frac{\bar{X}_n - \theta}{\theta/\sqrt{n}} \xrightarrow{D} Z \sim N(0, 1)$$

which then by  Proposition 58, we have

$$\sqrt{n}(\bar{X}_n - \theta) \xrightarrow{D} \theta Z \sim N(0, \theta^2).$$

Let $g(x) = x^2$. Then g is continuous on any $x \in \text{supp}(X_n)$ and on θ .

Then using  Corollary 60, we have

$$\sqrt{n}(\bar{X}_n^2 - \theta^2) \sim \theta^2 Z \sim N(0, 4\theta^2).$$

So we just need to pick $b_n = \sqrt{n}$ and $a_n = \theta^2$.

17 Lecture 17 Jun 28th 2018

17.1 Estimation

SUPPOSE $X_1, \dots, X_n \sim f(x; \theta)$ is an IID sequence of rvs, where $f(x; \theta)$ is the pf of the X_i 's. The joint distribution of X_1, \dots, X_n is

$$\prod_{i=1}^n f(x_i; \theta)$$

where the unknown parameter θ can either be a scalar in Ω , where Ω is the parameter space or the set of possible values of θ , or a vector,

$$\text{i.e. } \theta = \begin{pmatrix} \theta_1 & \theta_2 & \dots & \theta_n \end{pmatrix}^T.$$

We are interested in making inferences about the unknown parameter θ , i.e. we want to find **estimators** (point and interval) of θ and we want to test our hypothesis about θ .

Before proceeding the the rest of this chapter (actual chapter, ahem...), we require the following definitions.

Definition 57 (Statistic)

A **statistic**, $T = T(X) = T(X_1, \dots, X_n)$, is a function of the data which does not depend on any unknown parameter(s).

Example 17.1.1

Suppose X_1, \dots, X_n is a random sample from a distribution with $E[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$, where μ and σ^2 are unknown. The sample mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ and the sample variance $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$ are statistics, while $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$ is not a statistic.

Definition 58 (Estimators and Estimates)

A statistic $T = T(X) = T(X_1, \dots, X_n)$ that is used to estimate $\tau(\theta)$, a function of θ , is called an **estimator** of $\tau(\theta)$, and an observed value of the statistic $t = t(x) = t(x_1, \dots, x_n)$ is called an **estimate** of $\tau(\theta)$.

Example 17.1.2

Suppose X_1, \dots, X_n are IID rvs with $E[X_i] = \mu$. The rv \bar{X} is an estimator of μ . For a given set of observations, x_1, \dots, x_n , the number $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$ is an estimate of μ .

There are certain “properties” of an estimator that we look for, in particular, for an estimator, $\tilde{\theta}$, of an unknown parameter θ , we want that the estimator

- is not biased, aka an **unbiased estimator**, i.e. $E[\tilde{\theta}] = \theta$;
- has small variance;
- is consistent, i.e. $\tilde{\theta} \xrightarrow{P} \theta$.

17.1.1 Maximum Likelihood Estimation

SUPPOSE X is a discrete rv with pf $P(X = x; \theta) = f(x; \theta)$, $\theta \in \Omega$ where the scalar parameter θ is unknown. Suppose x is an observed value of the rv X . Then the probability of observing this value is

$$P(X = x; \theta) = f(x; \theta).$$

With the observed value of x substituted into $f(x; \theta)$, we have a function of the parameter θ , referred to as the **likelihood function**, and denoted $L(\theta)$. In the absence of any other information, it seems logical (temptingly so) that we should estimate the parameter θ using a value that is “the most compatible” with the data. E.g., we might choose the value of θ of which it maximizes the probability of the observed data, or equivalently, the value of θ which maximizes the likelihood function $L(\theta)$.

But first, let us formally state the definition of a likelihood func-

tion.

 **Definition 59 (Likelihood function)**

Suppose X is an rv with pf $f(x; \theta)$, where $\theta \in \Omega$ is a scalar. If x is the observed data, then the **likelihood function** for θ based on x is

$$L(\theta) = P(X = x; \theta) = f(x; \theta) \quad \text{for } \theta \in \Omega.$$

Similarly so, suppose X_1, \dots, X_n is a random sample from a distribution, each with pf $f(x; \theta)$, and let x_1, \dots, x_n be the observed data. Then the **likelihood function** for θ based on x_1, \dots, x_n is

$$L(\theta) = P(X_1 = x_1, \dots, X_n = x_n; \theta) = \prod_{i=1}^n f(x_i; \theta) \quad \text{for } \theta \in \Omega.$$

“ Note

As discussed earlier, a natural estimate of θ is the value which maximizes the probability of the observed sample, and we denote this notion as

$$\hat{\theta} = \arg \max_{\theta \in \Omega} L(\theta)$$

and call $\hat{\theta}$ the **maximum likelihood estimate** (ML estimate). In practice, it is often convenient to work, instead, with the natural logarithm of the likelihood function, in which we call the **Log-likelihood**:

$$\ell(\theta) = \ln L(\theta).$$

Note that the maximum likelihood estimate for both the likelihood function and the log-likelihood are the same since \log is a strictly increasing function.

Example 17.1.3 (Example 6.1)

Consider flipping a coin repeatedly, where for $i \in \mathbb{N}$

$$X_i = \begin{cases} 1 & \text{if the } i^{\text{th}} \text{ flip lands on heads} \\ 0 & \text{otherwise} \end{cases}$$

Based on 4 independent flips, we are given that $X_1, X_2, X_3, X_4 \sim \text{Bernoulli}(p)$ are IID rvs. The sample has been observed as $x_1 = 1, x_2 = 1, x_3 = 0, x_4 = 1$. Write the probability of this sample as a function of p . Then, compute the likelihood function and find the ML estimate.

Solution

Since the X_i 's are IID rvs, we have

$$\begin{aligned} f(x_1, x_2, x_3, x_4; p) &= \prod_{i=1}^4 P(X_i = x_i; p) \\ &= \prod_{i=1}^4 p^{x_i} (1-p)^{1-x_i} = p^{\sum_{i=1}^4 x_i} (1-p)^{\sum_{i=1}^4 (1-x_i)}. \end{aligned}$$

Note that the likelihood function for p under the observed values is therefore

$$L(p) = p^3(1-p).$$

To find the ML estimate, we do

$$0 = \frac{dL(p)}{dp} = 3p^2 - 4p^3 = p^2(3 - 4p) \quad (17.1)$$

and we observe that $p = 0$ or $\frac{3}{4}$. Clearly, $p = \frac{3}{4}$ maximizes the likelihood function.

“ Note

WE HAVE BEEN talking almost solely about the discrete case, so what about the continuous case? We have that $P(X = x) = 0$, and so we consider a small neighbourhood of radius $\delta > 0$. Then for a small $\delta > 0$ around any point $x \in \text{supp}(X)$, we have that

$$P(x - \delta < X < x + \delta; \theta) = \int_{x-\delta}^{x+\delta} f(t; \theta) dt \approx 2\delta \cdot f(x; \theta). \quad (17.2)$$

And so for an observed value x , since δ is fixed in Equation (17.2), we

have that the value of θ that maximizes $f(t; \theta)$ also maximizes $2\delta \cdot f(t; \theta)$.

⚠ Warning

We shall clarify the following notations: we use

- $\tilde{\theta}$ as the estimator, which is an rv; and
- $\hat{\theta}$ as an estimate, which is a fixed value.

Example 17.1.4 (Example 6.2 (Course Notes 6.2.4))

Recall the coin flip example in Example 17.1.3. Suppose $X_1, \dots, X_n \sim \text{Bernoulli}(p)$ is a sequence of IID rvs. Calculate the ML estimate of p .

✎ Solution

Since X_1, \dots, X_n are IID, let $\vec{x} = (x_1 \ x_2 \ \dots \ x_n)$, then the joint distribution is

$$\begin{aligned} f(\vec{x}; p) &= \prod_{i=1}^n f(x_i; p) \\ &= \prod_{i=1}^n p^{x_i} (1-p)^{1-x_i} = p^{\sum_{i=1}^n x_i} (1-p)^{\sum_{i=1}^n (1-x_i)} \end{aligned}$$

To get an ML estimator, we shall use the log-likelihood:

$$\ell(p) = \left(\sum_{i=1}^n x_i \right) \ln p + \left(n - \sum_{i=1}^n x_i \right) \ln(1-p)$$

and so to get the ML estimator

$$\begin{aligned} 0 &= \frac{\partial}{\partial p} \ell(p) \Big|_{p=\hat{p}} = \left(\sum_{i=1}^n x_i \right) \left(\frac{1}{\hat{p}} \right) - \left(n - \sum_{i=1}^n x_i \right) \left(\frac{1}{1-\hat{p}} \right) \\ &= \frac{\sum x_i}{\hat{p}} - \frac{n - \sum x_i}{1-\hat{p}} = \frac{(1-\hat{p})\sum x_i - n\hat{p} + \hat{p}\sum x_i}{\hat{p}(1-\hat{p})} \\ &= \frac{\sum x_i - n\hat{p}}{\hat{p}(1-\hat{p})} \end{aligned}$$

where we represent $\sum_{i=1}^n x_i$ by $\sum x_i$ for sanity. Thus we have that

$$n\hat{p} = \sum_{i=1}^n x_i \implies \hat{p} = \frac{1}{n} \sum_{i=1}^n x_i = \bar{x}$$

Example 17.1.5 (Example 6.3 (Course Notes 6.2.5))

Note that when looking for an **ML estimate**, we should also check for the case when $\frac{\partial^2}{\partial p^2} \ell(p) \Big|_{p=\hat{p}} < 0$ to ensure maximality (instead of minimality).

Note that the **ML estimator** in this case would be represented as

$$\tilde{p} = \bar{X}.$$

Suppose we have the data $\vec{x} = (x_1 \ x_2 \ \dots \ x_n)$ for the sequence of IID rvs $X_1, X_2, \dots, X_n \sim \text{Poi}(\theta)$. Find the likelihood function, log-likelihood, the ML estimate $\hat{\theta}$, and the ML estimator $\tilde{\theta}$.

 **Solution**

Since each of the X_i 's are IID, we have that their joint pmf, and in particular, their likelihood function, is

$$L(\theta) = \prod_{i=1}^n \frac{e^{-\theta} \theta^{x_i}}{x_i!} = e^{-n\theta} \theta^{\sum_{i=1}^n x_i} \left(\prod_{i=1}^n x_i! \right)^{-1}.$$

Then the log-likelihood is

$$\ell(\theta) = -n\theta + \left(\sum_{i=1}^n x_i \right) \log \theta - \sum_{i=1}^n \log(x_i!).$$

To get the ML estimate, note that

$$0 = \frac{\partial}{\partial \theta} \ell(\theta) \Big|_{\theta=\hat{\theta}} = -n + \frac{1}{\hat{\theta}} \sum_{i=1}^n x_i$$

and so

$$\hat{\theta} = \frac{1}{n} \sum_{i=1}^n x_i = \bar{x}.$$

Consequently, we have that the ML estimator is

$$\tilde{\theta} = \bar{X}.$$

18 Lecture 18 Jul 3rd 2018

18.1 Estimation (Continued)

18.1.1 Maximum Likelihood Estimation (Continued)

Definition 60 (Score Function)

The **score function** is defined as

$$S(\theta) = S(\theta; x) = \frac{d}{d\theta} \ell(\theta) = \frac{d}{d\theta} \ln L(\theta) \quad \theta \in \Omega.$$

“ Note

Notice that to find the ML estimate, we usually set the score function to zero and solve for $S(\theta) = 0$.

Definition 61 (Information Function)

The **information function** is defined as

$$I(\theta) = I(\theta; x) = -\frac{d^2}{d\theta^2} \ell(\theta) = -\frac{d^2}{d\theta^2} \ln L(\theta) \quad \theta \in \Omega$$

If $\hat{\theta}$ is the ML estimate of θ , then $I(\hat{\theta})$ is called the **observed information**.

Example 18.1.1

Going back to our example in Example 17.1.4, we had that the likeli-

hood function was

$$L(p) = p^{\sum_{i=1}^n x_i} (1-p)^{n-\sum_{i=1}^n x_i}$$

the log-likelihood was

$$\ell(p) = \left(\sum_{i=1}^n x_i \right) \ln p + \left(n - \sum_{i=1}^n x_i \right) \ln(1-p).$$

So the score function is

$$S(p) = \frac{d}{dp} \ell(p) = \frac{1}{p} \sum_{i=1}^n x_i - \frac{1}{1-p} \left(n - \sum_{i=1}^n x_i \right)$$

Consequently, the information function is

$$\begin{aligned} I(p) &= -\frac{d^2}{dp^2} \ell(p) = -\frac{d}{dp} S(p) \\ &= \frac{1}{p^2} \sum_{i=1}^n x_i + \frac{1}{(1-p)^2} \left(n - \sum_{i=1}^n x_i \right) \end{aligned}$$

“ Note

Recall from the calculus knowledge of the 2nd derivative being the “curvature” of the original function, and so $I(\theta)$ is a function about the curvature of the log-likelihood. In particular, we have that $I(\hat{\theta})$ tells us about the concavity of the log-likelihood function at the ML estimate.

Note that the information function is a function that has two variables: the unknown parameter θ , and the data $\vec{X} = (X_1 \dots X_n)$.

In a later lecture, we shall see how the observed information $I(\hat{\theta})$ can be used to construct approximate confidence intervals for the unknown parameter θ .

Definition 62 (Fisher Information)

If θ is a scalar, then the **expected information**, or **Fisher information** (function) is given by

$$J(\theta) = E[I(\theta; \vec{X})] = E \left[-\frac{\partial^2}{\partial \theta^2} \ell(\theta; \vec{X}) \right] \quad \theta \in \Omega$$

Note

Just to take away the layers of definitions and compare the Fisher information with our pf for the rv(s), if X_1, \dots, X_n is a random sample (i.e. IID rvs), each with pf $f(x; \theta)$, then

$$J(\theta) = E \left[-\frac{\partial^2}{\partial \theta^2} \ell(\theta; \vec{X}) \right] = nE \left[-\frac{\partial^2}{\partial \theta^2} \ln f(\vec{X}; \theta) \right]$$

where $\vec{X} = (X_1 \dots X_n)$.

Example 18.1.2 (Example 6.4 (Course Notes 6.2.10))

Suppose $X_1, \dots, X_n \sim \text{Bernoulli}(p)$ is a sequence of IID rvs. We have showed in Example 17.1.4 that the ML estimator of p is $\hat{p} = \bar{X}$. Calculate the Fisher information and compare it with the variance of the ML estimator of p .

Solution

Recall that from Example 18.1.1, we had

$$\begin{aligned} \ell(p) &= \left(\sum_{i=1}^n x_i \right) \ln p + \left(n - \sum_{i=1}^n x_i \right) \ln(1-p) \\ S(p) &= \frac{1}{p} \sum_{i=1}^n x_i - \frac{1}{1-p} \left(n - \sum_{i=1}^n x_i \right) \\ I(p) &= \frac{1}{p^2} \sum_{i=1}^n x_i + \frac{1}{(1-p)^2} \left(n - \sum_{i=1}^n x_i \right), \end{aligned}$$

So the Fisher information is

$$\begin{aligned} E[I(p)] &= E \left[\frac{1}{p^2} \sum_{i=1}^n X_i + \frac{1}{(1-p)^2} \left(n - \sum_{i=1}^n X_i \right) \right] \\ &= \frac{1}{p^2} \sum_{i=1}^n E[X_i] + \frac{1}{(1-p)^2} \left(n - \sum_{i=1}^n E[X_i] \right) \\ &= \frac{1}{p^2} (np) + \frac{1}{(1-p)^2} (n - np) \\ &= \frac{n}{p} + \frac{n}{1-p} = \frac{n}{p(1-p)}. \end{aligned}$$

On the other hand, note that the variance of the ML estimator is

$$\text{Var}(\hat{p}) = \text{Var}(\bar{X}) = \frac{1}{n} \text{Var}(X_i) = \frac{p(1-p)}{n}.$$

Example 18.1.3 (Example 6.5 (Course Notes 6.2.10))

Suppose $X_1, \dots, X_n \sim \text{Poi}(\theta)$ is a sequence of IID rvs. We showed in Example 17.1.5 the ML estimator of θ is $\tilde{\theta} = \bar{X}$. Calculate the Fisher information and compare it with the variance of the ML estimator of θ .

 Solution

We had the that log-likelihood is

$$\ell(\theta) = -n\theta + \left(\sum_{i=1}^n x_i \right) \log \theta - \sum_{i=1}^n \log(x_i!).$$

So the score function is

$$S(\theta) = -n + \frac{1}{\theta} \sum_{i=1}^n x_i,$$

and so the information function is

$$I(\theta; x) = \frac{1}{\theta^2} \sum_{i=1}^n x_i.$$

Then the Fisher information is

$$E[I(\theta; \bar{X})] = \frac{n}{\theta^2} E[X_i] = \frac{n}{\theta}.$$

The variance of the ML estimator of θ is

$$\text{Var}(\tilde{\theta}) = \frac{1}{n} \text{Var}(X_i) = \frac{\theta}{n}.$$

“ Note

We observe from the above two examples that

$$J(\theta) = \frac{1}{\text{Var} \tilde{\theta}}$$

where $\tilde{\theta} = \bar{X}$. It is tempting to verify if this is the case in general, but there does not seem to be reliable sources that points to this case (at least, online searches have yet to yield me results). The lecture claims that this is only true for certain families of distributions.

Example 18.1.4 (Example 6.6 (Course Notes 6.2.12))

Suppose X_1, \dots, X_n is a random sample, each from a distribution with pdf

$$f(x; \theta) = \theta x^{\theta-1} \quad 0 \leq x \leq 1, \quad \theta > 0.$$

Find the score function, the ML estimator of θ , the information function and the observed information.

 **Solution**

Since X_1, \dots, X_n are IID, we have

$$L(\theta) = \prod_{i=1}^n \theta x_i^{\theta-1} = \theta^n \left(\prod_{i=1}^n x_i \right)^{\theta-1}.$$

So the log-likelihood is

$$\ell(\theta) = n \ln \theta + (\theta - 1) \ln \left(\prod_{i=1}^n x_i \right) = n \ln \theta + (\theta - 1) \sum_{i=1}^n \ln x_i.$$

The score function is therefore

$$S(\theta) = \frac{n}{\theta} + \sum_{i=1}^n \ln x_i,$$

and so the ML estimator of θ is

$$\hat{\theta} = -\frac{n}{\sum_{i=1}^n \ln x_i}.$$

From the score function, the information function is

$$I(\theta) = \frac{n}{\theta^2},$$

and so the observed information is

$$I(\hat{\theta}) = \frac{n}{\left(-\frac{n}{\sum_{i=1}^n \ln x_i} \right)^2} = \frac{1}{n} \left(\sum_{i=1}^n \ln x_i \right)^2$$

Note

In the above example, note that the Fisher information is

$$J(\theta) = E[I(\theta; \vec{X})] = \frac{n}{\theta^2}.$$

Example 18.1.5 (Example 6.7 (Course Notes 6.2.13))

Suppose $X_1, \dots, X_n \sim \text{Unif}(0, \theta)$ is a random sample. Find the ML estimator of θ .

 **Solution**

Note that $f_{X_i}(x_i; \theta) = \frac{1}{\theta}$ for $0 \leq x_i \leq \theta$. So the likelihood function is

$$\begin{aligned} L(\theta) &= \prod_{i=1}^n \frac{1}{\theta} \mathbb{1}_{\{0 \leq x_i \leq \theta\}} \\ &= \theta^{-n} \mathbb{1}_{\{0 \leq x_1 \leq \theta, 0 \leq x_2 \leq \theta, \dots, 0 \leq x_n \leq \theta\}} \\ &= \theta^{-n} \mathbb{1}_{\{0 \leq x_{(n)} \leq \theta\}} \end{aligned}$$

where we use the order statistics notation to simplify the equation. Notice that if we take the derivative of the likelihood function from this point and try to get the ML estimate/estimator, we would run into the following equation:

$$-\frac{n}{\theta^{n+1}} = 0$$

in which we would have trouble getting the MLE.

Example 18.1.6 (Example 6.8)

Suppose X_1, \dots, X_n is a random sample, each from the $\text{Unif}(\theta, \theta + 1)$ distribution. Find the ML estimator of θ .


 **Solution**

This time, we have that the pdf for each X_i is

$$f_{X_i}(x_i; \theta) = \mathbb{1}_{\{\theta \leq x_i \leq \theta + 1\}}.$$


Then the likelihood function is

$$\begin{aligned} L(\theta) &= \prod_{i=1}^n f(x_i; \theta) = \mathbb{1}_{\{\theta \leq x_1 \leq \theta + 1, \dots, \theta \leq x_n \leq \theta + 1\}} \\ &= \mathbb{1}_{\{\theta \leq x_{(1)}\}} \mathbb{1}_{\{x_{(n)} \leq \theta + 1\}} \\ &= \mathbb{1}_{\{\theta \leq x_{(1)}\}} \mathbb{1}_{\{x_{(n)} - 1 \leq \theta\}} = \mathbb{1}_{\{x_{(n)} - 1 \leq \theta \leq x_{(1)}\}} \end{aligned}$$



where we, once again, use the order statistics notation. Consequently, we have the graph of θ versus $L(\theta)$ on the right.

Thus we observe that the MLE is not unique, since it can be any number between $x_{(n)} - 1$ and $x_{(1)}$.

 **Theorem 61 (Invariance Property of the MLE)**


Suppose $\tau = h(\theta)$ is an injective function of θ . Suppose also that $\hat{\theta}$ is the ML estimator of θ . Then $\hat{\tau} = h(\hat{\theta})$ is the ML estimator of τ .

 **Warning**

We are short on certain tools to actually prove this theorem.


Example 18.1.7 (Example 6.9)

Suppose $X_1, \dots, X_n \sim f(x; \theta) = \theta x^{\theta-1} \mathbb{1}_{\{0 < x < 1\}}$, where $\theta > 0$. Find the MLE of the median of the distribution.

 **Solution**

Recall from Example 18.1.4 that we had

$$\hat{\theta} = -\frac{n}{\sum_{i=1}^n \ln x_i}.$$

Let m be the median. The goal is to use  Theorem 61. **Something feels off...**

19 Lecture 19 July 5th 2018

19.1 Estimation (Continued 2)

19.1.1 Maximum Likelihood Estimation (Continued 2)

Definition 63 (Relative Likelihood)

Suppose $X_1, \dots, X_n \sim f(x; \theta)$ is an IID sequence of rvs where the likelihood function is $L(\theta)$ and the MLE of θ is $\hat{\theta}$. The **relative likelihood** function is defined by

$$R(\theta) = R(\theta; x) = \frac{L(\theta)}{L(\hat{\theta})}, \quad \theta \in \Omega$$

“ Note

Note that since $L(\hat{\theta})$ is maximum since $\hat{\theta}$ is the maximum likelihood estimate, we have that $L(\theta) \leq L(\hat{\theta})$, and so¹

$$0 \leq R(\theta) = \frac{L(\theta)}{L(\hat{\theta})} \leq 1.$$

¹ Note that the likelihood function $L(\theta) = f(x; \theta)$ is nonnegative, since the cdf of a distribution is nondecreasing, and so the pf must be nonnegative.

“ Note

Given the data $\theta = (x_1 \dots x_n)$, we say that

- if $R(\theta)$ is small (usual threshold is 0.1), then we say that θ is **implausible**;
- if $R(\theta)$ is large (usual threshold is 0.5), then we say that θ is **plausible**.

Definition 64 (Likelihood Region & Likelihood Interval)

The set of θ values for which $R(\theta) \geq p$ is called a $100p\%$ **likelihood region** for θ . If the region is an interval of real values, then it is called a $100p\%$ **likelihood interval (LI)** for θ .

Note

Using the definition, we can extend on the previous note:

- Values inside the 10% LI are referred to as **plausible** and values outside of this interval as **implausible**;
- Values inside a 50% LI are very plausible; and
- Values outside a 1% LI are very implausible in light of the data.

As how we had the log-likelihood from the likelihood function, we can have the log relative likelihood.

Definition 65 (Log Relative Likelihood)


The **log relative likelihood** is the natural logarithm of the relative likelihood function:

$$r(\theta) = r(\theta; x) = \ln[R(\theta)] = \log[L(\theta)] - \log[L(\hat{\theta})] = \ell(\theta) - \ell(\hat{\theta})$$

for $\theta \in \Omega$.

Note (Unimodality)

If for $\theta \in \Omega$, $R(\theta)$ is unimodal as shown in the graph to the right, then $R(\theta) \geq p$ will give us the likelihood interval, in which this case we have that the interval is $[a_p, b_p]$.

**Example 19.1.1 (Example 6.10)**

Let $X_1, \dots, X_{100} \sim \text{Poi}(\theta)$ be an IID sequence of rvs. Based on the observed values, you are given that $\sum_{i=1}^{100} x_i = 980$. Find a 10% and 50% likelihood intervals for θ .

 **Solution**

From Example 17.1.5, we had

$$L(\theta) = e^{-n\theta} \theta^{n\bar{x}} \left(\prod_{i=1}^n x_i! \right)^{-1} \quad \text{and } \hat{\theta} = \bar{x}.$$

The relative likelihood is therefore

$$R(\theta) = \frac{L(\theta)}{L(\hat{\theta})} = \frac{e^{-n\theta} \theta^{n\bar{x}} \left(\prod_{i=1}^n x_i! \right)^{-1}}{e^{-n\hat{\theta}} \hat{\theta}^{n\bar{x}} \left(\prod_{i=1}^n x_i! \right)^{-1}} = e^{-n(\theta - \bar{x})} \left(\frac{\theta}{\bar{x}} \right)^{n\bar{x}}.$$

We were given that

$$\bar{x} = \frac{1}{100} \sum_{i=1}^{100} x_i = \frac{980}{100} = 9.8,$$

and so get

$$R(\theta) = e^{-100(\theta - 9.8)} \left(\frac{\theta}{9.8} \right)^{980}.$$

Using WolframAlpha, we get that the 50% likelihood interval for θ is [9.436, 10.173]. Unfortunately so, I cannot get the solution for the 10% likelihood interval since the allowed free computation time is exceeded, and so I shall leave the likelihood interval as

$$\left\{ \theta \in \Omega : e^{-100(\theta - 9.8)} \left(\frac{\theta}{9.8} \right)^{980} > 0.1 \right\}.$$

Example 19.1.2 (Example 6.11 (Course Notes 6.2.24))

Suppose X_1, \dots, X_n is a random sample from $\text{Exp}(1, \theta)$. Plot the relative likelihood function for θ if $n = 20$ and $x_{(1)} = \min\{x_1, \dots, x_n\} = 1$. Find the 10% and 50% likelihood intervals for θ .

 **Solution**

Recall the definition of the Double Parameter Exponential Distribution. We have that the pdf of each X_i is

$$f_{X_i}(x_i) = e^{-\frac{x_i - \theta}{1}} = e^{-(x_i - \theta)}.$$

Since the X_i 's form a random sample, we have that

$$L(\theta) = \prod_{i=1}^n e^{-(x_i-\theta)} \mathbb{1}_{\{x_i \geq \theta\}} = e^{-\sum_{i=1}^n x_i + n\theta} \mathbb{1}_{\{x_{(1)} \geq \theta\}}.$$

Firstly, note that $L(\theta)$ is an increasing function with respect to θ . Now, note that if the indicator function is 1, we would have that $\hat{\theta}_0 = \bar{x}$, where we denote the "supposed" ML estimate as $\hat{\theta}_0$. However, since the condition for the indicator function to be 1 is that $\theta \leq x_{(1)}$, we must then have that $\hat{\theta} = x_{(1)}$, since $x_{(1)} \leq \bar{x}$ as $x_{(1)} = \min\{x_1, \dots, x_n\}$.

With that, we can calculate the relative likelihood function:

$$R(\theta) = \frac{L(\theta)}{L(\hat{\theta})} = e^{n(\theta - x_{(1)})} \mathbb{1}_{\{x_{(1)} \geq \theta\}}.$$

Given $n = 20$ and $x_{(1)} = 1$, we have that

$$R(\theta) = e^{20(\theta-1)} \mathbb{1}_{\{\theta \leq 1\}}.$$

To get the 50% likelihood interval, we have

$$\begin{aligned} e^{20(\theta-1)} &\geq 0.5 \\ \theta &\geq 1 + \frac{1}{20} \ln 0.5. \end{aligned}$$

Note that $\ln 0.5 < 0$, so we have that the 50% likelihood interval is $[1 + 0.05 \ln 0.5, 1]$. For the 10% likelihood interval, we have

$$\begin{aligned} e^{20(\theta-1)} &\geq 0.1 \\ \theta &\geq 1 + \frac{1}{20} \ln 0.1. \end{aligned}$$

Again, note that $\ln 0.1 < 0$, and so we have that the 10% likelihood interval is $[1 + 0.05 \ln 0.1, 1]$.

WE CAN EXTEND our methods analogously for the multiparameter case. Assume $\theta = (\theta_1 \dots \theta_k)^T$. Then the likelihood function, $L(\theta)$, is a function of k parameters. Then the log-likelihood function is $\ell(\theta) = \ln L(\theta)$. From there, we can then use the same approach to get the ML estimate of θ ,

$$\hat{\theta} = (\hat{\theta}_1 \dots \hat{\theta}_k)^T = \arg \max_{\theta_1, \dots, \theta_k} \ell(\theta).$$

We shall also note that we can extend [Theorem 61](#) to the multivariate case.

Furthermore, we will restate some of the definitions that were stated in univariate context for the multivariate case.

Definition 66 (Score Vector)

Suppose $X_1, \dots, X_n \sim f(x; \theta)$ is a sequence of IID rvs, where $\theta = (\theta_1 \dots \theta_k)^T$. The **score vector** is defined as

$$S(\theta) = S(\theta; x) = \left[\frac{\partial \ell}{\partial \theta_1} \quad \dots \quad \frac{\partial \ell}{\partial \theta_k} \right]^T$$

Definition 67 (Information Matrix)

Suppose $X_1, \dots, X_n \sim f(x; \theta)$ be a sequence of IID rvs, where $\theta = (\theta_1 \dots \theta_k)^T$. The **information matrix**, $I(\theta) = I(\theta; x)$, is a $k \times k$ symmetric matrix whose (i, j) entry is given by

$$-\frac{\partial^2}{\partial \theta_i \partial \theta_j} \ell(\theta).$$

The constant matrix $I(\hat{\theta})$ is called the **observed information matrix**.

Definition 68 (Fisher Information Matrix)

Suppose $X_1, \dots, X_n \sim f(x; \theta)$ is a sequence of IID rvs, where $\theta = (\theta_1 \dots \theta_k)^T$. The **Fisher information matrix**, $J(\theta)$ is a $k \times k$ symmetric matrix whose (i, j) entry is given by

$$E \left[-\frac{\partial^2}{\partial \theta_i \partial \theta_j} \ell(\theta; X) \right],$$

where $X = (X_1 \dots X_k)^T$.

Definition 69 (Likelihood Region - Multivariate)

Suppose $X_1, \dots, X_n \sim f(x; \theta)$ is a sequence of IID rvs, where $\theta =$

$(\theta_1 \dots \theta_k)^T$. The set of $\theta = (\theta_1 \dots \theta_k)^T$ for which $R(\theta) \geq p$, for some $0 \leq p \leq 1$, is called a 100p% **likelihood region** for θ , which is a region in a k -dimensional space \mathbb{R}^k .

“ Note

Recall that in the univariate case, we can certainly check if our choice of the $\hat{\theta}$ is indeed a maximizing value by taking the second derivative of $\ell(\theta)$ or $L(\theta)$, and checking that $I(\hat{\theta}) > 0$. The analogous approach for the multivariate case is to ensure that the **observed information matrix**, $I(\hat{\theta})$ is **positive definite**².

We have two methods to verify that a matrix M is positive definite:

- $\det M > 0$;
- eigenvalues of M are all positive.

² A matrix M is called a **positive definite** if the scalar $z^T M z > 0$ for every non-zero column vector z .

Example 19.1.3 (Example 6.12 Part 1)

Suppose $X_1, \dots, X_n \sim N(\mu, \sigma^2)$ is a sequence of IID rvs.

1. Derive the MLE of the parameters μ and σ^2 .
2. Derive the MLE of the coefficient of variation $CV = \frac{\sigma}{\mu}$.

Solution

1. Since the X_i 's are IID, we have

$$L(\mu, \sigma^2) = (2\pi)^{-\frac{n}{2}} (\sigma^2)^{-\frac{n}{2}} \exp \left\{ -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \right\}.$$

Then the log-likelihood is

$$\ell(\mu, \sigma^2) = -\frac{n}{2} \ln(2\pi) - \frac{n}{2} \log \sigma^2 - \frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2.$$

Now, note that the first partial derivatives of $\ell(\mu, \sigma^2)$ are

$$\begin{aligned} \frac{\partial \ell}{\partial \mu} &= \frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu) = \frac{1}{\sigma^2} \left(\sum_{i=1}^n x_i - n\mu \right) \\ \frac{\partial \ell}{\partial \sigma^2} &= -\frac{n}{2\sigma^2} + \frac{1}{2\sigma^4} \sum_{i=1}^n (x_i - \mu)^2. \end{aligned}$$

So the score vector is

$$S(\mu, \sigma^2) = \begin{bmatrix} \frac{\partial \ell}{\partial \mu} \\ \frac{\partial \ell}{\partial \sigma^2} \end{bmatrix} = \begin{bmatrix} \frac{1}{\sigma^2} (\sum_{i=1}^n x_i - n\mu) \\ -\frac{n}{2\sigma^2} + \frac{1}{2\sigma^4} \sum_{i=1}^n (x_i - \mu)^2 \end{bmatrix}.$$

So the possible candidates for $\hat{\mu}$ and $\hat{\sigma}^2$ is

$$\hat{\mu} = \bar{x} \quad \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2.$$

It remains to verify that $\hat{\mu}$ and $\hat{\sigma}^2$ are indeed the “maximizers”, note that

$$\begin{aligned} \frac{\partial^2 \ell}{\partial \mu^2} &= -\frac{n}{\sigma^2} \\ \frac{\partial^2 \ell}{\partial (\sigma^2)^2} &= \frac{n}{2\sigma^4} - \frac{1}{\sigma^8} \sum_{i=1}^n (x_i - \mu)^2 \\ \frac{\partial^2 \ell}{\partial \mu \partial \sigma^2} &= \frac{n\mu}{\sigma^4} - \frac{1}{\sigma^4} \sum_{i=1}^n x_i, \end{aligned}$$

and so the information matrix is

$$I(\mu, \sigma^2) = \begin{bmatrix} \frac{n}{\sigma^2} & \frac{n\mu}{\sigma^4} - \frac{1}{\sigma^4} \sum_{i=1}^n x_i \\ \frac{n\mu}{\sigma^4} - \frac{1}{\sigma^4} \sum_{i=1}^n x_i & \frac{1}{\sigma^8} \sum_{i=1}^n (x_i - \mu)^2 - \frac{n}{2\sigma^4} \end{bmatrix}.$$

Then the observed matrix is

$$\begin{aligned} I(\hat{\mu}, \hat{\sigma}^2) &= \begin{bmatrix} \frac{n}{\hat{\sigma}^2} & \frac{n\bar{x}}{(\hat{\sigma}^2)^2} - \frac{1}{(\hat{\sigma}^2)^2} \sum_{i=1}^n x_i \\ \frac{n\bar{x}}{(\hat{\sigma}^2)^2} - \frac{1}{(\hat{\sigma}^2)^2} \sum_{i=1}^n x_i & \frac{1}{(\hat{\sigma}^2)^4} \sum_{i=1}^n (x_i - \bar{x})^2 - \frac{n}{2(\hat{\sigma}^2)^2} \end{bmatrix} \\ &= \begin{bmatrix} \frac{n}{\hat{\sigma}^2} & 0 \\ 0 & \frac{1}{(\hat{\sigma}^2)^2} (\hat{\sigma}^2) - \frac{n}{2(\hat{\sigma}^2)^2} \end{bmatrix} \\ &= \begin{bmatrix} \frac{n}{\hat{\sigma}^2} & 0 \\ 0 & \frac{n}{(\hat{\sigma}^2)^2} \end{bmatrix}. \end{aligned}$$

We have that the determinant of the observed information matrix

is

$$\det I(\hat{\mu}, \hat{\sigma}^2) = \frac{n}{\hat{\sigma}^2} \cdot \frac{n}{(\hat{\sigma}^2)^2} > 0$$

since $\hat{\sigma}^2 > 0$.

2. By [Theorem 61](#), we have that the MLE of the coefficient of variation, CV , is³

$$\hat{CV} = \frac{\hat{\sigma}}{\hat{\mu}} = \frac{1}{\bar{x}} \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}} = \frac{\hat{\sigma}}{\sqrt{n\bar{x}}}$$

³Note that the function $f(x, y) = \frac{x}{y}$ is not injective if x and y are not coprime. Since both $\hat{\sigma}$ and $\hat{\mu}$ has a common multiple of $\sqrt{\frac{1}{n}}$, they are not coprime, so **I do not think we can actually invoke [Theorem 61](#)**. I shall, however, leave this answer here from the lecture.

20 Lecture 20 Jul 10th 2018

20.1 Estimation (Continued 3)

20.1.1 Maximum Likelihood Estimation (Continued 3)

Example 20.1.1 (Example 6.12 Part 2)

Continuing with Example 19.1.3, answer the following questions:

- Derive the Fisher information for the parameter vector (μ, σ^2) .

Solution

Recall that the information matrix was

$$I(\mu, \sigma^2) = \begin{bmatrix} \frac{n}{\sigma^2} & \frac{n\mu}{\sigma^4} - \frac{1}{\sigma^4} \sum_{i=1}^n x_i \\ \frac{n\mu}{\sigma^4} - \frac{1}{\sigma^4} \sum_{i=1}^n x_i & \frac{1}{\sigma^8} \sum_{i=1}^n (x_i - \mu)^2 - \frac{n}{2\sigma^4} \end{bmatrix}.$$

Thus the Fisher information matrix is

$$\begin{aligned} J(\mu, \sigma^2) &= \begin{bmatrix} E\left(\frac{n}{\sigma^2}\right) & E\left[\frac{n\mu}{\sigma^4} - \frac{1}{\sigma^4} \sum_{i=1}^n x_i\right] \\ E\left[\frac{n\mu}{\sigma^4} - \frac{1}{\sigma^4} \sum_{i=1}^n x_i\right] & E\left[\frac{1}{\sigma^8} \sum_{i=1}^n (x_i - \mu)^2 - \frac{n}{2\sigma^4}\right] \end{bmatrix} \\ &= \begin{bmatrix} \frac{n}{\sigma^2} & \frac{n\mu}{\sigma^4} - \frac{1}{\sigma^4} \sum_{i=1}^n E[X_i] = 0 \\ 0 & \frac{n}{\sigma^6} - \frac{n}{2\sigma^4} \end{bmatrix}. \end{aligned}$$

“ Note

Notice that we have, yet again, a similar scenario as when we first intro-

duced the Fisher information in the univariate case: notice that

$$\tilde{\mu} = \bar{X} \implies \text{Var}(\tilde{\mu}) = \text{Var}(\bar{X}) = \frac{\sigma^2}{n} = \frac{1}{J(\mu, \sigma^2)_{11}}$$

and

$$\begin{aligned} \tilde{\sigma}^2 &= \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 \implies \text{Var}(\tilde{\sigma}^2) = \frac{1}{n^2} \text{Var}((n-1)S^2) \\ &= \frac{\sigma^4}{n^2} \text{Var}\left(\frac{n-1}{\sigma^2} S^2\right) = \frac{\sigma^4 2(n-1)}{n^2} \end{aligned}$$

by Cochran's Theorem. Note that $\text{Var}(\tilde{\sigma}^2) \approx J(\mu, \sigma^2)_{22}$.

Example 20.1.2 (Example 6.13 (Course Notes 6.3.8))

Suppose $Y_i \sim N(\alpha + \beta x_i, \sigma^2)$, $i = 1, \dots, n$, is a sequence of IID rvs. Show that the ML estimators of α, β and σ^2 are given by

Note that $\tilde{\alpha}$ and $\tilde{\beta}$ are also the **least squares estimators** of α and β .

$$\begin{aligned} \tilde{\alpha} &= \bar{Y} - \tilde{\beta} \bar{x} \\ \tilde{\beta} &= \frac{\sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y})}{\sum_{i=1}^n (x_i - \bar{x})^2} \\ \tilde{\sigma}^2 &= \frac{1}{n} \sum_{i=1}^n (Y_i - \tilde{\alpha} - \tilde{\beta} x_i)^2 \end{aligned}$$

Solution

Since the Y_i 's constitute a random sample, the likelihood function is

$$\begin{aligned} L(\alpha, \beta, \sigma) &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(y_i - \alpha - \beta x_i)^2}{2\sigma^2}\right\} \\ &= (2\pi)^{-\frac{n}{2}} (\sigma^2)^{-\frac{n}{2}} \exp\left\{-\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \alpha - \beta x_i)^2\right\} \end{aligned}$$

and consequently the log-likelihood is

$$\ell(\alpha, \beta, \sigma) = -\frac{n}{2} \ln(2\pi) - \frac{n}{2} \ln \sigma^2 - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \alpha - \beta x_i)^2.$$

The first partial derivatives are

$$\frac{\partial \ell}{\partial \alpha} = \frac{1}{\sigma^2} \sum_{i=1}^n (y_i - \alpha - \beta x_i) = \frac{n}{\sigma^2} [\bar{y} - \alpha - \beta \bar{x}]$$

$$\frac{\partial \ell}{\partial \beta} = \frac{1}{\sigma^2} \sum_{i=1}^n [x_i (y_i - \alpha - \beta x_i)] = \frac{1}{\sigma^2} \left[\sum_{i=1}^n x_i y_i - \alpha \sum_{i=1}^n x_i - \beta \sum_{i=1}^n x_i^2 \right]$$

$$\frac{\partial \ell}{\partial \sigma} = -\frac{n}{\sigma} + \frac{1}{\sigma^3} \sum_{i=1}^n (y_i - \alpha - \beta x_i)^2.$$

The score vector is therefore

$$S(\alpha, \beta, \sigma) = \begin{bmatrix} \frac{n}{\sigma^2} [\bar{y} - \alpha - \beta \bar{x}] \\ \frac{1}{\sigma^2} [\sum_{i=1}^n x_i y_i - \alpha \sum_{i=1}^n x_i - \beta \sum_{i=1}^n x_i^2] \\ \frac{1}{\sigma^3} \sum_{i=1}^n (y_i - \alpha - \beta x_i)^2 - \frac{n}{\sigma} \end{bmatrix}.$$

To get the candidates for respective MLE,

$$\hat{\alpha} = \bar{y} - \hat{\beta} \bar{x}$$

$$\begin{aligned} \hat{\beta} &= \frac{\sum_{i=1}^n x_i y_i - \hat{\alpha} \sum_{i=1}^n x_i}{\sum_{i=1}^n x_i^2} = \frac{\sum_{i=1}^n x_i y_i - (\bar{y} - \hat{\beta} \bar{x}) \sum_{i=1}^n x_i}{\sum_{i=1}^n x_i^2} \\ &= \frac{\sum_{i=1}^n x_i y_i - n \bar{x} \bar{y} + n \hat{\beta} \bar{x}^2}{\sum_{i=1}^n x_i^2} = \frac{\sum_{i=1}^n x_i y_i - n \bar{x} \bar{y}}{\sum_{i=1}^n x_i^2 - n \bar{x}^2} \end{aligned}$$

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \alpha - \beta x_i)^2$$

For $\hat{\beta}$, note that

$$\begin{aligned} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) &= \sum_{i=1}^n [x_i y_i - \bar{x} y_i - \bar{y} x_i + \bar{x} \bar{y}] \\ &= \sum x_i y_i - \bar{x} \sum y_i - \bar{y} \sum x_i + n \bar{x} \bar{y} \\ &= \sum x_i y_i - \bar{x} (n \bar{y}) - \bar{y} (n \bar{x}) + n \bar{x} \bar{y} \\ &= \sum x_i y_i - n \bar{x} \bar{y} \end{aligned}$$

and similarly so we have

$$\sum_{i=1}^n (x_i - \bar{x})^2 = \sum_{i=1}^n x_i^2 - n \bar{x}^2.$$

Therefore

$$\hat{\beta} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

Thus the potential ML estimators are

$$\begin{aligned}\tilde{\alpha} &= \bar{Y} - \tilde{\beta}\bar{x} \\ \tilde{\beta} &= \frac{\sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y})}{\sum_{i=1}^n (x_i - \bar{x})^2} \\ \tilde{\sigma}^2 &= \frac{1}{n} \sum_{i=1}^n (Y_i - \tilde{\alpha} - \tilde{\beta}x_i)^2.\end{aligned}$$

To verify that these are indeed maximizing estimates, we need the following second order partial derivatives:

$$\begin{aligned}\frac{\partial^2 \ell}{\partial \alpha^2} &= -\frac{n}{\sigma^2} \\ \frac{\partial^2 \ell}{\partial \beta \partial \alpha} &= -\frac{n\bar{x}}{\sigma^2} \\ \frac{\partial^2 \ell}{\partial \sigma \partial \alpha} &= -\frac{2n}{\sigma^3} [\bar{y} - \alpha - \beta\bar{x}] \\ \frac{\partial^2 \ell}{\partial \alpha \partial \beta} &= -\frac{n\bar{x}}{\sigma^2} \\ \frac{\partial^2 \ell}{\partial \beta^2} &= -\frac{1}{\sigma^2} \sum_{i=1}^n x_i^2 \\ \frac{\partial^2 \ell}{\partial \sigma \partial \beta} &= -\frac{2}{\sigma^3} \left(\sum x_i y_i - \alpha \sum x_i - \beta \sum x_i^2 \right) \\ \frac{\partial^2 \ell}{\partial \alpha \partial \sigma} &= -\frac{2n}{\sigma^3} [\bar{y} - \alpha - \beta\bar{x}] \\ \frac{\partial^2 \ell}{\partial \beta \partial \sigma} &= -\frac{2}{\sigma^3} \left(\sum x_i y_i - \alpha \sum x_i - \beta \sum x_i^2 \right) \\ \frac{\partial^2 \ell}{\partial \sigma^2} &= \frac{n}{\sigma^2} - \frac{3}{\sigma^4} \sum_{i=1}^n (y_i - \alpha - \beta x_i)^2.\end{aligned}$$

It remains to show that the observed information matrix is positive definite, which will not be shown since the working is too tedious.

20.1.2 Asymptotic Properties of ML Estimators

Theorem 62 (Asymptotic Distribution of the ML Estimator)

Suppose $X = (X_1 \dots X_n)$ be a random sample from $f(x; \theta)$. Let $\tilde{\theta}_n = \tilde{\theta}_n(X_1, \dots, X_n)$ be the ML estimator of θ based on X . Then under certain regularity conditions¹:

¹ See Appendix A.

1. Consistency:

$$\tilde{\theta}_n \xrightarrow{P} \theta. \quad (20.1)$$

2. *Asymptotic Normality:*

$$\sqrt{J(\theta)}(\tilde{\theta}_n - \theta) \xrightarrow{D} Z \sim N(0,1). \tag{20.2}$$

3. *Asymptotic Distribution of Relative Likelihood:*

$$-2 \ln R(\theta; X) = 2[\ell(\tilde{\theta}_n; X) - \ell(\theta; X)] \xrightarrow{D} W \sim \chi^2(1)$$

where θ is the unknown true parameter. If **consistency** holds, then we call $\hat{\theta}_n$ a **consistent estimator** of θ .

For a proof of the above theorem see Casella & Berger 2002².

² Casella, G. & Berger, R. L. (2002). *Statistical Inference*. Thomson Learning

Note

The above theorem implies that for sufficiently large n , $\hat{\theta}_n$ has, approximately, an $N(\theta, [J(\theta)]^{-1})$ distribution (using methods from

Proposition 58). $[J(\theta)]^{-1}$ is called the **asymptotic variance** of $\hat{\theta}_n$. Consequently, for sufficiently large n , we have that

$$\text{Var}(\hat{\theta}_n) \approx [J(\theta)]^{-1}.$$

Note that $J(\theta)$ is **unknown since θ is unknown**.³ Since $\tilde{\theta}_n \xrightarrow{P} \theta$, by

Proposition 58, we have $J(\tilde{\theta}_n) \xrightarrow{P} J(\theta)$ ⁴. Then for large n , we have

$$\text{Var}(\tilde{\theta}_n) \approx [J(\theta)]^{-1} \approx [J(\tilde{\theta}_n)]^{-1}$$

and so $\tilde{\theta}_n \sim N(\theta, [J(\theta)]^{-1}) \approx N(\theta, [J(\tilde{\theta}_n)]^{-1})$, and therefore we get

$$[J(\tilde{\theta}_n)]^{\frac{1}{2}}(\tilde{\theta}_n - \theta) \xrightarrow{D} Z \sim N(0,1).$$

Now, note that the definition of the information function is

$$I(\theta; X) = - \sum_{i=1}^n \frac{d^2}{d\theta^2} \ell(\theta; X_i).$$

According to Geyer⁵, by the **Law of Large Numbers**,

$$\frac{1}{n} I(\tilde{\theta}) \xrightarrow{P} \frac{1}{n} J(\theta).$$

³ Comments from here on are my own after discussion with the lecturer. The explanation here will try to stay within the syllabus.

⁴ This is if J is continuous on both θ and $\tilde{\theta}_n$.

⁵ Geyer, C. J. (2007). Stat 5102 notes: Fisher information and confidence intervals using maximum likelihood. <https://www.stat.umn.edu/geyer/s06/5102/notes/fish.pdf>

Therefore, by Limit Theorems, we have

$$\frac{I(\tilde{\theta})}{J(\theta)} \xrightarrow{P} 1$$

Then by the Continuous Mapping Theorem, we have

$$[I(\tilde{\theta})]^{1/2} (\tilde{\theta} - \theta) \xrightarrow{D} Z \sim N(0, 1).$$

This result implies, by Limit Theorems, that for sufficiently large n , we have

$$\text{Var}(\tilde{\theta}_n) \approx [I(\hat{\theta})]^{-1}.$$

In the next section, we shall study how these results can be used to construct approximate **confidence intervals** for θ .

Example 20.1.3 (Example 6.14 (Course Notes 6.4.2))

Suppose X_1, \dots, X_n is a random sample from the $\text{Wei}(\theta, 2)$ distribution. Verify that the *consistency* and *Normal distribution assumption* hold for this distribution. Also, show that

$$[I(\tilde{\theta}_n; X)]^{1/2} (\tilde{\theta}_n - \theta) \xrightarrow{D} Z \sim N(0, 1).$$

Note that if $X \sim \text{Wei}(\theta, 2)$, then $E(X^k) = \theta^k \Gamma\left(\frac{k}{2} + 1\right)$ and its pf is

$$f_{X_i}(x_i) = \frac{2}{\theta^2} x_i e^{-(\frac{x_i}{\theta})^2}$$

Solution

Since the X_i 's form a random sample, we have that the likelihood function is

$$L(\theta) = \prod_{i=1}^n \frac{2}{\theta^2} x_i e^{-(\frac{x_i}{\theta})^2} = \left(\frac{2}{\theta^2}\right)^n e^{-\frac{1}{\theta^2} \sum_{i=1}^n x_i^2} \prod_{i=1}^n x_i.$$

Then the log-likelihood is

$$\ell(\theta) = n \ln 2 - 2n \ln \theta - \frac{1}{\theta^2} \sum_{i=1}^n x_i^2 + \sum_{i=1}^n \ln x_i.$$

To get a candidate for the MLE, we have

$$\frac{d\ell}{d\theta} = -\frac{2n}{\theta} + \frac{2}{\theta^3} \sum_{i=1}^n x_i^2,$$

and so

$$\hat{\theta}^2 = \frac{1}{n} \sum_{i=1}^n x_i \text{ and } \tilde{\theta}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2.$$

To verify that this candidate is indeed a maximum, note that we have

$$\frac{d^2 \ell}{d\theta^2} = \frac{2n}{\theta^2} - \frac{6}{\theta^4} \sum_{i=1}^n x_i^2. \quad (20.3)$$

Evaluating the above at $\hat{\theta}^2 = \frac{1}{n} \sum_{i=1}^n x_i^2$, we have

$$\frac{2n}{\frac{1}{n} \sum_{i=1}^n x_i^2} - \frac{6}{\left(\frac{1}{n}\right)^2 \left(\sum_{i=1}^n x_i^2\right)^2} \sum_{i=1}^n x_i^2 = -\frac{4n^2}{\sum_{i=1}^n x_i^2} < 0.$$

Thus $\hat{\theta}$ is indeed the maximum likelihood estimate. Now to show consistency, note that we have

$$E(X_i^2) = \theta^2 \Gamma(1+1) = \theta^2,$$

and so

$$E(\hat{\theta}^2) = E\left(\frac{1}{n} \sum_{i=1}^n X_i^2\right) = \frac{1}{n} \sum_{i=1}^n E(X_i^2) = \theta^2.$$

From here, to simplify notations, we shall write⁶

⁶Note that $\overline{X^2} \neq \bar{X}^2$.

$$\overline{X^2} = \frac{1}{n} \sum_{i=1}^n X_i^2 \text{ and } \sum x_i^2 = \sum_{i=1}^n x_i^2.$$

By the **Weak Law of Large Numbers**, we have that $\overline{X^2} \xrightarrow{P} \theta^2$. Since $\overline{X^2} \geq 0$, the function $g(x) = \sqrt{x}$ is continuous on its support, and so using **Proposition 58**, we have

$$\tilde{\theta} = \sqrt{\overline{X^2}} \xrightarrow{P} \theta.$$

To show normality, note that the Fisher information is

$$J(\theta) = E\left[-\frac{d^2}{d\theta^2} \ell(\theta)\right] = \frac{6}{\theta^4} \sum_{i=1}^n E(X_i^2) - \frac{2n}{\theta^2} = \frac{4n}{\theta^2} \quad (20.4)$$

where we shall note the use of **Equation (20.3)**. We want to invoke **CLT**, and so we also need $\text{Var}(\overline{X^2})$.

$$\text{Var}(\overline{X^2}) = \frac{1}{n^2} \sum_{i=1}^n \text{Var}(X_i^2) = \frac{1}{n^2} \sum_{i=1}^n \left[E(X_i^4) - [E(X_i^2)]^2 \right] = \frac{\theta^4}{n}.$$

Then by the CLT, we have

$$\frac{\sqrt{n}(\bar{X}^2 - \theta^2)}{\theta^2} \xrightarrow{D} Z \sim \mathbf{N}(0, 1).$$

By \heartsuit Proposition 58,

$$\sqrt{n}(\bar{X}^2 - \theta^2) \xrightarrow{D} \theta^2 Z \sim \mathbf{N}(0, \theta^4).$$

Using the same $g(x) = \sqrt{x}$ as above, by \spadesuit Corollary 60, we have

$$\sqrt{n}(\tilde{\theta} - \theta) \xrightarrow{D} \frac{\theta}{2} Z \sim \mathbf{N}\left(0, \frac{\theta^2}{4}\right)$$

Since Equation (20.4), using \heartsuit Proposition 58 again, we have

$$[J(\theta)]^{\frac{1}{2}} (\tilde{\theta} - \theta) = \frac{2\sqrt{n}}{\theta} (\tilde{\theta} - \theta) \xrightarrow{D} \frac{2}{\theta} \cdot \frac{\theta}{2} Z = Z \sim \mathbf{N}(0, 1)$$

Now from Equation (20.3), we have that the information function is

$$I(\theta; X) = \frac{6}{\theta^4} \sum_{i=1}^n X_i^2 - \frac{2n}{\theta^2}.$$

Then

$$I(\tilde{\theta}; X) = \frac{6n^2}{\left(\sum_{i=1}^n X_i^2\right)^2} \sum_{i=1}^n X_i^2 - \frac{2n^2}{\sum_{i=1}^n X_i^2} = \frac{4n^2}{\sum_{i=1}^n X_i^2} = \frac{4n}{\tilde{\theta}^2}.$$

Since $\tilde{\theta} \xrightarrow{P} \theta$, again by Limit Theorems, we have $\frac{\theta^2}{\tilde{\theta}^2} \xrightarrow{P} 1$, and so

$$\frac{\sqrt{I(\tilde{\theta}; X)}}{\sqrt{J(\theta)}} = \frac{\theta^2}{\tilde{\theta}^2} \xrightarrow{P} 1.$$

Then

$$[I(\tilde{\theta}; X)]^{\frac{1}{2}} (\tilde{\theta} - \theta) = \frac{\sqrt{I(\tilde{\theta}; X)}}{\sqrt{J(\theta)}} [J(\theta)]^{\frac{1}{2}} (\tilde{\theta} - \theta) \xrightarrow{D} 1 \cdot Z = Z \sim \mathbf{N}(0, 1).$$

21 Lecture 21 Jul 12th 2018

21.1 Estimation (Continued 4)

21.1.1 Asymptotic Properties of ML Estimators (Continued)

Example 21.1.1 (Example 6.15 (Course Notes 6.4.3))

Suppose X_1, \dots, X_n is a random sample from the $\text{Unif}(0, \theta)$ distribution. We showed in Example 18.1.5 that $\tilde{\theta}_n = X_{(n)}$ is the ML estimator of θ . Since the support of X_i depends on θ , Theorem 62 does not hold. Show, however, that $X_{(n)}$ is still a consistent estimator of θ .

Solution

Since θ is a fixed (but unknown) value, we have the option of use

 Proposition 55. We have


$$\begin{aligned} F_{X_{(n)}}(t) &= P(X_{(n)} \leq t) = P(X_1 \leq t, X_2 \leq t, \dots, X_n \leq t) \\ &= \prod_{i=1}^n P(X_i \leq t) \quad \because \text{independence} \\ &= \prod_{i=1}^n \frac{t}{\theta} = \left(\frac{t}{\theta}\right)^n \end{aligned}$$

So the full expression for the CDF of $X_{(n)}$ is

$$F_{X_{(n)}}(t) = \begin{cases} 0 & t \leq 0 \\ \left(\frac{t}{\theta}\right)^n & 0 < t < \theta \\ 1 & t \geq \theta \end{cases}$$

Observe that

$$\lim_{n \rightarrow \infty} F_{X_{(n)}}(t) = \begin{cases} 0 & t < \theta \\ 1 & t \geq \theta \end{cases}$$

and so as $n \rightarrow \infty$, we have that $X_{(n)}$ has a degenerate distribution, and so the proof for consistency follows from  Proposition 55.

21.1.2 Interval Estimators

Definition 70 (Interval Estimators)

Suppose X is an rv whose distribution depends on θ . Suppose that $A(x)$ and $B(x)$ are functions such that $A(x) \leq B(x)$ for all $x \in \text{supp}(X)$ and $\theta \in \Omega$. Let x be the observed data. Then $(A(x), B(x))$ is an **interval estimate** for θ . The interval $(A(X), B(X))$ is an **interval estimator** for θ .

Note

Two interval estimators that we already know¹:

1. Confidence intervals
2. Likelihood intervals

¹ Well, one from STAT231, that is.

WE NOW CONSIDER a general approach for constructing confidence intervals based on pivotal quantities.

Definition 71 (Pivotal Quantity)

Suppose X is an rv whose distribution depends on θ . The rv $Q(X; \theta)$ is called a **pivotal quantity** if the distribution of Q does not depend on θ . $Q(X; \theta)$ is called an **asymptotic pivotal quantity** if the limiting distribution of Q as $n \rightarrow \infty$ does not depend on θ .

Example 21.1.2 (Example 6.16)

Suppose X_1, \dots, X_n is a random sample from the $N(\mu, \sigma^2)$ distribution. The following rvs are pivotal quantities.:

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}}, \frac{\bar{X} - \mu}{S/\sqrt{n}}, \frac{\sum_{i=1}^n (X_i - \mu)^2}{\sigma^2} \text{ and } \frac{(n-1)S^2}{\sigma^2} = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{\sigma^2}$$

Exercise 21.1.1
Show that the given rvs are pivotal quantities.


Example 21.1.3 (Example 6.17 (Course Notes 6.5.4))

Suppose X_1, \dots, X_n is a random sample from the $\text{Poi}(\theta)$ distribution. Show that


$$\sqrt{n} \frac{(\bar{X}_n - \theta)}{\sqrt{\bar{X}_n}}$$

is an asymptotic pivotal quantity.

 **Solution**

By  Theorem 57, we have that


$$\sqrt{n} \frac{\bar{X}_n - \theta}{\sqrt{\theta}} \xrightarrow{D} Z \sim N(0, 1).$$

By  Proposition 58, we have

$$\sqrt{n}(\bar{X}_n - \theta) \xrightarrow{D} \sqrt{\theta}Z \sim N(0, \theta).$$

Now by the **Weak Law of Large Numbers**, we have that

$$\bar{X}_n \xrightarrow{P} \theta.$$


Since \bar{X}_n takes only non-negative values, and $\theta > 0$, the function $g(x) = \sqrt{x}$ is well-defined on these “points”, and so by  Proposition 58, we have

$$\sqrt{\bar{X}_n} \xrightarrow{P} \sqrt{\theta}.$$

Then by Slutsky’s Theorem, we have

$$\sqrt{n} \frac{\bar{X}_n - \theta}{\sqrt{\bar{X}_n}} \xrightarrow{D} Z \sim N(0, 1).$$

We observe that the limiting distribution of $\sqrt{n}(\bar{X}_n - \theta) / \sqrt{\bar{X}_n}$ depends not on the unknown parameter θ . Therefore it is a asymptotic pivotal quantity.

This may be a little late but at this point, it should be clear how Limit Theorems work when it comes to playing with the Normal distribution as the asymptotic distribution. If this is not clear to you, read back from  Proposition 58.

 **Definition 72 (Confidence Interval)**

Suppose $A(X)$ and $B(X)$ are statistics, and are called the **lower** and

upper confidence limits, respectively. If $P[A(X) < \theta < B(X)] = p$, where $0 < p < 1$, then $(a(x), b(x))$ is called a $100p\%$ **confidence interval (CI)** for θ .

Pivotal quantities can be used for constructing CIs in the following way: since the distribution of $Q(X; \theta)$ is known, we can write down a probability statement of the form

$$P(q_1 \leq Q(X; \theta) \leq q_2) = p.$$

If Q is a monotone function of θ , then this statement can be rewritten as

$$P[A(X) \leq \theta \leq B(X)] = p,$$

and the “realized” interval $[a(x), b(x)]$ is a $100p\%$ CI.

Example 21.1.4 (Example 6.18 (Course Notes 6.5.6))

Suppose X_1, \dots, X_n is a random sample from the $N(\mu, \sigma^2)$ distribution. Use the pivotal quantities in Example 21.1.2 to find:

1. a $100p\%$ CI for μ if σ^2 is unknown;
2. a $100p\%$ CI for σ^2 if μ is known.


Solution

1. Since σ^2 is unknown, we cannot use

$$\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

as our pivotal quantity. We shall in fact, use

$$\frac{\bar{X}_n - \mu}{S/\sqrt{n}} \sim t(n-1)$$

since it contains the unknown parameter whose confidence interval is that of what we want to construct, where we know that the rv follows a t -Distribution with degrees of freedom $n-1$ from  Theorem 47. Then since the t -Distribution is symmetric², we have

$$P\left[-a \leq \frac{\bar{X}_n - \mu}{S/\sqrt{n}} \leq a\right] = p$$

$$P\left[\bar{X}_n - (S/\sqrt{n})a \leq \mu \leq \bar{X}_n + (S/\sqrt{n})a\right] = p.$$

² See Wikipedia for a graph of the t -Distribution.

And so the $100p\%$ CI for μ when σ^2 is unknown is

$$[\bar{X}_n - (S/\sqrt{n})a, \bar{X}_n + (S/\sqrt{n})a]$$

2. Since μ is known, we shall use the rv

$$\frac{\sum_{i=1}^n (X_i - \mu)^2}{\sigma^2} \sim \chi^2(n-1),$$

where the following of the chi-squared Distribution is from Cochran's Theorem. To construct the CI for σ^2 ,

$$P \left(a \leq \frac{\sum_{i=1}^n (X_i - \mu)^2}{\sigma^2} \leq b \right) = p$$

$$P \left(\frac{1}{b} \sum_{i=1}^n (X_i - \mu)^2 \leq \sigma^2 \leq \frac{1}{a} \sum_{i=1}^n (X_i - \mu)^2 \right) = p.$$

Thus the CI for σ^2 is

$$\left[\frac{1}{b} \sum_{i=1}^n (X_i - \mu)^2, \frac{1}{a} \sum_{i=1}^n (X_i - \mu)^2 \right].$$

◆ **Proposition 63 (MLE of a Location/Scale Parameter as a Pivotal Quantity)**

Let $X = (X_1, \dots, X_n)$ be a random sample from $f(x; \theta)$ and let $\tilde{\theta} = \tilde{\theta}(X)$ be the ML estimator of the scalar parameter θ based on X .


1. If θ is a location parameter, then $Q = \tilde{\theta} - \theta$ is a pivotal quantity.
2. If θ is a scale parameter, then $Q = \tilde{\theta}/\theta$ is a pivotal quantity.

There is a proof out there and I wish to hunt for it but I do not have the time.

Example 21.1.5 (Example 6.19)


Suppose $X_1, \dots, X_n \sim \text{Exp}(\theta)$ is a random sample. Find a $100p\%$ equal tails CI for θ . For the data $n = 15$ and $\sum_{i=1}^{15} x_i = 36$, find a 95% equal tail CI for θ .

 **Solution**

Referring to Example 3.3.3, we know that $\text{Exp}(\theta)$ belongs to a scale parameter family of distributions. We shall proceed with finding the MLE of θ and then use  Proposition 63 to get a CI for θ . Since the X_i 's are IID,

$$\begin{aligned} L(\theta) &= \prod_{i=1}^n \frac{1}{\theta} e^{-\frac{x_i}{\theta}} = \theta^{-n} \exp \left\{ -\frac{1}{\theta} \sum_{i=1}^n x_i \right\} \\ \ell(\theta) &= -n \ln \theta - \frac{1}{\theta} \sum_{i=1}^n x_i \\ \frac{d\ell}{d\theta} &= -\frac{n}{\theta} + \frac{1}{\theta^2} \sum_{i=1}^n x_i \\ \implies \hat{\theta} &= \frac{1}{n} \sum_{i=1}^n x_i = \bar{x}. \end{aligned}$$

Thus the MLE of θ is $\tilde{\theta} = \bar{X}$. To get the 100p% CI for θ , we have by

 Proposition 63, $Y = \bar{X}/\theta$ is a pivotal quantity. Then

$$\begin{aligned} P \left[a \leq \frac{\bar{X}}{\theta} \leq b \right] &= p \\ P \left[\frac{1}{b} \bar{X} \leq \theta \leq \frac{1}{a} \bar{X} \right] &= p. \end{aligned}$$

And so the 100p% CI for θ is

$$\left[\frac{1}{b} \bar{X}, \frac{1}{a} \bar{X} \right].$$

To solve for a and b , we would need to know what the distribution of $Y = \bar{X}/\theta$ is:

$$\begin{aligned} M_Y(t) &= E \left(\exp \left\{ \frac{t}{\theta n} \sum_{i=1}^n X_i \right\} \right) = E \left(e^{\frac{tX_1}{n\theta}} \right)^n \quad \because \text{IID} \\ &= M_{X_1} \left(\frac{t}{n\theta} \right)^n = \left(1 - \frac{t}{n} \right)^{-n}, \end{aligned}$$

which we observe that then $Y \sim \text{Gam} \left(n, \frac{1}{n} \right)$.

For $n = 15$, $\sum_{i=1}^{15} x_i = 36$ (and so $\bar{x} = 36/15 = 2.4$), and to find the 95% equal tail CI for θ , we have that

$$\begin{aligned} P \left[0.559 \leq \frac{2.4}{\theta} \leq 1.566 \right] &= 0.95 \\ P [2.4/1.566 \leq \theta \leq 2.4/0.559] &= 0.95. \end{aligned}$$

where the values of a and b are verified on R using the following:

Listing 21.1: R interactive code to get a and b

```
qgamma(0.025, shape=15, rate=15)
qgamma(0.975, shape=15, rate=15)
```

Thus

$$\left[\frac{2.4}{1.566}, \frac{2.4}{0.559} \right]$$


$$[1.533, 4.293]$$

Example 21.1.6 (Example 6.20 (Course Notes 6.5.11))

Suppose $X_1, \dots, X_n \sim \text{Unif}(0, \theta)$ is a random sample. Find an appropriate pivotal quantity. Determine a such that

$$[\hat{\theta}, a\hat{\theta}]$$

is a $100p\%$ CI for θ .

 **Solution**

Since the X_i 's are IID, we have that

$$L(\theta) = \prod_{i=1}^n \frac{1}{\theta} \mathbb{1}_{\{0 \leq x_i \leq \theta\}} = \theta^{-n} \mathbb{1}_{\{0 \leq x_{(1)} \leq x_{(n)} \leq \theta\}}$$

$$\ell(\theta) = -n \ln \theta + \ln \mathbb{1}_{\{0 \leq x_{(1)} \leq x_{(n)} \leq \theta\}}.$$

Using a similar argument as in [Example 19.1.2](#), notice that $-n \ln \theta$ is a decreasing function, and so to get the maximum likelihood estimate, we want θ to be as close to 0 as possible, but the closest θ can get to is $x_{(n)}$. Note that if the indicator function turns out to be 1, then θ must also be 1, but this does not maximize the likelihood function.

So we have $\hat{\theta} = x_{(n)}$ and $\tilde{\theta} = X_{(n)}$. Note that $\text{Unif}(0, \theta)$ is from a scale parameter family of distributions since

$$f_{X_i}(x_i) = \frac{1}{\theta} \mathbb{1}_{\{0 \leq x_i \leq \theta\}} = \frac{1}{\theta} f_{X_i}\left(\frac{x_i}{\theta}\right).$$

Then by  [Proposition 63](#), $Y = \tilde{\theta}/\theta = X_{(n)}/\theta$ is a pivotal quantity.

22 Lecture 22 Jul 17th 2018

22.1 Estimate (Continued 5)

22.1.1 Interval Estimators (Continued)

Example 22.1.1 (Example 6.21)

Suppose $X_1, \dots, X_n \sim \text{Exp}(1, \theta)$ is a random sample. Find an appropriate pivotal quantity. Determine a such that


$$[\hat{\theta} - a, \hat{\theta}]$$

is a $100p\%$ CI for θ .

Solution

Recall from *Example 19.1.2* that the ML estimate is $\hat{\theta} = x_{(1)}$ and the estimator is $\tilde{\theta} = X_{(1)}$. Note that $\text{Exp}(1, \theta)$ is from a location parameter family of distributions, since

$$f(x; 1, 0) = e^{-x} = f(x - \theta; 1, \theta).$$

We can then use  *Proposition 63*, and use $Y = \tilde{\theta} - \theta$ as our pivotal quantity. Note that

$$\begin{aligned}P(\hat{\theta} - a \leq \theta \leq \hat{\theta}) &= p \\P(-\hat{\theta} \leq -\theta \leq a - \hat{\theta}) &= p \\P(0 \leq \hat{\theta} - \theta \leq a) &= p \\P(0 \leq Y \leq a) &= p\end{aligned}$$

It remains, again, to find the CDF.

$$\begin{aligned}
 F_Y(y) &= P(X_{(1)} - \theta \leq y) = 1 - P(X_{(1)} > y + \theta) \\
 &= 1 - P(X_1 > y + \theta)^n \quad \because \text{IID} \\
 &= 1 - \left[\int_{y+\theta}^{\infty} e^{-(x+\theta)} dx \right]^n \\
 &= 1 - \left[-e^{-(x+\theta)} \Big|_{y+\theta}^{\infty} \right]^n \\
 &= 1 - e^{-ny}
 \end{aligned}$$

Thus

$$\begin{aligned}
 F(a) - F(0) &= p \\
 1 - e^{-na} - 1 + 1 &= p \\
 a &= -\frac{1}{n} \ln(1 - p)
 \end{aligned}$$

IN CASES WHERE we cannot construct an exact pivotal quantity, we can use the limiting distribution of the ML estimator $\tilde{\theta} = \tilde{\theta}(X_1, \dots, X_n)$ and construct approximate CIs.

♦ Proposition 64 (Asymptotic Confidence Intervals)


Since¹

$$[J(\tilde{\theta})]^{1/2} (\tilde{\theta} - \theta) \xrightarrow{D} Z \sim N(0, 1),$$

then $[J(\tilde{\theta})]^{1/2} (\tilde{\theta} - \theta)$ is an asymptotic pivotal quantity. An **asymptotic 100p% CI**² based on this asymptotic pivotal quantity is given by

$$[\hat{\theta} - a[J(\hat{\theta})]^{-1/2}, \hat{\theta} + a[J(\hat{\theta})]^{-1/2}]$$

where $P(-a < Z < a) = p$ and $Z \sim N(0, 1)$.

¹ This is based on  Theorem 62, which we could not prove, and its consequences are more than just fishy. I am also unsure whether this is presented as a proof or definition in the lecture, since there are no indications of that whatsoever both during the lectures and in the course notes.

² Also called **approximate 100p% CI**.

“ Note

Similarly, since

$$[I(\tilde{\theta})]^{1/2} (\tilde{\theta} - \theta) \xrightarrow{D} Z \sim N(0, 1),$$

then $[I(\tilde{\theta})]^{1/2} (\tilde{\theta} - \theta)$ is an asymptotic pivotal quantity. An asymptotic 100p% CI based on this asymptotic pivotal quantity is given by

$$[\hat{\theta} - a[I(\hat{\theta})]^{-1/2}, \hat{\theta} + a[I(\hat{\theta})]^{-1/2}]$$

where $I(\hat{\theta})$ is the observed information.

Example 22.1.2 (Example 6.22)

Suppose $X \sim \text{Bin}(n, \theta)$. Show how you would construct an approximate 100p% CI for θ .

Solution

Since this is just a single rv, we have

$$\begin{aligned} L(\theta) &= \binom{n}{x} \theta^x (1-\theta)^{n-x} \\ \ell(\theta) &= \ln \binom{n}{x} + x \ln \theta + (n-x) \ln(1-\theta) \\ \frac{d\ell}{d\theta} &= \frac{x}{\theta} - \frac{n-x}{1-\theta} \\ \frac{d^2\ell}{d\theta^2} &= -\frac{x}{\theta^2} - \frac{n-x}{(1-\theta)^2} < 0 \end{aligned}$$

where we note that the ML estimate is $\hat{\theta} = \frac{x}{n}$ and the estimator $\tilde{\theta} = \frac{X}{n}$, and we know that this will be the maximum since the second order derivative is always negative³. The information function is


$$I(\theta; X) = \frac{X}{\theta} + \frac{n-X}{(1-\theta)^2}$$

and so the Fisher information is

$$J(\theta) = \frac{E(X)}{\theta^2} + \frac{n-E(X)}{(1-\theta)^2} = \frac{n}{\theta} + \frac{n}{1-\theta} = \frac{n}{\theta(1-\theta)}.$$

Thus

$$J(\hat{\theta}) = \frac{n^3}{x(n-x)}.$$

Then by  Proposition 64, we have that the asymptotic 100p% CI for θ is

$$\left[\frac{x}{n} - zn \sqrt{\frac{x(n-x)}{n}}, \frac{x}{n} + zn \sqrt{\frac{x(n-x)}{n}} \right],$$

where $P(-z \leq Z \leq z) = p$ where $Z \sim N(0, 1)$.

Example 22.1.3 (Example 6.23 (Course Notes 6.5.15))

Suppose X_1, \dots, X_n is a random sample from the $\text{Poi}(\theta)$ distribution. Show how you would construct an asymptotic 100p% CI for θ .

³ In fact, by rearranging $\frac{d\ell}{d\theta}$, we can show that it is a linear function.

 **Solution**

Since the X_i 's are IID, we have

$$L(\theta) = \prod_{i=1}^n \frac{e^{-\theta} \theta^{x_i}}{x_i!} = e^{-n\theta} \theta^{\sum_{i=1}^n x_i} \prod_{i=1}^n \frac{1}{x_i!}$$

$$\ell(\theta) = -n\theta + \left(\sum_{i=1}^n x_i \right) \ln \theta - \sum_{i=1}^n \ln x_i!$$

$$\frac{d\ell}{d\theta} = -n + \frac{1}{\theta} \sum_{i=1}^n x_i$$

$$\frac{d^2\ell}{d\theta^2} = -\frac{1}{\theta^2} \sum_{i=1}^n x_i.$$

The candidate for MLE from equating the score function to 0 is $\hat{\theta} = \frac{1}{n} \sum_{i=1}^n x_i = \bar{x}$. We verify that it is indeed a maximum point since

$$\left. \frac{d^2\ell}{d\theta^2} \right|_{\theta=\hat{\theta}} = -\frac{1}{\bar{x}} \sum_{i=1}^n x_i = -\frac{1}{\bar{x}} < 0.$$

The information function is


$$I(\theta; X) = \frac{1}{\theta^2} \sum_{i=1}^n X_i$$

and so the Fisher information is

$$J(\theta) = E[I(\theta; X)] = \frac{1}{\theta^2} n\theta = \frac{n}{\theta}.$$

Thus

$$J(\hat{\theta}) = \frac{n}{\bar{x}}$$

Assuming that our random sample satisfies the regularity conditions, we have that the asymptotic 100p% CI for θ , by  Proposition 64, is

$$\left[\bar{x} - z \sqrt{\frac{\bar{x}}{n}}, \bar{x} + z \sqrt{\frac{\bar{x}}{n}} \right]$$

 **Definition 73 (Likelihood Interval)**

Given a random sample X_1, \dots, X_n from $f(x; \theta)$, the 100p% **likelihood interval** for θ is defined by

$$\{\theta : R(\theta) > p\}.$$

“ Note

Recall the 3rd property from  Theorem 6.2 that

$$-2 \log R(\theta; X) \stackrel{D}{\rightarrow} W \sim \chi^2(1).$$

Since the $R(\theta)$ is a unimodal graph, we have

$$\begin{aligned} P[R(\theta; X) \geq p] &= P[-2 \ln R(\theta; X) \leq -2 \ln p] \\ &\approx P(W \leq -2 \ln p) \\ &= P(Z^2 \leq -2 \ln p) \\ &= P\left(-\sqrt{-2 \ln p} \leq Z \leq \sqrt{-2 \ln p}\right) \\ &= 2P\left(Z \leq \sqrt{-2 \ln p}\right) - 1. \end{aligned}$$

Consequently, if $p = 0.15$, then

$$P[R(\theta; X) \geq 0.15] \approx 0.95$$

i.e. a 15% likelihood interval is, approximately, a 95% CI.

The amount of things built on this shady theorem is insane...

Example 22.1.4 (Example 6.24)

In Example 19.1.1, based on $X_1, \dots, X_{100} \sim \text{Poi}(\theta)$ is a random sample, where $\sum_{i=1}^{100} x_i = 980$, we derived the 10% and 50% likelihood intervals for θ . Compare the approximate 95% confidence interval based on the asymptotic distribution of the MLE to the 15% likelihood interval for θ .

 Solution

Recall that

$$R(\theta) = e^{-n(\theta - \bar{x})} \left(\frac{\theta}{\bar{x}}\right)^{n\bar{x}} = e^{-100(\theta - 9.8)} \left(\frac{\theta}{980}\right)^{980}$$

and that the MLE is $\hat{\theta} = \bar{x}$. Using R⁴, one can find that the 15% likelihood interval is

$$9.2 < \theta < 10.4.$$

⁴ This is actually a non-trivial process...

Again, assuming that the regularity conditions are satisfied, by

23 Lecture 23 Jul 19th 2018

23.1 Hypothesis Testing

23.1.1 Introduction

Definition 74 (Hypothesis)

In inferential statistics, given a model $f(x; \theta)$, where $\theta = (\theta_1 \dots \theta_n)^T \in \Omega$, and Ω is a parameter space, a **hypothesis** is a general statement about the parameters for the model, denoted by

$$H : \theta \in \Omega_0 \subseteq \Omega.$$

Definition 75 (Null Hypothesis and Alternative Hypothesis)

A **null hypothesis** is a hypothesis of which its default position is that there is no relationship between the two phenomena measured by the model $f(x; \theta)$. We usually denote the null hypothesis by

$$H_0 : \theta \in \Omega_0.$$

The **alternative hypothesis** is the other case of the null hypothesis; it is a hypothesis that states that there is a relationship between the two phenomena measured by the model $f(x; \theta)$. We usually denote the alternative hypothesis as

$$H_a : \theta \notin \Omega_0.$$

Definition 76 (Test of Hypothesis)

A **test of hypothesis** is a procedure for evaluating the strength of the evidence provided by the data against the null hypothesis.

To measure the evidence against H_0 based on the observed data, we use a test statistic or a discrepancy measure.

Definition 77 (Test Statistic)

A **test statistic** is a statistic that is used for measuring the evidence against H_0 using the observed data.

Note

1. A small observed value of the test statistic shows close agreement between the observed data and the null hypothesis; while
2. A large observed value of the test statistic shows poor agreement.

Definition 78 (Significance Level and p -Values)

Assuming that H_0 is true, we compute the probability of observing a value of the test statistic being greater than or equal to the actual observed value. This probability is called the **significance level** or **p -value** of the data in relation with the null hypothesis.

Note

The p -value is the probability of observing a poor agreement between the null hypothesis and the data.

- A small p -value indicates that the probability that our observed test statistic occurs under the null hypothesis is low, i.e. an **evidence against the null hypothesis**;
- A large p -value indicates that the probability that our observed test statistic occurs under the null hypothesis is high, i.e. we have **little evidence against the null hypothesis**.

This also means that the smaller the p -value, the stronger the evidence against the null hypothesis.

There are two types of errors that we may run into while performing hypothesis testing:

Definition 79 (Type I and Type II Errors)

A **Type I Error** is the case where we reject a true null hypothesis. A **Type II Error** is the case where we fail to reject a false null hypothesis.

	$H_0 : \theta \in \Omega_0$	$H_a : \theta \notin \Omega_0$
Reject H_0	Type I Error	✓ (True Negatives)
Fail to Reject H_0	✓ (True Positives)	Type II Error

Note

The common practice is to reduce the occurrence of a Type I Error. It is analogous to the **Presumption of Innocence**, where the “null hypothesis” is “correct until proven wrong”. In other words, the goal of this practice is to reduce the occurrence of **false positives**.

Example 23.1.1 (Example 7.11 (Course Notes 7.1.1))

Suppose $X_1, \dots, X_n \sim N(\mu, \sigma^2)$ is a random sample, where σ^2 is known. Suppose $n = 25$, $\sigma = 1$, $\bar{x} = 0.5$, and $\mu_0 = 0$. Perform a test of hypothesis on the null hypothesis $H_0 : \mu = \mu_0$ and state the conclusion.

Solution

By the CLT, we have

$$Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

which contains all the parameters of our interest and is a statistic¹.

¹ Note that it is also a pivotal quantity.

We shall use this as a test statistic. Under H_0 , we have $\mu = \mu_0$, i.e.

$$\frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1).$$

Then, the observed test statistic is

$$z_0 = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} = \frac{0.5 - 0}{1/\sqrt{25}} = 2.5,$$

and so the p -value is

$$p = 2P(Z \geq 2.5) \approx 0.01241923$$

by the Z-table. Therefore, we observe that the p -value is small, and so we have evidence against the null hypothesis $H_0 : \mu_0 = 0$.

23.1.2 Likelihood Ratio Tests for Simple Hypothesis

Definition 80 (Simple Hypothesis)

Suppose X_1, \dots, X_n is a random sample from $f(x; \theta)$, where $\theta = (\theta_1 \dots \theta_k)^T$. Suppose we wish to test $H : \theta = \theta_0$, where θ_0 is a completely known $k \times 1$ vector. This hypothesis is called a **simple hypothesis**, which specifies the values of all unknown parameters in the mode.

Definition 81 (Likelihood Ratio Statistic)

The **likelihood ratio (LR) statistic** is defined as

$$\Lambda(\theta_0) = -2 \ln R(\theta_0; X) = -2 \ln \left[\frac{L(\theta_0; X)}{L(\tilde{\theta}; X)} \right] = 2 [\ell(\tilde{\theta}; X) - \ell(\theta_0; X)],$$

where $X = (X_1, \dots, X_n)$ are the data and $\tilde{\theta} = \tilde{\theta}(X_1, \dots, X_n)$ is the ML estimator of θ . The observed likelihood ratio statistic is thus

$$\lambda(\theta_0) = -2 \ln R(\theta_0; x)$$

where $x = (x_1, \dots, x_n)$.

Assuming H_0 is true, under certain regularity conditions², we have

$$\Lambda(\theta_0) = -2 \ln R(\theta_0, X) \xrightarrow{D} W \sim \chi^2(k).$$

² This probably involves  Theorem 62

where k is the number of parameters under the general hypothesis minus the number of parameters under H_0 . Thus an approximate p -value is given by

$$p \approx P(W \geq -2 \ln R(\theta_0; x)),$$

where $x = (x_1, \dots, x_n)$ is the observed data.

Example 23.1.2 (Example 7.2 (Course Notes 7.2.1))

Continuing with Example 23.1.1, find the LR statistic and compare it to the test statistic used in the example.

 **Solution**

Let $H_0 : \mu = \mu_0$ and $H_a : \mu \neq \mu_0$. We need the MLE of μ . Since the X_i 's forms a random sample, we have

$$\begin{aligned} L(\mu) &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x_i - \mu)^2}{2\sigma^2}} = (2\pi)^{-\frac{n}{2}} \sigma^{-n} \exp \left\{ -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \right\} \\ \ell(\mu) &= -\frac{n}{2} \ln(2\pi) - n \ln \sigma - \frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \\ \frac{d\ell}{d\mu} &= \frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu) = \frac{1}{\sigma^2} (\sum_{i=1}^n x_i - n\mu) \\ \frac{d^2\ell}{d\mu^2} &= -\frac{n}{\sigma^2} < 0 \end{aligned}$$

and so the ML estimate for μ is

$$\hat{\mu} = \frac{1}{n} \sum_{i=1}^n x_i = \bar{x}.$$

Thus the likelihood ratio statistic is

$$\begin{aligned} \Lambda(\mu_0) &= -2 \ln R(\mu_0, X) = 2[\ell(\hat{\mu}; X) - \ell(\mu_0; X)] \\ &= \frac{1}{\sigma^2} \left(\sum_{i=1}^n (X_i - \mu_0)^2 - \sum_{i=1}^n (X_i - \hat{\mu})^2 \right) \\ &= \frac{n}{\sigma^2} \left[2\bar{x}(\hat{\mu} - \mu_0) + \mu_0^2 - \hat{\mu}^2 \right] \end{aligned}$$

By our argument before this example, we have that

$$\begin{aligned} k &= \text{number of unknown parameters} \\ &\quad - \text{number of unknown parameters under } H_0 \\ &= 1 - 0 = 1. \end{aligned}$$

Thus

$$\Lambda(\theta_0) \stackrel{D}{\rightarrow} W \sim \chi^2(1).$$

The observed likelihood ratio statistic is

$$\begin{aligned} \lambda(\mu_0) &= \frac{n}{\sigma^2} [2\bar{x}(\bar{x} - \mu_0) + \mu_0^2 - \bar{x}^2] = \frac{n(\bar{x}^2 - 2\mu_0\bar{x} + \mu_0^2)}{\sigma^2} \\ &= \frac{n(\bar{x} - \mu_0)^2}{\sigma^2}. \end{aligned}$$

The p -value is therefore

$$p \approx P(W \geq \lambda(\mu_0)) = P\left(W \geq \frac{n(\bar{x} - \mu_0)^2}{\sigma^2}\right).$$

Using the values given in Example 23.1.1, we have

$$p \approx P(W \geq 6.25) \approx 0.01241923,$$

where our value is obtained by the R code:

```
1 - pchisq(6.25, 1)
```

We can now conclude that the two tests are similar.

Example 23.1.3 (Example 7.3 (Course Notes 7.2.4))

Suppose X_1, \dots, X_n is a random sample from the $\text{Poi}(\theta)$ distribution.

Find the LR statistic for testing $H : \theta = \theta_0$. Another test statistic

which could be used is

$$T = \frac{|\bar{X} - \theta_0|}{\sqrt{\theta_0/n}}.$$

What is the approximate distribution of

$$\frac{\bar{X} - \theta_0}{\sqrt{\theta_0/n}}$$

for large n if H_0 is true, and how could you use this to find an approximate p -value?

Solution

Let $H_0 : \theta \in \Omega_0$ and $H_a : \theta \notin \Omega_0$. By Example 22.1.3, we have that

$\hat{\theta} = \bar{x}$, and the log-likelihood is

$$\ell(\theta) = -n\theta + \left(\sum_{i=1}^n x_i\right) \ln \theta - \sum_{i=1}^n \ln x_i! .$$

Thus the likelihood ratio statistic is

$$\Lambda(\theta_0) = 2[n\bar{X} \ln \tilde{\theta} - n\bar{X} \ln \theta_0] = 2n\bar{X} \ln \left(\frac{\tilde{\theta}}{\theta_0} \right)$$

and so the observed likelihood ratio statistic is

$$\lambda(\theta_0) = 2n\bar{x} \ln \left(\frac{\bar{x}}{\theta_0} \right).$$

Thus the p -value under the assumption of H_0 is

$$p \approx P(W \geq \lambda(\theta_0)).$$

On the other hand, by the CLT, we have that

$$\frac{\bar{X} - \theta_0}{\sqrt{\theta_0/n}} \xrightarrow{D} Z \sim N(0, 1),$$

which then the p -value under this method is

$$p \approx 2P \left(Z \geq \left| \frac{\bar{x} - \theta_0}{\sqrt{\theta_0/n}} \right| \right).$$

However, it is not necessary that

$$P(W \geq \lambda(\theta_0)) = 2P \left(Z \geq \left| \frac{\bar{x} - \theta_0}{\sqrt{\theta_0/n}} \right| \right)$$

Example 23.1.4 (Example 7.4 (Course Notes 7.2.6))

The following table gives the observed frequencies of the six faces in 100 rolls of a die:

Face: i	1	2	3	4	5	6	Total
Freq: f_i	16	15	14	20	22	13	100

Are these observations consistent with the hypothesis that the die is fair?

 **Solution**

Let θ_i be the probability that we get face i , and let

$$H_0 : \theta_1 = \theta_2 = \dots = \theta_6 = \frac{1}{6}$$

$$H_a : \text{at least one of the } \theta_i \neq \theta_j \text{ for } i \neq j, 0 \leq i, j \leq 6$$

Note that the above experiment is of a multinomial model. Let $\theta =$

$(\theta_1, \dots, \theta_6)$. Thus we have

$$L(\theta) = \frac{100!}{\prod_{i=1}^6 x_i!} \theta_1^{x_1} \theta_2^{x_2} \dots \theta_6^{x_6}.$$

With the constraint that $\sum_{i=1}^6 \theta_i = 1$, we can use Lagrange's Multiplier³ to find that

$$\hat{\theta}_i = \frac{x_i}{100}.$$

Then the likelihood ratio statistic is

$$\begin{aligned} \Lambda(\theta_0) &= -2 \ln \left[\frac{L(\theta_0)}{L(\hat{\theta})} \right] = -2 \ln \left[\frac{\prod_{i=1}^6 \left(\frac{1}{6}\right)^{x_i}}{\prod_{i=1}^6 \left(\frac{x_i}{100}\right)^{x_i}} \right] \\ &= 2 \sum_{i=1}^6 x_i \ln \frac{6x_i}{100} \xrightarrow{D} W \sim \chi^2(5) \end{aligned}$$

where $\hat{\theta} = (\hat{\theta}_1, \dots, \hat{\theta}_6)$, and the degree of freedom is because $\theta_6 = 1 - \sum_{i=1}^5 \theta_i$. The observed likelihood ratio statistic is

$$\begin{aligned} \lambda(\theta_0) &= 2 \sum_{i=1}^6 x_i \ln \frac{6x_i}{100} \\ &= -1.3063 - 3.1608 - 4.8819 + 7.2929 + 12.2158 - 6.4600 \\ &= 3.6997. \end{aligned}$$

And so the p -value, by

```
pchisq(3.6997, 5, lower.tail=FALSE)
```

is

$$p \approx P(W \geq 3.6997) \approx 0.5934.$$

This shows that we do not have sufficient evidence to reject H_0 , i.e. we have a fair die.

Example 23.1.5 (Example 7.5)

Suppose $X_1, \dots, X_n \sim N(\mu_0, \sigma^2)$ is a random sample. Test the hypothesis $H_0 : \sigma^2 = \sigma_0^2$, where the mean μ_0 is known.

³ A step of that can be found here: [MathSE](#).

Solution

Let $H_a : \sigma^2 \neq \sigma_0^2$. Since the X_i 's are IID, we have

$$\begin{aligned} L(\theta) &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(x_i - \mu_0)^2}{2\sigma^2}\right\} \\ &= (2\pi)^{-\frac{n}{2}} (\sigma^2)^{-\frac{n}{2}} \exp\left\{-\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu_0)^2\right\} \\ \ell(\sigma^2) &= -\frac{n}{2} \ln 2\pi - \frac{n}{2} \ln \sigma^2 - \frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu_0)^2 \\ \frac{d\ell}{d\sigma^2} &= -\frac{n}{2\sigma^2} + \frac{1}{2\sigma^4} \sum_{i=1}^n (x_i - \mu_0)^2 \\ \implies \hat{\sigma} &= \frac{1}{n} \sum_{i=1}^n (x_i - \mu_0)^2 \\ \frac{d^2\ell}{d(\sigma^2)^2} &= \frac{n}{2\sigma^4} - \frac{1}{\sigma^6} \sum_{i=1}^n (x_i - \mu_0)^2 \\ \frac{d^2\ell}{d(\sigma^2)^2} \Big|_{\sigma^2=\hat{\sigma}^2} &= \frac{n^3}{\left(\sum_{i=1}^n (x_i - \mu_0)^2\right)^2} \left(\frac{1}{2} - n^3\right) < 0 \end{aligned}$$

Then, under the null hypothesis, note that

$$\begin{aligned} R(\sigma^2) &= \frac{L(\sigma_0^2)}{L(\hat{\sigma}^2)} = \frac{(\sigma_0^2)^{-\frac{n}{2}}}{(\hat{\sigma}^2)^{-\frac{n}{2}}} \exp\left\{\left(\frac{1}{2\hat{\sigma}^2} - \frac{1}{2\sigma_0^2}\right) \sum_{i=1}^n (X_i - \mu_0)^2\right\} \\ &= \left(\frac{\hat{\sigma}^2}{\sigma_0^2}\right)^{\frac{n}{2}} \exp\left\{\frac{n}{2} \left(1 - \frac{\hat{\sigma}^2}{\sigma_0^2}\right)\right\} \end{aligned}$$

and thus

$$\begin{aligned} \Lambda(\sigma_0^2) &= -2 \ln R(\sigma_0^2) = -n \ln \left(\frac{\hat{\sigma}^2}{\sigma_0^2}\right) - n \left(1 - \frac{\hat{\sigma}^2}{\sigma_0^2}\right) \\ &= n \left[\frac{\hat{\sigma}^2}{\sigma_0^2} - 1 - \ln \left(\frac{\hat{\sigma}^2}{\sigma_0^2}\right)\right] \stackrel{D}{\rightarrow} W \sim \chi^2(1). \end{aligned}$$

The observed likelihood ratio statistic is therefore

$$\lambda(\sigma_0^2) = n \left[\frac{\hat{\sigma}^2}{\sigma_0^2} - 1 - \ln \left(\frac{\hat{\sigma}^2}{\sigma_0^2}\right)\right].$$

Then the p -value can be obtained by evaluating

$$p \approx P\left(W \geq n \left[\frac{\hat{\sigma}^2}{\sigma_0^2} - 1 - \ln \left(\frac{\hat{\sigma}^2}{\sigma_0^2}\right)\right]\right)$$

A Regularity Conditions

Throughout this course, the phrase “regularity conditions” have been regularly coined but never formally introduced. To clarify that, I have sought out for the instructor and confirmed that these are the regularity conditions that are used. This is taken out from *Casella & Berger (2002)*¹.

“(R)egularity conditions” are typically very technical, rather boring, and usually satisfied in most reasonable problems. But they are a necessary evil...

Furthermore,

These conditions mainly relate to differentiability of the density² and the ability to interchange differentiation and integration.

The following are the regularity conditions:

1. $X_1, \dots, X_n \sim f(x; \theta)$ are IID;
2. The unknown parameter is **identifiable**, i.e. if $\theta \neq \theta'$, then $f(x; \theta) \neq f(x; \theta')$;
3. The pf's $f(x; \theta)$ have common support, and $f(x; \theta)$ is differentiable in θ ;
4. The parameter space Ω contains an open set $\omega \subset \Omega$ of which the true parameter value θ_0 is an interior point³.
5. $\forall x \in \text{supp}(X)$, the pf $f(x; \theta)$ is three times differentiable with respect to θ , and the third derivative is continuous on θ , and $\int f(x; \theta)$ can be differentiated three times under the integral sign.
6. $\forall \theta_0 \in \Omega, \exists c \in \mathbb{R}_{>0}, \exists$ a function $M(x)$ ⁴ such that $\forall x \in \text{supp}(X)$,

$$|\theta - \theta_0| < c \implies \left| \frac{\partial^3}{\partial \theta^3} \ln f(x; \theta) \right| \leq M(x)$$

¹ Casella, G. & Berger, R. L. (2002). *Statistical Inference*. Thomson Learning

² probability function

³ ω is an open set, and so we can really only talk about interior points. The laymen definition of an interior point, is a point that is in a set but is not the limit point of the set.

⁴ Note that both c and $M(x)$ can depend on θ_0 , by the way that the statement is laid out.

B Useful References

B.1 Commonly Used Distributions

Distribution	pf	Mean	Var	mgf
Binomial Distribution : $X \sim \text{Bin}(n, p)$				
$x \in \mathbb{N} \cup \{0\}$				
$n \in \mathbb{N}$	$\binom{n}{x} p^x (1-p)^{n-x}$	np	$np(1-p)$	$(pe^t + 1 - p)^n$
$0 < p < 1$				
Geometric Distribution : $X \sim \text{Geo}(p)$				
$x \in \mathbb{N} \cup \{0\}$				
	$p(1-p)^x$	$\frac{1-p}{p}$	$\frac{1-p}{p^2}$	$\frac{p}{1-(1-p)e^t}$
$0 < p < 1$				$t < -\log(1-p)$
Poisson Distribution : $X \sim \text{Poi}(\mu)$				
$x \in \mathbb{N} \cup \{0\}$				
	$\frac{e^{-\mu} \mu^x}{x!}$	μ	μ	$e^{\mu(e^t-1)}$
$\mu > 0$				
Uniform Distribution : $X \sim \text{Unif}(a, b)$				
$a \leq x \leq b$				
	$\frac{1}{b-a}$	$\frac{a+b}{2}$	$\frac{(b-a)^2}{12}$	$\frac{e^{bt} - e^{at}}{(b-a)t}$
$a < b \in \mathbb{R}$				$t \neq 0$
Normal Distribution : $X \sim \text{N}(\mu, \sigma)$				
$x \in \mathbb{R}$				
	$\frac{1}{\sqrt{2\pi}\sigma^2} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$	μ	σ^2	$e^{\mu t + \frac{\sigma^2 t^2}{2}}$

$$\mu \in \mathbb{R}$$

$$\sigma > 0$$

Gamma Distribution : $X \sim \Gamma(\alpha, \beta)$

$$x \in \mathbb{R}_{\geq 0}$$

$$\alpha > 0$$

$$\beta > 0$$

$$\frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-\frac{x}{\beta}}$$

$$\alpha\beta$$

$$\alpha\beta^2$$

$$(1 - \beta t)^{-\alpha}$$

$$t < \frac{1}{\beta}$$

Exponential Distribution : $X \sim \text{Exp}(\theta)$

$$x \in \mathbb{R}_{> 0}$$

$$\theta > 0$$

$$\frac{1}{\theta} e^{-\frac{x}{\theta}}$$

$$\theta$$

$$\theta^2$$

$$(1 - \theta t)^{-1}$$

$$t < \frac{1}{\theta}$$

Negative Binomial Distribution : $X \sim \text{NB}(r, p)$

$$x \in \mathbb{N} \cup \{0\}$$

$$r > 0$$

$$p \in (0, 1)$$

$$\binom{x+r-1}{x} \cdot (1-p)^r p^x$$

$$\frac{pr}{1-p}$$

$$\frac{pr}{(1-p)^2}$$

$$\left(\frac{1-p}{1-pe^t}\right)^r$$

$$t < -\log p$$

Beta Distribution : $X \sim \text{Beta}(\alpha, \beta)$

$$x \in [0, 1] \text{ or}$$

$$x \in (0, 1)$$

$$\alpha > 0$$

$$\beta > 0$$

$$\frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)} x^{\alpha-1} (1-x)^{\beta-1}$$

$$\frac{\alpha}{\alpha+\beta}$$

$$\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$$

$$1 + \sum_{k=1}^{\infty} \left(\prod_{r=0}^{k-1} \frac{\alpha+r}{\alpha+\beta+r} \right) \frac{t^k}{k!}$$

Chi-Squared Distribution : $X \sim \chi^2(k)$

$$k \in \mathbb{N}$$

$$\text{if } k = 1$$

$$x \in (0, \infty)$$

otherwise

$$x \in [0, \infty]$$

$$\frac{1}{2^{\frac{k}{2}} \Gamma(\frac{k}{2})} x^{\frac{k}{2}-1} e^{-\frac{x}{2}}$$

$$k$$

$$2k$$

$$(1 - 2t)^{-\frac{k}{2}}$$

$$t < \frac{1}{2}$$

Bibliography

Casella, G. & Berger, R. L. (2002). *Statistical Inference*. Thomson Learning.

Geyer, C. J. (2007). Stat 5102 notes: Fisher information and confidence intervals using maximum likelihood. <https://www.stat.umn.edu/geyer/s06/5102/notes/fish.pdf>.

Index

- δ -Method, 155
- σ -algebra, 15
- σ -field, 15
- p -Value, 202

- Alternative Hypothesis, 201
- Asymptotic Confidence Intervals, 196
- Asymptotic Distribution, 135
- Asymptotic Distribution of Relative Likelihood, 183
- Asymptotic Normality, 183
- Asymptotic Pivotal Quantity, 188
- asymptotic variance, 183

- Bernoulli's Theorem, 143
- Beta Distribution, 88
- Binomial Distribution, 29
- Bivariate Normal Distribution, 114
- Bonferroni's Inequality, 19
- Boole's Inequality, 19

- Central Limit Theorem, 148
- Chebyshev's Inequality, 54
- Clairaut's Theorem, 71
- Cochran's Theorem, 132
- Conditional Distributions, 83
- Conditional Expectation, 101
- conditional mean, 101
- Conditional PF, 83
- Conditional Probability, 19
- conditional variance, 101
- Confidence Interval, 189
- Consistency, 182
- consistent estimator, 183
- Continuity Property, 20

- Continuous Mapping Theorem, 152
- Continuous Random Variable, 27
- Convergence in Distribution, 135
- Convergence in Probability, 139
- Correlation Coefficient, 99
- Covariance, 93
- Cumulative Distribution Function, 21

- Degenerate Distribution, 138
- Discrete Random Variable, 22
- Double Parameter Exponential Distribution, 144

- Estimate, 158
- Estimator, 158
- expected value, 44, 45
- Exponential Distribution, 31

- F Distribution, 132
- factorial moment, 51
- Factorization Theorem, 82
- Fisher Information, 164
- Fisher Information Matrix, 175

- Gamma Distribution, 31
- Gaussian Distribution, 30, 131
- Geometric Distribution, 29

- Hypothesis, 201

- implausible, 171
- Independence, 70, 78
- Independent Events, 19
- Indicator Function, 57
- Information Function, 163

- Information Matrix, 175
- Interval Estimate, 188
- Interval Estimators, 188
- Invariance Property of MLE, 169

- Jacobian, 121, 122
- Joint CDF, 63
- Joint Continuous Random Variables, 73
- joint density function, 74
- Joint Discrete Random Variables, 66
- Joint Expectation, 89
- Joint Moment Generating Functions, 107
- Joint Moments, 107
- Joint PMF, 67

- k-variate CDF, 111
- k-Variate Joint MGF, 112
- k-variate Support Set, 111
- Kolmogorov Axioms, 16

- Law of the Unconscious Statistician, 49, 89
- Law of Total Expectation, 104
- Law of Total Variance, 106
- Likelihood function, 159
- Likelihood Interval, 172, 198
- Likelihood Ratio Statistic, 204
- Likelihood Region, 172, 175
- Limiting Distribution, 135
- Linearity - Expectation, 49, 91
- Location Family, 41
- Location Parameter, 41
- Location-Scale Family, 42

Log Relative Likelihood, 172
 Log-likelihood, 159
 Lower Confidence Limit, 189

Marginal CDF, 66
 Marginal Distribution, 68
 Marginal MGF, 107
 Marginal Probability Density Function, 77
 Markov's Inequality, 52
 Markov's Inequality 2, 53
 maximum likelihood estimate, 159
 Measurable Space, 16
 Moment Generating Function, 56
 Moments, 51
 Multinomial Distribution, 113

Normal Distribution, 30
 Null Hypothesis, 201

observed information, 163
 observed information matrix, 175
 One-to-One Bivariate Transformations, 121
 One-to-One Transformation, 120

Pearson Correlation Coefficient, 99
 Pivotal Quantity, 188
 plausible, 171
 Poisson Distribution, 30
 power set, 15
 probability axioms, 16
 probability density function, 27
 Probability Integral Transformation, 40
 probability mass function, 22
 Probability Measure, 16
 probability set function, 16
 probability space, 16
 Product Rule, 87
 Properties of pdf, 27
 Properties of pmf, 22
 Properties of the cdf, 21

Random Sample, 115
 Random Variable, 20
 Relative Likelihood, 171
 right-continuous, 21

Sample Space, 15
 Scale Family, 42
 Scale Parameter, 42

Score Function, 163
 Score Vector, 175
 Significance Level, 202
 Simple Hypothesis, 204
 Slutsky's Theorem, 152
 Standard Normal Distribution, 31
 Statistic, 157
 support set, 22, 67, 74

t-test, 132
 Taylor Series with Lagrange's Remainder, 135
 Test of Hypothesis, 202
 Test Statistic, 202
 Type I Error, 203
 Type II Error, 203

unbiased estimator, 158
 uncorrelated, 93
 Uniform Distribution, 31
 Upper Confidence Limit, 189

Variance, 50

Weak Law of Large Numbers, 143